

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 September 2006 (08.09.2006)

PCT

(10) International Publication Number
WO 2006/093847 A1

(51) International Patent Classification:
C12N 15/11 (2006.01) *C12N 15/00* (2006.01)
C12N 15/06 (2006.01)

(21) International Application Number:
PCT/US2006/006752

(22) International Filing Date:
24 February 2006 (24.02.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

11/068,155	28 February 2005 (28.02.2005)	US
60/683,686	23 May 2005 (23.05.2005)	US
11/193,750	29 July 2005 (29.07.2005)	US
60/733,669	4 November 2005 (04.11.2005)	US
11/296,119	7 December 2005 (07.12.2005)	US

(71) Applicant (for all designated States except US): **AVIGENICS, INC.** [US/US]; Legal Department, 111 Riverbend Road, Athens, GA 30605 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **CHRISTMANN, Leandro** [US/US]; Watkinsville, GA 30677 (US). **EBERHARDT, Dawn, M.** [US/US]; Danielsville, GA 30633 (US). **LEAVITT, Markley, C.** [US/US]; Watkinsville, GA 30677 (US). **HARVEY, Alex, J.** [US/US]; Athens, GA 30606 (US).

(74) Agent: **YESLAND, Kyle**; AVIGENICS, INC., Legal Department, 111 Riverbend Road, Athens, GA 30605 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,

[Continued on next page]

(54) Title: ARTIFICIAL CHROMOSOMES AND TRANSCROMOSOMIC AVIANS



(57) Abstract: The invention includes avians containing an artificial chromosome in their genome and methods of making the avians.

WO 2006/093847 A1



SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*
~ *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ARTIFICIAL CHROMOSOMES AND TRANSCHROMOSOMIC AVIANS

This application claims priority to US patent application No. 11/296,119, filed
5 December 7, 2005; US provisional application No. 60/683,686, filed May 23, 2005;
US provisional application No. 60/733,669, filed November 4, 2005; US patent
application No. 11/193,750, filed July 29, 2005; and US patent application No.
11/068,155, filed February 28, 2005.

10

Field of the Invention

The present invention relates to the field of biotechnology, and more
specifically to the field of genome modification. Disclosed herein are compositions
including chromosomes and vectors, and methods of use thereof, for the generation of
15 genetically transformed cells and animals including avians.

Background

Transgenic technology to convert animals into "bioreactors" for the production
20 of specific proteins or other substances of pharmaceutical interest (Gordon et al, 1987,
Biotechnology 5: 1183-1187; Wilmut et al, 1990, Theriogenology 33: 113-123) offers
significant advantages over more conventional methods of protein production by gene
expression. For example, recombinant nucleic acid molecules have been engineered
and incorporated into transgenic animals so that an expressed heterologous protein
25 may be joined to a protein or peptide that allows secretion of the transgenic expression
product into milk or urine, from which the protein may then be recovered.

Another system useful for heterologous protein production is the avian
reproductive system. The production of an avian egg begins with formation of a large
yolk in the ovary of the hen. The unfertilized oocyte or ovum is positioned on top of
30 the yolk sac. After ovulation the ovum passes into the infundibulum of the oviduct
where it is fertilized, if sperm are present, and then moves into the magnum of the

oviduct which is lined with tubular gland cells. These cells secrete the egg-white proteins, including ovalbumin, lysozyme, ovomucoid, conalbumin and ovomucin into the lumen of the magnum where they are deposited onto the avian embryo and yolk. The hen oviduct offers outstanding potential as a protein bioreactor because of the
5 high levels of protein production, the promise of proper folding and post-translation modification of the target protein, the ease of product recovery, and the relatively short developmental period of chickens.

One method for creating permanent genomic modification of a eukaryotic cell is to integrate an introduced DNA into an existing chromosome. Retroviruses have so
10 far proven to be the method of choice for efficient integration. However, retroviral integration is directed to a number of insertion sites within the recipient genome so that positional variation in heterologous gene expression can be evident. Unpredictability as to which insertion site is targeted introduces an undesirable lack of control over the procedure. An additional limitation of the use of retroviruses is that
15 the size of the nucleic acid molecule encoding the virus and heterologous sequences may be limited to about 8 kb. In addition, retroviruses may include undesirable features such as splice sites. Although wild-type adeno-associated virus (AAV) often integrates at a specific region in the human genome, replication deficient vectors derived from AAV do not integrate site-specifically possibly due to the deletion of the
20 toxic rep gene. In addition, homologous recombination produces site-specific integration, but the frequency of such integration usually is typically low.

An alternative method for delivering a heterologous nucleic acid into the genome is the use of one or more site-specific enzymes that can catalyze the insertion of nucleic acids into chromosomes. These enzymes recognize relatively short unique
25 nucleic acid sequences that serve for both recognition and recombination. Examples include Cre (Sternberg & Hamilton, 1981, J. Mol. Biol. 150: 467-486, 1981), FIp (Broach et al, 1982, Cell 29: 227-234, 1982) and R (Matsuzaki et al, 1990, J. Bact. 172: 610-618, 1990).

A novel class of phage integrases, that includes the integrase from the phage phiC3 1, can mediate highly efficient integration of transgenes in mammalian cells both in vitro and in vivo (Thyagarajan et al, Mol. Cell Biol. 21: 3926-3934, 2001). Constructs and methods of using recombinase to integrate heterologous DNA into a plant, insect or mammalian genome are described by Calos in U.S. Patent Serial No. 6,632,672, the disclosure of which is incorporated in its entirety herein by reference.

The phiC3 1 integrase is a member of a subclass of integrases, termed serine recombinases, that include, for example, R4 and TP901-1. Unlike the phage lambda integrases, which belong to a tyrosine class of recombinases, the serine integrases do not require cofactors such as integration host factor. The phiC31 integrase normally mediates integration of the phiC31 bacteriophage into the genome of *Streptomyces* via recombination between the attP recognition sequence of the phage genome and the attB recognition sequence within the bacterial genome. When a plasmid is equipped with a single attB site, phiC3 1 integrase will detect and mediate crossover between the attB site and a pseudo-attP site within the mammalian genome. Such pseudo-attP integration sites have now been identified in the mouse and human genomes. If the heterologous DNA is in a circular or supercoiled form, the entire plasmid becomes integrated with attL and attR arms flanking the nucleic acid insert.

Integration mediated by certain integrases, such as PhiC3 1 integrase-mediated integration, results in the alteration of the recognition or recombination sites themselves so that the integration reaction is irreversible. This will bypass the primary concern inherent with other recombinases, i.e., the reversibility of the integration reaction and excision of the inserted DNA.

Another method for the stable introduction of heterologous nucleic acid (e.g., large heterologous nucleic acids) into a genome is by the use of an artificial chromosome. Artificial chromosomes for expression of heterologous genes in yeast are available, but artificial chromosomes being delivered to avians has not previously been achieved.

Therefore, it is an object of the invention to produce transgenic animals with large nucleic acid segments integrated into their genome and to provide avians which include an artificial chromosome in their genome.

In one useful embodiment, the transgenic avians of the invention can be used to produce polyclonal antibodies to antigens of interest for therapeutic applications. Fully human polyclonal antibodies have proven to be effective therapeutics, and in certain circumstances may be more efficacious than monoclonal antibodies. Polyclonal antibodies, opposed to monoclonal antibodies, are of particular therapeutic value for use against antigenic targets that are either complex in nature, subject to resistance via mutational escape or are highly polymorphic. For example, toxins can require multiple antibodies for effective neutralization. Also, pathogenic virus and bacteria, which can quickly mutate into resistant strains, are targets for polyclonal antibodies. In addition, polyclonal antibodies can be used as a masking therapeutic agent. For example, the polyclonal antibodies may be used in Rh disease therapy and immunosuppressive regimens associated transplant rejection and autoimmune disease.

At present, there are approximately 20 therapeutic polyclonals on the market. Existing polyclonal therapeutics are derived either from animal or human serum which imposes certain drawbacks. For example, polyclonal antibodies have a limited in vivo half-life. In addition, these polyclonals usually cannot be re-administered to a patient due to immune reaction. In addition, human serum derived antibodies, while fully human, have both inherent production limitations as well as certain bio-safety concerns.

Although human polyclonal antibodies have been produced in transgenic mice and cattle (reviewed in, for example, Bruggemann (2004) in *Molecular Biology of B cells* pp 547-561 . Academic Press and Kuroiwa et al (2002) *Nature Biotechnol* 20:889-894), there are certain limitations to each of these platforms with respect to large-scale manufacture of therapeutic polyclonals. For example, the levels of antibody production achievable in mice is extremely small by virtue of their body size. In cattle, the endogenous immunoglobulin genes are not "knocked-out", since embryonic stem cell lines necessary for knock-out procedures do not exist. Therefore,

contaminating bovine immunoglobulins will be present which will be difficult to separate from human antibodies by standard protein A/G affinity purification procedures. In addition, since the antibodies are produced in animal serum, there are biosafety and serum protein contamination problems.

5 In order to fully realize the potential of therapeutic polyclonals, a production platform is needed that can efficiently produce large quantities of fully human polyclonal antibody.

Transgenic chickens, which express fully human polyclonal antibodies in response to antigenic stimulation and deposit the antibodies into their eggs, would
10 present such an ideal production system. For example, a single hen has a production capacity of over 250 eggs/year and about 50 to about 100 mg of chicken IgG (also termed IgY) is naturally transported into each egg produced.

The present invention is also directed to methods of producing artificial chromosomes which contain large nucleic acid inserts, such as Ig loci. Producing
15 artificial chromosomes containing a transgene by integrating the transgene into the chromosome can have certain limitations. For example, in some integration methodologies the transgene can integrate into any of the available chromosomes within the cell, including the host cells chromosomes. In certain instances homologous recombination, can overcome this problem. However, homologous
20 recombination has a number of limitations including the requirement that the transgene be specifically engineered for the procedure. In certain useful site specific integration methodologies, the transfected nucleotide sequence must be circular, otherwise integration will introduce a double-stranded break into the artificial chromosome. To by-pass the need for a circular insert the vector can be equipped with two RRSs that
25 flank the desired transgene. However, two recombinases would be required for the integration event and the artificial chromosome would also harbor two recombination sites. The complexity involved in this type of integration would result in an overall low rate of integration. Regardless of the integration methodology employed, the efficiency of integration for large transgenes is typically very much reduced relative to
30 the integration of smaller transgenes, (e.g., up to 1000 fold reduction in efficiency for

transgenes over 80 kb (kilobases) relative to smaller transgenes, for example, less than 10 kb). This may be due to certain factors such as the large size of the transgene lowering the rate of transfection. In addition, large transgenes can be susceptible to nicking and breaking due to shear forces and/or nuclease degradation.

5 One potential difficulty in the use of artificial chromosomes in the production of transchromosomic animals such as avians can be difficulty in preparing a sufficiently homogeneous mixture of artificial chromosomes. Fluorescent synthetic polyamide probes have been used to obtain human chromosomes from their native environment by tagging repeated sequences in the chromosome with labeled
10 polyamides (see, for example, Gygi et al. (2002) Use of fluorescent sequence-specific polyamides to discriminate human chromosomes by microscopy and flow cytometry. Nucleic Acids Res 30: 2790-9).

Purification of artificial chromosomes by methods such as flow cytometry can be limited to only metaphase chromosomes, for example, larger than 60Mb in size. In
15 certain instances artificial chromosomes which cannot be purified using conventional technologies (e.g., artificial chromosomes less than about 60Mb in size) could be useful for the production of transchromosomic animals including transchromosomic avians.

What is needed are methods which provide for the efficient introduction of
20 artificial chromosomes into animal genomes such as avian genomes.

Summary of the Invention

One useful aspect of the invention relates to methods of producing transchromosomic avians. In one embodiment, the methods include substantially
25 purifying a chromosome followed by introducing the purified chromosome into an avian embryo and thereafter maintaining the embryo under conditions suitable for the embryo to develop and hatch as a chick. In one embodiment, the methods include inserting a heterologous nucleotide sequence into the chromosome before or after substantially purifying the chromosome. In one embodiment, the chromosome is
30 introduced into the avian embryo by microinjection; however, any useful method to

introduce the chromosome into the avian embryo is within the scope of the present invention.

It is contemplated that the chromosome may be introduced into the embryo by delivering the chromosome to an avian cell before or after fertilization. For example, the chromosome may be introduced into an ovum or a sperm before fertilization. In another example, the chromosome is introduced into a cell of an embryo (e.g., stage I to stage XII embryo). In one embodiment, the chromosome is introduced into an early stage embryo, for example, and without limitation, a stage I embryo. In one embodiment, the chromosome is introduced into a germinal disc.

The methods provide for the introduction of any useful number of chromosomes into the avian embryo in order to produce a transchromosomal avian. For example, and without limitation, between 1 and about 10,000 chromosomes may be introduced into the embryo. In another example, between 1 and about 1,000 chromosomes may be introduced into the embryo.

The invention also provides for transchromosomal avian cells wherein the artificial chromosome includes a nucleotide sequence which encodes a therapeutic substance. The cells may be isolated from transchromosomal avians and thereafter grown in culture. The invention also contemplates the production of the transchromosomic avian cells by stable introduction of the artificial chromosome into cultured avian cells. Any useful method may be employed for the introduction of the artificial chromosome into the cultured cells including, without limitation, lipofection or microinjection.

The invention also contemplates methods which include isolating an artificial chromosome; introducing the artificial chromosome into an avian embryo; maintaining the embryo under conditions suitable for the embryo to develop and hatch as a chick; and maintaining the chick under conditions suitable to obtain a mature avian wherein the artificial chromosome is present in the genome of the mature avian.

In one aspect, the invention relates to methods which include isolating an artificial chromosomes by flow cytometry. The flow cytometry may be facilitated by a probe which is associated with the artificial chromosome. For example, the probe may

be a polyamide probe. In one embodiment, the probe (e.g., polyamide probe) may include a fluorescent molecule or tag.

In one embodiment, the artificial chromosome is present in micronuclei. For example, the artificial chromosome may be present in micronuclei prior to flow
5 cytometry purification of the artificial chromosome. In one embodiment, the micronuclear environment protects the artificial chromosome from degradation or fragmentation that may occur before during or after introduction (e.g., by injection or lipofection) of the artificial chromosome into the avian embryo. In one embodiment, the micronuclei contain diploid mitotic artificial chromosomes. Production of
10 micronuclei can be accomplished by any useful method known in the art, for example, as disclosed in Labidi et al (1987) *Experimental Cell Research* 617-627, the disclosure of which is incorporated in its entirety herein by reference.

Typically the methods include transferring the embryo, into which the artificial chromosome has been introduced, into a recipient female avian. In one embodiment,
15 the artificial chromosome is an early stage embryo such as a stage I, stage II, stage III, stage IV, stage V or Stage VI embryo. In one useful embodiment, the embryo is a stage I embryo.

In one useful embodiment, the artificial chromosome comprises one or more heterologous recombination sites, for example, between 1 and about 100
20 recombination sites may be employed. Typically, the chromosome will include a heterologous coding sequence. In one useful embodiment, the heterologous coding sequence consists of or contains a pharmaceutical protein coding sequence. Any useful pharmaceutical protein coding sequence may be employed, such as those disclosed elsewhere herein. In addition, the artificial chromosome can include a
25 promoter, for example, and without limitation, a promoter which functions in tubular gland cells. For example, the promoter may be linked to a pharmaceutical protein coding sequence such that the promoter initiates transcription of the heterologous coding sequence (i.e., the promoter is operably linked to the heterologous coding sequence). The invention also contemplates the inclusion of an IRES (internal
30 ribosome entry site) in the artificial chromosome.

In one aspect, the invention provides for transgenic avians which produce eggs containing polyclonal antibodies, for example, human polyclonal antibodies. The invention also relates to the eggs produced by such an avian. The avians employed in the invention may be any useful avians, such as those avians disclosed elsewhere
5 herein, for example chickens, quail and turkeys. The invention contemplates the production of chimeric birds and germline transgenic birds including G1 and G2 transgenic or transchromosomic avians which produce polyclonal antibodies.

In one useful embodiment of the invention, one or more cells of the transgenic avian contain an artificial chromosome which has coding sequences for a polyclonal
10 antibody. Any useful artificial chromosome may be employed such as those having a centromere selected from the group consisting of an insect centromere, a mammalian centromere and an avian centromere. In one specific embodiment, the artificial chromosome is a satellite artificial chromosome.

The invention also provides for methods of producing artificial chromosomes
15 in cells. In one aspect, methods of the invention include introducing one or more transgenes into an artificial chromosome during assembly of the artificial chromosome. In one useful embodiment, the transgenes contain at least one of a promoter and a coding sequence for a therapeutic protein. In one embodiment, the coding sequence encodes one or more Ig loci such as Ig λ , IgK₅ IgH, or portions thereof
20 or combinations thereof in its germline. The methods for producing artificial chromosomes containing a transgene are particularly useful for the introduction of large transgenes into the chromosome such as portions of Ig genes, for example, portions of human Ig genes (c.g., an Ig λ gene, an Ig H gene and/or an IgK gene). Certain references which include disclosure that can be useful in certain aspects of the
25 invention include Csonka, et al (2000) Journal of Cell Science 113: 3207-3216 and Nicholson, et al (1999) J. Immunology 163(12):6898-6906. The disclosures of each of these two journal articles are incorporated in their entirety herein by reference.

Integration of a transgene into a defined chromosomal site is useful to improve the predictability of expression of the transgene, which is particularly advantageous
30 when creating transgenic vertebrate animals such as, transgenic avians. Transgenesis

by methods that randomly insert a transgene into a genome are often inefficient since the transgene may not be expressed at the desired levels or in desired tissues.

The present invention relates to methods of modifying the genome of vertebrate cells (e.g., production of transgenic vertebrates, in particular, transgenic avians) and to such cells with modified genomes and their progeny. In one embodiment, the methods provide for introducing into vertebrate cells a first recombination site such that the recombination site is inserted into the vertebrate cell genome. Typically, in such embodiments, the genome does not normally include this first recombination site prior to the recombination site introduction. Methods of the invention may also include introducing a nucleotide sequence comprising a second recombination site and a sequence of interest such as a coding sequence into the vertebrate cell or progeny of the vertebrate cell. The nucleotide sequence comprising the second recombination site and the sequence of interest such as a coding sequence may be introduced into the vertebrate cell before, at about the same time as or after the introduction of the first recombination site. Additionally, the present methods may include introducing into the vertebrate cell or progeny cell thereof a substance which facilitates insertion of the nucleotide sequence comprising the second recombination site and the sequence of interest proximal to the first recombination site. For example, the nucleotide sequence comprising the second recombination site and the sequence of interest may be inserted adjacent to or internally in the first recombination site. In one very useful embodiment, the first recombination site and/or the nucleotide sequence comprising the second recombination site and the sequence of interest are stably incorporated into the genome of the cell.

The present invention contemplates the genomic modification of any useful vertebrate cells including, but not limited to, avian cells. Examples of cells which may have their genomes modified in accordance with the present invention include, without limitation, reproductive cells including sperm, ova and embryo cells and nonreproductive cells such as tubular gland cells.

The present invention also relates to methods of producing transgenic vertebrate animals and to the transgenic animals produced by the methods and to their

transgenic progeny or descendants. The invention also includes the transgenic cells included in or produced by the transgenic vertebrate animals. Examples of such cells include, without limitation, germline cells, ova, sperm cells and protein producing cells such as tubular gland cells. In one useful embodiment, the transgenic vertebrate animals of the invention are transgenic avians. Transgenic avians of the invention may include, without limitation, chickens, turkeys, ducks, geese, quail, pheasants, parrots, finches, hawks, crows or ratites including ostrich, emu or cassowary.

In accordance with the present invention, methods of producing transgenic vertebrate animals can include introducing into an embryo of a vertebrate animal a first recombination site such that the recombination site is present in sperm or ova of a mature vertebrate animal developed from the embryo. In one useful embodiment, the embryo does not normally include the first recombination site in its genome prior to the recombination site introduction. The methods may also include introducing a nucleotide sequence comprising a second recombination site and a sequence of interest such as a coding sequence into the embryo of the vertebrate animal. The first recombination site and/or the nucleotide sequence comprising the second recombination site and a sequence of interest may be introduced into the embryo of the vertebrate animal before the embryo is fertilized (i.e., when an ovum), at about the same time as introduction of the sperm into the ovum or after fertilization.

The methods can also include introducing the nucleotide sequence comprising a second recombination site and a sequence of interest into an ovum or a sperm of a mature vertebrate animal developed from the embryo (or its descendants) into which the first recombination site was introduced. In one embodiment, the nucleotide sequence comprising a second recombination site and a sequence of interest is introduced into the ovum from the mature vertebrate animal before the ovum is fertilized. In another embodiment, the nucleotide sequence comprising a second recombination site and a sequence of interest is introduced into the ovum at about the time of fertilization. In one particularly useful embodiment, the nucleotide sequence comprising a second recombination site and a sequence of interest is introduced into the ovum after the ovum is fertilized (when an embryo).

The methods may include, upon addition of the nucleotide sequence comprising a second recombination site and a sequence of interest to an embryo, ovum or sperm, introducing into the embryo, ovum or sperm, a substance which facilitates insertion of the nucleotide sequence comprising the second recombination site and the sequence of interest proximal to the first recombination site. For example, the nucleotide sequence comprising the second recombination site and the sequence of interest may be inserted adjacent to or internally in the first recombination site. In one useful embodiment, the methods include introducing into an embryo comprising the first recombination site in its genome, a substance which facilitates insertion of the nucleotide sequence comprising the second recombination site and the sequence of interest proximal to the first recombination site.

In one useful embodiment, these methods include fertilizing an ovum with sperm comprising the first recombination site. The methods can include also introducing into the ovum a nucleotide sequence comprising a second recombination site and a sequence of interest such as a coding sequence and a substance which facilitates insertion of the nucleotide sequence comprising the second recombination site and sequence of interest proximal to (e.g., adjacent to or internally in) the first recombination site. It is contemplated that the nucleotide sequence comprising a second recombination site and a sequence of interest may be introduced into the ovum before or after fertilization by the sperm or at about the same time as fertilization.

In one very useful embodiment of the methods disclosed herein, the nucleotide sequence comprising the second recombination site and the sequence of interest is stably incorporated into the genome of the embryo, ovum or sperm.

The methods disclosed herein typically eventually include exposing a fertilized ovum to conditions which lead to the development of a viable transgenic vertebrate animal.

In one embodiment, the nucleotide sequence of interest includes an expression cassette. Optionally, the nucleotide sequence of interest may include a marker such as, but not limited to, a puromycin resistance gene, a luciferase gene, EGFP-encoding gene, and the like.

Typically, in accordance with methods known in the art or methods disclosed herein, the embryo of the vertebrate animal or fertilized ovum of a mature vertebrate animal of the invention is exposed to conditions which lead to the development of a viable transgenic vertebrate animal.

5 Embryos that are useful in the present methods include, without limitation, stage I, stage II, stage III, stage IV, stage V, stage VI, stage VII, stage VIII, stage IX, stage X, stage XI and stage XII embryos.

In one embodiment, the nucleotide sequence included with the second recombination site of interest is a coding sequence. The nucleotide sequence of
10 interest included with the second recombination site can be of any useful size. For example, and without limitation, the nucleotide sequence of interest may be from about 0.1 kb to about 10 mb, for example, about 1 kb to about 1 mb. In one embodiment, the nucleotide sequence of interest is about 5 kb to about 5 mb in size, for example, about 5 kb to about 2 mb, e.g., about 8 kb to about 1 mb. In one
15 embodiment, the nucleotide sequence of interest is about 0.5 kb to about 500 kb.

The first recombination site and/or the nucleotide sequence which includes the second recombination site and a sequence of interest such as a coding sequence may be introduced into cells, embryos (i.e., fertilized ova) or sperm by any useful method. These useful methods include, without limitation, cell fusion, lipofection, transfection,
20 microinjection, calcium phosphate co-precipitation, electroporation, protoplast fusion, particle bombardment and the like. In addition, the first recombination site or nucleotide sequence comprising the second recombination site and the sequence of interest may be introduced into cells, embryos, ova or sperm in the presence of a cationic polymer such as PEI and/or other substances disclosed elsewhere herein or
25 known in the art.

In one embodiment, recombination sites employed in the present invention are isolated from bacteriophage and/or bacteria. For example, the recombination sites may be attP sites or attB sites.

The substance which facilitates insertion of the second recombination site and
30 a sequence of interest may be an enzyme. In one embodiment, the substance is a site

specific recombinase. In one useful embodiment, the substance which facilitates insertion of the nucleotide sequence is nucleic acid, for example, DNA or RNA. The DNA or RNA may include modified nucleosides as described elsewhere herein or are known to those of skill in the art. In one embodiment, modified nucleosides are employed to extend the half-life of RNA or DNA molecules employed in the present invention. For example, it may be desirable to extend the half life of the RNA or DNA molecules in the presence of a cellular environment. In one useful embodiment, the nucleic acid encodes an enzyme such as a site specific recombinase.

Nonlimiting examples of site specific recombinases which may be employed herein either as protein or encoded by nucleic acid include serine recombinases and tyrosine recombinases. Examples of serine recombinases which may be employed include, without limitation, EcoYBCK, Φ C31, SCH10.38c, SCC88.14, SC8F4.15c, SCD12A.23, Bxb1, WwK, Sau CcrB, Bsu CisB, TP901-1, Φ 370.1, Φ 105, Φ FC1, A118, CaeI956, CaeI951, Sau CcrA, Spn, TnpX, TndX, SPBc2, SC3C8.24, SC2E1.37, SCD78.04c, R4, Φ Rv1, Y4bA and Bja serine recombinases.

In one embodiment of the invention, the present methods include introducing an integration host factor into a cell (e.g., an embryo) to facilitate genomic integration. Such integration host factors may be particularly useful when employing certain substances such as tyrosine recombinases as disclosed herein.

The nucleotide sequence of interest may include a coding sequence. The coding sequence may encode any useful protein. In one useful embodiment, the sequence of interest encodes a pharmaceutical or therapeutic substance. The invention contemplates the production of any useful protein based pharmaceutical or therapeutic substance. Examples of pharmaceutical or therapeutic substances include without limitation at least one of a light chain or a heavy chain of an antibody (e.g., a human antibody) or a cytokine. In one embodiment, the pharmaceutical or therapeutic composition is interferon, erythropoietin, or granulocyte-colony stimulating factor, in one embodiment, the transgenic animal is an avian and the sequence of interest encodes a polypeptide present in eggs produced by the avian.

In one embodiment, integrases such as phage integrases, for example, serine recombinases, such as the integrase from phage phiC31, can mediate the efficient integration of transgenes into target cells both in vitro and in vivo. In one embodiment, when a plasmid is equipped with a single attB site, the integrase detects
5 attP homologous sequences, termed pseudo-attP sites, in a target genome and mediates crossover between the attB site and a pseudo attP site.

In one embodiment, once delivered to a recipient cell, for example, an avian cell, the phiC31 integrase mediates recombination between the att site within the nucleic acid molecule and a bacteriophage attachment site within the genomic DNA of
10 the cell. Both att sites are disrupted and the nucleic acid molecule, with partial att sequences at each end, is stably integrated into the genome attP site. The phiC31 integrase, by disrupting the att sites of the incoming nucleic acid and of the recipient site within the cell genome can preclude any subsequent reverse recombination event that would excise the integrated nucleic acid and reduce the overall efficiency of stable
15 incorporation of the heterologous nucleic acid.

Following delivery of the nucleic acid molecule and a source of integrase activity into a cell population and integrase-mediated recombination, the cells may be returned to an embryo, hi the case of avians, late stage blastodermal cells may be returned to a hard shell egg, which is resealed for incubation until hatching. Stage I
20 embryos may be directly microinjected with the polynucleotide and source of integrase activity, isolated, transfected and returned to a stage I embryo which is reimplanted into a hen for further development. Additionally, the transfected cells may be maintained in culture in vitro.

The present invention provides novel methods and recombinant polynucleotide
25 molecules for transfecting and integrating a heterologous nucleic acid molecule into the genome of a cell of a vertebrate animal, such as an avian. Certain methods of the invention provide for the delivery to a cell population a first nucleic acid molecule that comprises a region encoding a recombination site, such as a bacterial recombination site or a bacteriophage recombination site. In one embodiment, a source of integrase
30 activity is also delivered to the cell and can be in the form of an integrase-encoding

nucleic acid sequence and its associated promoter or as a region of a second nucleic acid molecule that may be co-delivered with the polynucleotide molecule. Alternatively, integrase protein itself can be delivered directly to the target cell.

5 The recombinant nucleic acid molecules of the present invention may further comprise a heterologous nucleotide sequence operably linked to a promoter so that the heterologous nucleotide sequence, when integrated into the genomic DNA of a recipient cell, can be expressed to yield a desired polypeptide. The nucleic acid molecule may also include a second transcription initiation site, such as an internal ribosome entry site (IRES), operably linked to a second heterologous polypeptide-
10 encoding region desired to be expressed with the first polypeptide in the same cell.

The present invention provides modified isolated artificial chromosomes useful as vectors to shuttle transgenes or gene clusters into a genome of an avian. By delivery of the modified chromosome to a recipient cell, the target cell, and progeny thereof, become trisomic or transchromosomic. The additional chromosome will
15 typically not affect the subsequent development of the recipient cell and/or embryo, nor interfere with the reproductive capacity of an adult bird developed from such cells or embryos. The chromosome will also be stable within the genome of the cells of the adult bird or within isolated avian cells. The invention provides methods to isolate a population of chromosomes for delivery into embryos or early cells of avians, for
20 example, chickens.

The methods can include inserting a lac-operator sequence into an isolated chromosome and, optionally, inserting a desired transgene sequence within the same chromosome. The lac operator region is typically a concatamer of a plurality of lac operators for the binding of multiple lac repressor molecules. A recombinant DNA
25 molecule is constructed that includes an identified region of the target chromosome, a recombination site such as attB or attP, and the lac-operator concatamer. The recombinant molecule is delivered to an avian cell, and homologous recombination will integrate the heterologous polynucleotide and the lac-operator concatamer into the targeted chromosome. A tag-polypeptide, such as the GFP-lac-repressor fusion
30 protein, binds to the lac-operator sequence for identification and isolation of the

genetically modified chromosome. The tagged mitotic chromosome can be isolated using, for instance, flow cytometry.

Among other things, the present invention relates to transchromosomic avians. In a particular aspect, the invention provides for GO founder transchromosomic avians (e.g., chimeric including, but not limited to, germline chimeric transchromosomic avians) which can give rise to germline transchromosomic offspring, for example, G1 and G2 germline transchromosomic offspring.

Examples of avians which are contemplated for use herein include, without limitation, chicken, turkey, duck, goose, quail, pheasants, parrots, finches, hawks, crows and ratites including ostrich, emu and cassowary.

In one useful aspect, the artificial chromosome employed herein includes a centromere. Any useful centromere may be employed in the present invention including, without limitation, centromeres from insects, mammals or avians.

In one particularly useful embodiment, the artificial chromosomes used herein include a heterologous nucleotide sequence. The nucleotide sequence may be heterologous to the avian and/or heterologous to the artificial chromosome. In one useful embodiment, the heterologous nucleotide sequence includes a coding sequence for a therapeutic substance. In addition, the heterologous nucleotide sequence may include a gene expression controlling region. Any useful gene expression controlling region may be employed in the invention. For example, and without limitation, the gene expression controlling region may include a lysozyme promoter, an ovomucin promoter, a conalbumin promoter, an ovomucoid promoter and/or an ovalbumin promoter or functional portions thereof. See, for example, US Patent Application No. 10/114,739, filed April 1, 2002; US Patent Application No. 10/856,218, filed May 28, 2004 and US Patent Application No. 10/733,042, filed December 11, 2003. The disclosure of each of these patent applications is incorporated herein by reference in its entirety. In one useful embodiment, the product of the heterologous nucleotide sequence (e.g., therapeutic substance) is delivered to the avian egg (e.g., the egg white) during production of the egg in the avian. The invention also includes the eggs

produced by the avians produced by these methods and other methods disclosed herein.

Another aspect of the present invention is a cell, for example, an avian cell, genetically modified with a transgene vector by the methods of the invention. For example, in one embodiment, the transformed cell can be a chicken early stage blastodermal cell or a genetically transformed cell line, including a sustainable cell line. The transfected cell may comprise a transgene stably integrated into the nuclear genome of the recipient cell, thereby replicating with the cell so that each progeny cell receives a copy of the transfected nucleic acid. One useful cell line for the delivery and integration of a transgene comprises a heterologous attP site that can increase the efficiency of integration of a polynucleotide by an integrase, such as phiC3 1 integrase and, optionally, a region for expressing the integrase.

Another aspect of the present invention is methods of expressing a heterologous polypeptide in a cell by stably transfecting a cell by using site-specific integrase-mediation and a recombinant nucleic acid molecule, as described above, and culturing the transfected cell under conditions suitable for expression of the heterologous polypeptide under the control of a transcriptional regulatory region.

Yet another aspect of the present invention concerns transgenic vertebrate animals, such as birds, for example chickens, comprising a recombinant nucleic acid molecule and which may (though optionally) express a heterologous gene in one or more cells in the animal. For example, in the case of avians, embodiments of the methods for the production of a heterologous polypeptide by the avian tissue involve providing a suitable vector and introducing the vector into embryonic blastodermal cells containing an attP site together with an integrase, for example, a serine recombinase such as phiC3 1 integrase, so that the vector can integrate into the avian genome at the attP site which has been engineered into the cell genome. A subsequent step may involve deriving a mature transgenic avian from the transgenic blastodermal cells by transferring the transgenic blastodermal cells to an embryo, such as a stage X embryo (e.g., an irradiated stage X embryo), and allowing that embryo to develop fully, so that the cells become incorporated into the bird as the embryo is allowed to

develop. In one embodiment, sperm from a GO bird positive for the transgene is used to inseminate a chicken giving rise to a fully transgenic G1 generation.

One approach may be to transfer a transfected nucleus to an enucleated recipient cell which may then develop into a zygote and ultimately an adult animal.

5 The resulting animal is then grown to maturity.

In the transgenic vertebrate of the present invention, the expression of the transgene may be restricted to specific subsets of cells, tissues or developmental stages utilizing, for example, trans-acting factors acting on the transcriptional regulatory region operably linked to the polypeptide-encoding region of interest of the present invention and which control gene expression in the desired pattern. Tissue-specific regulatory sequences and conditional regulatory sequences can be used to control expression of the transgene in certain spatial patterns. Moreover, temporal patterns of expression can be provided by, for example, conditional recombination systems or prokaryotic transcriptional regulatory sequences. By inserting an integration site such as attP into the genome, it is believed that expression of an integrated coding sequence will be much more predictable.

The invention can be used to express, in large yields and at low cost, a wide range of desired proteins including those used as human and animal pharmaceuticals, diagnostics, and livestock feed additives. Proteins such as growth hormones, cytokines, structural proteins and enzymes including human growth hormone, interferon, lysozyme, and β -casein may be produced by the present methods. In one embodiment, proteins are expressed in the oviduct and deposited in eggs of avians, such as chickens, according to the invention. The present invention includes these eggs and these proteins.

25 The present invention also includes methods of producing transgenic vertebrate animals, for example, transgenic chickens, which employ the use of integrase, cationic polymers and/ nuclear localization signals. The present invention also includes the transgenic vertebrate animals, such as the avians, produced by these methods and other methods disclosed herein. The invention also includes the eggs produced by the transgenic avians produced by these methods and other methods disclosed herein.

In one embodiment, the methods of the invention include introducing into a cell: 1) a nucleic acid comprising a transgene; 2) an integrase activity; and 3) a cationic polymer. Such methods provide for an increased efficiency of transgenic avian production relative to identical methods without the cationic polymer.

5 In another embodiment, the methods include introducing into a cell: 1) a nucleic acid comprising a transgene; 2) an integrase activity; and 3) a nuclear localization signal. Such methods provide for an increased efficiency of transgenic animal, for example, avian, production relative to identical methods without the nuclear localization signal.

10 In another embodiment, the methods include introducing into a cell: 1) a nucleic acid comprising a transgene; 2) an integrase activity; 3) a cationic polymer; and 4) a nuclear localization signal. Such methods provide for an increased efficiency of transgenic vertebrate animal production relative to identical methods without the cationic polymer or without the nuclear localization signal.

15 In one embodiment, the cell is a cell of an embryo, for example, an avian embryo. In one embodiment, the cell is a cell of an early stage avian embryo comprising a germinal disc. The avian cell may be, for example, a cell of a stage I avian embryo, a cell of a stage II avian embryo, a cell of a stage III avian embryo, a cell of a stage IV avian embryo, a cell of a stage V avian embryo, a cell of a stage VI
20 avian embryo, a cell of a stage VII avian embryo, a cell of a stage VIII avian embryo, a cell of a stage IX avian embryo, a cell of a stage X avian embryo, a cell of a stage XI avian embryo or a cell of a stage XII avian embryo. In one particularly useful embodiment, the avian cell is a cell of a stage X avian embryo. In another useful embodiment, the avian cell is a cell of a stage I avian embryo.

25 The methods provide for the introduction of nucleic acid into the avian cell by any suitable technique known to those of skill in the art. For example, the nucleic acid may be introduced into the avian cell by microinjecting, transfection, electroporation or lipofection. In one particularly useful embodiment, the introduction of the nucleic acid is accomplished by microinjecting.

The nucleic acid which includes a transgene may be DNA or RNA or a combination of RNA and DNA. The nucleic acid may comprise a single strand or may comprise a double strand. The nucleic acid may be a linear nucleic acid or may be an open or closed circular nucleic acid and may be naturally occurring or synthetic.

5 Integrase activity may be introduced into the cell, such as an avian cell, in any suitable form. In one embodiment, an integrase protein is introduced into the cell. In another embodiment, a nucleic acid encoding an integrase is introduced into the cell. The nucleic acid encoding the integrase may be double stranded DNA, single stranded DNA, double stranded RNA, single stranded RNA or a single or double stranded
10 nucleic acid which includes both RNA and DNA. In one particularly useful embodiment, the nucleic acid is mRNA. Integrase activity may be introduced into the cell by any suitable technique. Suitable techniques include those described herein for introducing the nucleic acid encoding a transgene into a cell. In one useful embodiment, the integrase activity is introduced into the cell with the nucleic acid
15 encoding the transgene. For example, the integrase activity may be introduced into the cell in a mixture with the nucleic acid encoding the transgene.

 In one embodiment, a nuclear localization signal (NLS) is associated with the nucleic acid which includes a transgene. For example, the NLS may be associated with the nucleic acid by a chemical bond. Examples of chemical bonds by which an
20 NLS may be associated with the nucleic acid include an ionic bond, a covalent bond, hydrogen bond and Van der Waal's force. In one particularly useful embodiment, the nucleic acid which includes a transgene is associated with an NLS by an ionic bond. NLS may be introduced into the cell by any suitable technique. Suitable techniques included those described herein for introducing the nucleic acid encoding a transgene
25 into a cell. In one useful embodiment, the NLS is introduced into the cell with the nucleic acid encoding the transgene. For example, the NLS may be introduced into the cell while associated with the nucleic acid encoding the transgene.

 Cationic polymers may be employed to facilitate the production of transgenic vertebrate animals such as avians. For example, the cationic polymers may be
30 employed in combination with integrase and/or NLS. Any suitable cationic polymer

may be used. For example, and without limitation, one or more of polyethylenimine, polylysine, DEAE-dextran, starburst dendrimers and starburst polyamidoamine dendrimers may be used. In a particularly useful embodiment, the cationic polymer includes polyethylenimine. The cationic polymer may be introduced into the cell by
5 any suitable technique. Suitable techniques included those described herein for introducing the nucleic acid encoding a transgene into a cell. In one useful embodiment, the cationic polymer is introduced into the cell in a mixture with the nucleic acid encoding the transgene. For example, the cationic polymer may be introduced into the avian cell while associated with the nucleic acid encoding the
10 transgene.

In one particularly useful embodiment of the invention, the transgene includes a coding sequence which is expressed in a cell of the transgenic vertebrate animal, for example, a transgenic avian, producing a peptide or a polypeptide (e.g., a protein). The coding sequence may be expressed in any or all of the cells of the transgenic
15 animal. For example, the coding sequence may be expressed in the blood, the magnum and/or the sperm of the animal. In a particularly useful embodiment of the invention, the polypeptide is present in an egg, for example, in the egg white, produce by a transgenic avian.

The present invention also includes methods of dispersing nucleic acid in a
20 cell, for example, in an avian cell (e.g., an avian embryo cell). For example, the nucleic acid may be dispersed in the cytoplasm of a cell. These methods include introducing into a cell a nucleic acid and a dispersing agent, for example, a cationic polymer (e.g., polyethylenimine, polylysine, DEAE-dextran, starburst dendrimers and/or starburst polyamidoamine dendrimers) in an amount that will disperse the
25 nucleic acid in a cell. Typically, the dispersing of the nucleic acid is a homogeneous dispersing. In one embodiment, the dispersed nucleic acid includes a transgene. NLS or integrase activity may also be introduced into the cell. Dispersing of the nucleic acid may be particularly useful when the DNA is introduced into a cell containing a relatively large volume of cytoplasm, such as an avian embryo cell or a germinal disc.
30 Dispersing of the nucleic acid in the cell can increase the likelihood that the nucleic

acid will contact and enter the nucleus of the cell into which the nucleic acid has been introduced. Without such dispersing, the nucleic acid may localize to one or more areas within the cell and may not contact the nucleus of the cell. In addition, where the quantity of nucleic acid introduced into the cell is known, dispersing of the nucleic acid can assist in exposing the nucleus in the cell to known or specific concentrations of the nucleic acid.

The methods of the invention include introducing the cell into a recipient animal, for example, an avian such as a chicken, wherein the recipient avian produces an offspring which includes the transgene. The cell may be introduced into a recipient animal by any suitable technique.

The present invention also includes the identification of certain regions in the genome which are advantageous for heterologous gene expression. These regions can be identified by analysis, using methods known in the art, of the transgenic vertebrate animals or cells produced as disclosed herein.

The production of vertebrate animals or avians which are the mature animals developed from the recombinant embryos, ovum and/or sperm of the invention typically are referred to as the G0 generation and are usually hemizygous for each inserted transgene. The G0 generation may be bred to non-transgenic animals to give rise to G1 transgenic offspring which are also hemizygous for the transgene. The G1 hemizygous offspring may be bred to non-transgenic animals giving rise to G2 hemizygous offspring or may be bred together to give rise to G2 offspring homozygous for the transgene. In one embodiment, hemizygotic G2 offspring from the same line can be bred to produce G3 offspring homozygous for the transgene. In one embodiment, hemizygous G0 animals are bred together to give rise to homozygous G1 offspring. These are merely examples of certain useful breeding schemes. The present invention contemplates the employment of any useful breeding scheme such as those known to individuals of ordinary skill in the art.

In one embodiment, the production of transchromosomic avians which are mature avians developed from the recombinant embryos, ovum and/or sperm of the invention typically are referred to as the G0 generation and are usually chimeric for the

artificial chromosome. The G0 generation may be bred to non-transgenic (non-transchromosomal) birds to give rise to G1 transchromosomal offspring which contain the artificial chromosome in their genome in all or most cells in the bird. The G1 offspring may in turn be bred to non-transchromosomal birds giving rise to G2 offspring with a single copy of the artificial chromosome in their genome. It is also contemplated that birds which contain the artificial chromosome in their genome in all or most cells (e.g., G1 and/or G2 birds) may be bred together to give rise to offspring containing two of the artificial chromosome in their genome. It is contemplated that this process can be repeated, for example, by crossing the offspring containing two copies of the artificial chromosome in their genome, thus producing birds containing multiple copies, for example, four copies of the artificial chromosome in their genome. It is contemplated that this process can be repeated or modified as would be understood by a practitioner of skill in the art to obtain a bird with a desired number of artificial chromosomes contained in its genome. In one useful embodiment, artificial chromosomes of different types or which contain different nucleotide sequences are individually introduced into the genome of individual avians which are bred to produce an avian containing more than one type of artificial chromosome in its genome.

In one aspect, transchromosomal avians of the invention have a genome which includes a transgene of greater than about 5,000 nucleotides in length. In another aspect, transchromosomal avians of the invention have a genome which includes a transgene of between about 5,000 and about 50,000,000 nucleotides in length. For example, the transgene may be between about 5,000 nucleotides in length and about 5,000,000 nucleotides in length. In one embodiment, the transgene is between about 5,000 nucleotides in length and about 1,000,000 nucleotides in length. For example, the transgene may be between about 5,000 nucleotides in length and about 500,000 nucleotides in length.

In one aspect, transchromosomal avians of the invention have a genome which includes a transgene greater than about 8,000 nucleotides in length. In another aspect, transchromosomal avians of the invention have a genome which includes a transgene

of between about 8,000 and about 50,000,000 nucleotides in length. For example, the transgene may be between about 8,000 nucleotides in length and about 5,000,000 nucleotides in length. In one embodiment, the transgene is between about 8,000 nucleotides in length and about 1,000,000 nucleotides in length. For example, the transgene may be between about 8,000 nucleotides in length and about 500,000 nucleotides in length.

In one particularly useful embodiment, the transchromosomal avians of the invention lay eggs which contain one or more heterologous proteins, for example, one or more proteins (e.g., certain pharmaceutical proteins) which are heterologous or exogenous to the egg. The eggs may contain any useful amount of heterologous protein. In one embodiment, the eggs contain the heterologous protein in an amount greater than about 0.01 μg per hard-shell egg. For example, the eggs may contain the heterologous protein in an amount in a range of between about 0.01 μg per hard-shell egg and about 2 grams per hard-shell egg. In one embodiment, the eggs contain between about 0.1 μg per hard-shell egg and about 1 gram per hard-shell egg. For example, the eggs may contain between about 1 μg per hard-shell egg and about 1 gram per hard-shell egg. In one embodiment, the eggs contain between about 1 μg per hard-shell egg and about 1 gram per hard-shell egg. For example, the eggs may contain between about 10 μg per hard-shell egg and about 1 gram per hard-shell egg (e.g., the eggs may contain between about 10 μg per hard-shell egg and about 100 mg per hard-shell egg).

In one useful embodiment, the heterologous protein is present in the egg white of the eggs. In another useful embodiment, the heterologous protein is present in the egg white and is substantially not present in the egg yolk of the eggs.

In one embodiment, the heterologous protein is present in egg white in an amount greater than about 0.01 μg per ml of the egg. In another embodiment, the heterologous protein is present in egg white in an amount in a range of between about 0.01 μg per ml of the egg white and about 0.2 gram per ml of the egg white. For example, the heterologous protein may be present in egg white in an amount in a range of between about 0.1 μg per ml of the egg white and about 0.5 gram per ml of the egg

white. In one embodiment, the heterologous protein is present in egg white in an amount in a range of between about 1 µg per ml of the egg white and about 0.2 gram per ml of the egg white. For example, the heterologous protein may be present in egg white in an amount in a range of between about 1 µg per ml of the egg white and about 0.1 gram per ml of the egg white (e.g., the heterologous protein may be present in egg white in an amount in a range of between about 1 µg per ml of the egg white and about 10 mg per ml of the egg white).

Certain publications considered to be useful in the present invention, the disclosures of which are incorporated in their entirety herein by reference, include:

10 Iadonato et al (1996) RARE-cleavage analysis of YACs, *Methods Mol Biol* 54: 75-85; Popov et al. (1999) A human immunoglobulin lambda locus is similarly well expressed in mice and humans, *J Exp Med* 189(10): 1611-20; Call et al. (2000) A cre-lox recombination system for the targeted integration of circular yeast artificial chromosomes into embryonic stem cells, *Hum Mol Genet* 9(12): 1745-51; Csonka et al. (2000) Novel generation of human satellite DNA-based artificial chromosomes in mammalian cells, *Journal of Cell Science* 113, 3207-3216; Gogel et al. (1996) Mapping of replication initiation sites in the mouse ribosomal gene cluster, *Chromosoma* 104, 511-518; Peterson et al. (1998) LCR-dependent gene expression in beta-globin YAC transgenics: detailed structural studies validate functional analysis even in the presence of fragmented YACs, *Hum Mol Genet* 7(13): 2079-88; Marschall et al. (1999) Transfer of YACs up to 2.3 mb intact into human cells with polyethylenimine, *Gene Ther* 6(9): 1634-7; Basu, J., G. Stromberg et al. (2005) Rapid creation of BAC-based human artificial chromosome vectors by transposition with synthetic alpha-satellite arrays, *Nucleic Acids Res* 33(2): 587-96; Lindenbaum et al. (2004) A mammalian artificial chromosome engineering system (ACE System) applicable to biopharmaceutical protein production, transgenesis and gene-based cell therapy, *Nucleic Acids Res* 32(21): e172; Nicholson et al. (1999) Antibody repertoires of four- and five-feature translocus mice carrying human immunoglobulin heavy chain and kappa and lambda light chain yeast artificial chromosomes, *J Immunol* 163(12): 6898-906; Huxley (1994) *Genetic Engineering*. J. K. Setlow, New York, NY, Plenum

Press, 16: 65-91; Harvey et al. (2002) Consistent Production of Transgenic Chickens using Replication Deficient Retroviral Vectors and High-throughput Screening Procedures, Poultry Science 81(2): 202-12; Tomizuka et al (1997) Functional expression and germline transmission of a human chromosome fragment in chimeric mice, Nature Genetics 16:133-143; and Williams et al (1993) Cloning and sequencing of human immunoglobulin V-lambda gene segments, Eur J Immunol 23:1456-1461.

Any useful combination of features described herein is included within the scope of the present invention provided that the features included in any such combination are not mutually inconsistent as will be apparent from the context, this specification, and the knowledge of one of ordinary skill in the art. For example, the term transgenic can encompass the term transchromosomal and methodologies useful for transgenic animals (e.g., avians) and cells disclosed herein may also be employed for transchromosomal avians and avian cells.

Additional objects and aspects of the present invention will become more apparent upon review of the detailed description set forth below when taken in conjunction with the accompanying figures, which are briefly described as follows.

Brief Description of the Figures

Fig. 1 illustrates phage integrase-mediated integration. A plasmid vector bearing the transgene includes the attB recognition sequence for the phage integrase. The vector along with integrase-coding mRNA, a vector expressing the integrase, or the integrase protein itself, are delivered into cells or embryos. The integrase recognizes DNA sequences in the avian genome similar to attP sites, termed pseudo-attP, and mediates recombination between the attB and pseudo-attP sites, resulting in the permanent integration of the transgene into the avian genome.

Fig. 2 illustrates the persistent expression of luciferase from a nucleic acid molecule after phiC3 1 integrase-mediated integration into chicken cells.

Fig. 3 illustrates the results of a puromycin resistance assay to measure phiC3 1 integrase-mediated integration into chicken cells.

Fig. 4 illustrates phiC31 integrase-mediated integration into quail cells. Puromycin resistance vectors bearing attB sites were cotransfected with phiC31 integrase, or a control vector, into QT6 cells, a quail fibrosarcoma cell line. One day after transfection, puromycin was added. Puromycin resistant colonies were counted
5 12 days post-transfection.

Figs. 5A and 5B illustrate that phiC31 integrase can facilitate multiple integrations per avian cell. A puromycin resistance vector bearing an attB site was cotransfected with an enhanced green fluorescent protein (EGFP) expression vector bearing an attB site, and a phiC31 integrase expression vector. After puromycin
10 selection, many puromycin resistant colonies expressed EGFP in all of their cells. Figs. 5A and 5B are the same field of view with EGFP illuminated with ultraviolet light (Fig. 5A) and puromycin resistant colonies photographed in visible light (Fig. 5B). In Fig. 5B, there are 4 puromycin resistant colonies, two of which are juxtaposed at the top. One of these colonies expressed EGFP.

15 Fig. 6 shows maps of the small vectors used for integrase assays.

Fig. 7 shows integrase promotes efficient integration of large transgenes in avian cells.

Fig. 8 shows maps of large vectors used for integrase assays.

Fig. 9a and b illustrates the nucleotide sequence of the integrase-expressing
20 plasmid pCMV-3 lint (SEQ ID NO: 1).

Fig. 10a and b illustrates the nucleotide sequence of the plasmid pCMV-luc-attB (SEQ ID NO: 2).

Fig. 11a and b illustrates the nucleotide sequence of the plasmid pCMV-luc-attP (SEQ ID NO: 3).

25 Fig. 12a and b illustrates the nucleotide sequence of the plasmid pCMV-pur-attB (SEQ ID NO: 4).

Fig. 13a and b illustrates the nucleotide sequence of the plasmid pCMV-pur-attP (SEQ ID NO: 5).

30 Fig. 14a and b illustrates the nucleotide sequence of the plasmid pCMV-EGFP-attB (SEQ ID NO: 6).

Fig. 15a to f illustrates the nucleotide sequence of the plasmid p12.0-lys-LSPIPNMM-CMV-pur-attB (SEQ ID NO: 7).

Fig. 16a to f illustrates the nucleotide sequence of the plasmid pOMIFN-Ins-CMV-pur-attB (SEQ ID NO: 8).

5 Fig. 17a and b illustrates the nucleotide sequence of the integrase-expressing plasmid pRSV-Int (SEQ ID NO: 9).

Fig. 18a and b illustrates the nucleotide sequence of the plasmid pCR-XL-TOPO-CMV-pur-attB (SEQ ID NO: 10).

10 Fig. 19 illustrates the nucleotide sequence of the attP containing polynucleotide SEQ ID NO: 11.

Fig. 20 illustrates in schematic form the integration of a heterologous att recombination site into an isolated chromosome. The attB sequence is linked to selectable marker such as a puromycin expression cassette and is flanked by sequences found in the target site of the chromosome to be modified. The DNA is transfected
15 into cells containing the chromosome and stable transfectants are selected for by drug resistance. Site specific integration may be confirmed by several techniques including PCR.

Fig. 21 illustrates the persistent expression of luciferase from a nucleic acid molecule after phiC31 integrase-mediated integration into chicken cells bearing a
20 wild-type attP sequence.

Fig. 22 illustrates the distribution of plasmid DNA in a stage I embryo.

Fig. 23 illustrates the distribution of plasmid DNA in a stage I embryo in the presence of low molecular weight polyethylenimine.

25 Fig. 24 illustrates the distribution of plasmid DNA in a stage I embryo in the presence of low molecular weight polyethylenimine.

Fig. 25 illustrates the integration of a gene of interest (i.e., transgene OMC24-IRES-EPO) into an artificial chromosome by integration (which takes place inside of a host cell) wherein cells containing the recombinant chromosome can be selected for based on hygromycin resistance.

Fig. 26 illustrates the insertion of a nucleotide sequence of interest (A) into an attP site contained in an ALV genome which has been integrated into a chicken chromosome (B). The nucleotide sequence can be introduced into a cell containing the ALV genome by any useful method such as microinjection or transduction. For example, the nucleotide sequence can be introduced into an avian egg or germinal disc at any useful stage of development. For example, the nucleotide sequence can be introduced into a stage X egg by transduction. In another example, the nucleotide sequence can be introduced into a stage I egg by microinjection.

Fig. 27 shows human light-chain locus (27A) and heavy-chain locus (27B) containing YACs. V regions are numbered according to their gene family and their position in the locus, following the system of Lefranc et al (1999) IMCT, the international ImMuunoGenTics database Nucleic Acids Res. 27:209, the disclosure of which is incorporated in its entirety herein by reference. The Ig Heavy YAC contains the complete D and J region loci, the intro enhancer (not marked) and the Ig μ and Ig δ C regions. The IgLambda YAC contains the seven paired λ J and C regions, four of which are functional, and the 3' enhancer.

Definitions and Abbreviations

For convenience, definitions of certain terms and certain abbreviations employed in the specification, examples and appended claims are collected here.

Abbreviations used in the present specification include the following: aa, amino acid(s); bp, base pair(s); kb, kilobase(s); mb, megabase(s); art, bacterial recombination attachment site; IU, infectious units; mg, milligram(s); μ g, microgram(s); ml, milliliter(s).

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise. Thus, for example, reference to "an antigen" includes a mixture of two or more such agents.

The term "antibody" as used herein refers to polyclonal and monoclonal antibodies and fragments thereof, and immunologic binding equivalents thereof.

Antibodies may include, but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

5 As used herein, an "artificial chromosome" is a nucleic acid molecule that can stably replicate and segregate alongside endogenous chromosomes in a cell. Artificial chromosomes have the capacity to act as gene delivery vehicles by accommodating and expressing foreign genes contained therein. A mammalian artificial chromosome (MAC) refers to chromosomes that have an active mammalian centromere(s). Plant
10 artificial chromosomes, insect artificial chromosomes and avian artificial chromosomes refer to chromosomes that include plant, insect and avian centromeres, respectively. A human artificial chromosome (HAC,) refers to chromosomes that include human centromeres. For exemplary artificial chromosomes, see, for example, U.S. Pat. Nos. 6,025,155, issued February 15, 2000; 6,077,697, issued June 6, 2000;
15 5,288,625, issued February 22, 1994; 5,712,134, issued January 27, 1998; 5,695,967, issued December 9, 1997; 5,869,294, issued February 9, 1999; 5,891,691, issued April 6, 1999 and 5,721,118, issued February 24, 1998 and published International PCT application Nos., WO 97/40183, published October 30, 1997; WO 98/08964,
20 published March 5, 1998, published US Patent Applications, Serial Nos. 08/835,682, filed April 10, 1997; 10/151,078, filed May 16, 2002; 10/235,119, filed September 3, 2002; 10/086,745, filed February 28, 2002, the disclosures of which are incorporated herein in their entireties by reference. The term "chromosome" may be used interchangeably with the term "artificial chromosome" as will be apparent based on the context of such use.

25 Foreign genes that can be contained in artificial chromosome expression systems can include, but are not limited to, nucleic acid that encodes therapeutically effective substances, such as anti-cancer agents, enzymes, hormones and antibodies. Other examples of heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins (reporter genes), such as fluorescent proteins, such
30 as green, blue or red fluorescent proteins (GFP, BFP and RFP, respectively), other

reporter genes, such as beta-galactosidase and proteins that confer drug resistance, such as a gene encoding hygromycin-resistance.

The term "avian" as used herein refers to any species, subspecies or race of organism of the taxonomic class avia, such as, but not limited to chicken, turkey, duck, 5 goose, quail, pheasants, parrots, finches, hawks, crows and ratites including ostrich, emu and cassowary. The term includes the various known strains of Gallus gallus, or chickens, (for example, White Leghorn, Brown Leghorn, Barred-Rock, Sussex, New Hampshire, Rhode Island, Australorp, Minorca, Amrook, California Gray), as well as strains of turkeys, pheasants, quails, duck, ostriches and other poultry commonly bred 10 in commercial quantities. It also includes an individual avian organism in all stages of development, including embryonic and fetal stages. The term "avian" also may denote "pertaining to a bird", such as "an avian (bird) cell."

The terms "chimeric animal" or "mosaic animal" are used herein to refer to an animal in which a nucleotide sequence of interest is found in some but not all cells of 15 the animal, or in which the recombinant nucleic acid is expressed, in some but not all cells of the animal. The term "tissue-specific chimeric animal" indicates that the recombinant gene is present and/or expressed in some tissues but not others.

The term "coding region" as used herein refers to a continuous linear arrangement of nucleotides which may be translated into a polypeptide. A full length 20 coding region is translated into a full length protein; that is, a complete protein as would be translated in its natural state absent any post-translational modifications. A full length coding region may also include any leader protein sequence or any other region of the protein that may be excised naturally from the translated protein.

The term "cytokine" as used herein refers to any secreted polypeptide that 25 affects a function of cells and modulates an interaction between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines. Examples of cytokines include, but are not limited to, interferon α 2b, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor- α (TNF- α) and Tumor Necrosis Factor α (TNF- α).

30 As used herein, "delivery," which is used interchangeably with "transfection,"

refers to the process by which exogenous nucleic acid molecules are transferred into a cell such that they are located inside the cell.

As used herein, "DNA" is meant to include all types and sizes of DNA molecules including cDNA, plasmids and DNA including modified nucleotides and
5 nucleotide analogs.

The term "expressed" or "expression" as used herein refers to the transcription from a gene to give an RNA nucleic acid molecule at least complementary in part to a region of one of the two nucleic acid strands of the gene. The term "expressed" or "expression" as used herein may also refer to the translation from an RNA molecule to
10 give a protein, a polypeptide or a portion thereof. In one embodiment, for heterologous nucleic acid to be expressed in a host cell, it must initially be delivered into the cell and then, once in the cell, ultimately reside in the nucleus.

The term "gene" or "genes" as used herein refers to nucleic acid sequences that encode genetic information for the synthesis of a whole RNA, a whole protein, or any
15 portion of such whole RNA or whole protein. Genes that are not naturally part of a particular organism's genome are referred to as "foreign genes," "heterologous genes" or "exogenous genes" and genes that are naturally a part of a particular organism's genome are referred to as "endogenous genes". The term "gene product" refers to an RNA or protein that is encoded by the gene. "Endogenous gene products" are RNAs
20 or proteins encoded by endogenous genes. "Heterologous gene products" are RNAs or proteins encoded by "foreign, heterologous or exogenous genes" and are, therefore, not naturally expressed in the cell.

As used herein, the terms "heterologous" and "foreign" with reference to nucleic acids, such as DNA and RNA, are used interchangeably and refer to nucleic
25 acid that does not occur naturally as part of a chromosome, a genome or cell in which it is present or which is found in a location(s) and/or in amounts that differ from the location(s) and/or amounts in which it occurs in nature. It can be nucleic acid that is not endogenous to the genome, chromosome or cell and has been exogenously introduced into the genome, chromosome or cell. Examples of heterologous DNA
30 include, but are not limited to, DNA that encodes a gene product or gene product(s) of

interest, for example, for production of an encoded protein. Examples of heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins, DNA that encodes therapeutically effective substances, such as anti-cancer agents, enzymes and hormones and as antibodies. The terms "heterologous" and "exogenous" in general refer to a biomolecule such as a nucleic acid or a protein that is not normally found in a certain cell, tissue or other component contained in or produced by an organism. For example, a protein that is heterologous or exogenous to an egg is a protein that is not normally found in the egg.

The term "immunoglobulin polypeptide" as used herein refers to a constituent polypeptide of an antibody or a polypeptide derived therefrom. An "immunological polypeptide" may be, but is not limited to, an immunological heavy or light chain and may include a variable region, a diversity region, joining region and a constant region or any combination, variant or truncated form thereof. The term "immunological polypeptides" further includes single-chain antibodies comprised of, but not limited to, an immunoglobulin heavy chain variable region, an immunoglobulin light chain variable region and optionally a peptide linker.

The terms "integrase" and "integrase activity" as used herein refer to a nucleic acid recombinase of the serine recombinase family of proteins.

The term "internal ribosome entry sites (IRES)" as used herein refers to a region of a nucleic acid, most typically an RNA molecule, wherein eukaryotic initiation of protein synthesis occurs far downstream of the 5' end of the RNA molecule. A 43S pre-initiation complex comprising the eIF2 protein bound to GTP and Met-tRNA^{iMet}, the 40S ribosomal subunit, and factors eIF3 and 3fIA may bind to an "IRES" before locating an AUG start codon. An "IRES" may be used to initiate translation of a second coding region downstream of a first coding region, wherein each coding region is expressed individually, but under the initial control of a single upstream promoter. An "IRES" may be located in a eukaryotic cellular mRNA.

As used herein, the term "large nucleic acid molecules" or "large nucleic acids" refers to a nucleic acid molecule of at least about 0.05 mb in size, greater than 0.5 mb, including nucleic acid molecules at least about 0.6, 0.7, 0.8, 0.9, 1, 5, 10, 30, 50 and

100, 200, 300, 500 mb in size. Large nucleic acid molecules typically can be on the order of about 10 to about 450 or more mb, and can be of various sizes, such as, for example, from about 250 to about 400 mb, about 150 to about 200 mb, about 90 to about 120 mb, about 60 to about 100 mb and about 15 to 50 mb. A large nucleic acid molecule may be larger than about 8 kb (e.g., about 8 kb to about 1 mb) as will be apparent based on the context.

Examples of large nucleic acid molecules include, but are not limited to, natural chromosomes and fragments thereof, especially mammalian chromosomes and fragments thereof which retain a centromere or retain a centromere and telomeres, artificial chromosome expression systems (ACEs which include a mouse centromere; also called satellite DNA-based artificial chromosomes (SATACs); see U.S. Pat. Nos. 6,025,155, issued February 15; and 6,077,697, issued June 20, 2000), mammalian artificial chromosomes (MACs), plant artificial chromosomes, insect artificial chromosomes, avian artificial chromosomes and minichromosomes (see, e.g., U.S. Pat. Nos. 5,712,134, issued January 27, 1998; 5,891,691, issued April 6, 1999; and 5,288,625, issued February 22, 1994). Useful large nucleic acid molecules can include a single copy of a desired nucleic acid fragment encoding a particular nucleotide sequence, such as a gene of interest (transgene of interest), or can carry multiple copies thereof or multiple genes or different heterologous sequences of nucleotides. For example, the chromosomes may carry 1 to about 100 or 1 to about 1000 or even more copies of a gene of interest. Large nucleic acid molecules can be associated with proteins, for example chromosomal proteins, that typically function to regulate gene expression and/or participate in determining overall structure.

A "monoclonal antibody" is an antibody in a population of antibodies each of which have the same primary structure.

"Native" as used herein means being naturally associated with or a substance that is produced by a component or organism of interest (in which case the substance would be native to the component or organism) or being in an original form.

A "nucleic acid fragment of interest" or "nucleotide sequence of interest" may be a trait-producing sequence, by which it is meant a sequence conferring a non-native

trait upon the cell in which the protein encoded by the trait-producing sequence is expressed. The term "non-native" when used in the context of a trait-producing sequence means that the trait produced is different than one would find in an unmodified organism which can mean that the organism produces high amounts of a natural substance in comparison to an unmodified organism, or produces a non-natural substance. For example, the genome of a bird could be modified to produce proteins not normally produced in birds such as, for example, useful animal proteins (e.g., human proteins) such as hormones, cytokines and antibodies.

A nucleic acid fragment of interest may additionally be a "marker nucleic acid" or expressed as a "marker polypeptide". Marker genes encode proteins that can be easily detected in transformed cells and are, therefore, useful in the study of those cells. Examples of suitable marker genes include β -galactosidase, green or yellow fluorescent proteins, enhanced green fluorescent protein, chloramphenicol acetyl transferase, luciferase, and the like. Such regions may also include those 5' noncoding sequences involved with initiation of transcription and translation, such as the enhancer, TATA box, capping sequence, CAAT sequence, and the like.

As used herein, "nucleic acid" refers to a polynucleotide containing at least two covalently linked nucleotide or nucleotide analog subunits. A nucleic acid can be a deoxyribonucleic acid (DNA), a ribonucleic acid (RNA), or an analog of DNA or RNA. Nucleotide analogs are commercially available and methods of preparing polynucleotides containing such nucleotide analogs are known (Lin et al. (1994) Nucl. Acids Res. 22:5220-5234; Jellinek et al. (1995) Biochemistry 34:11363-11372; Pagratis et al. (1997) Nature Biotechnol. 15:68-73). The nucleic acid can be single-stranded, double-stranded, or a mixture thereof. For purposes herein, unless specified otherwise, the nucleic acid is double-stranded, or if it is apparent from the context that the nucleic acid is not double stranded. Nucleic acids include any natural or synthetic linear and sequential array of nucleotides and nucleosides, for example cDNA, genomic DNA, mRNA, tRNA, oligonucleotides, oligonucleosides and derivatives thereof. For ease of discussion, certain nucleic acids may be collectively referred to herein as "constructs," "plasmids," or "vectors."

Techniques useful for isolating and characterizing the nucleic acids and proteins of the present invention are well known to those of skill in the art and standard molecular biology and biochemical manuals may be consulted to select suitable protocols without undue experimentation. See, for example, Sambrook et al,
5 1989, "Molecular Cloning: A Laboratory Manual", 2nd ed., Cold Spring Harbor, the content of which is herein incorporated by reference in its entirety.

A "nucleoside" is conventionally understood by workers of skill in fields related to the present invention as comprising a monosaccharide linked in glycosidic linkage to a purine or pyrimidine base. A "nucleotide" comprises a nucleoside with at
10 least one phosphate group appended, typically at a 3' or a 5' position (for pentoses) of the saccharide, but may be at other positions of the saccharide. A nucleotide may be abbreviated herein as "nt." Nucleotide residues occupy sequential positions in an oligonucleotide or a polynucleotide. Accordingly a modification or derivative of a nucleotide may occur at any sequential position in an oligonucleotide or a
15 polynucleotide. All modified or derivatized oligonucleotides and polynucleotides are encompassed within the invention and fall within the scope of the claims. Modifications or derivatives can occur in the phosphate group, the monosaccharide or the base.

By way of nonlimiting examples, the following descriptions provide certain
20 modified or derivatized nucleotides. The phosphate group may be modified to a thiophosphate or a phosphonate. The phosphate may also be derivatized to include an additional esterified group to form a triester. The monosaccharide may be modified by being, for example, a pentose or a hexose other than a ribose or a deoxyribose. The monosaccharide may also be modified by substituting hydroxyl groups with hydro or
25 amino groups, by esterifying additional hydroxyl groups. The base may be modified as well. Several modified bases occur naturally in various nucleic acids and other modifications may mimic or resemble such naturally occurring modified bases. Nonlimiting examples of modified or derivatized bases include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-
30 (carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-

carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, 5 beta-D-mannosylqueosine, S'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaininopurine. Nucleotides may also be modified to harbor a label. Nucleotides may also bear a fluorescent label or a biotin label.

The term "operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Control sequences operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control sequences need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

"Therapeutic proteins" or "pharmaceutical proteins" include an amino acid sequence which in whole or in part makes up a drug. In one embodiment, a pharmaceutical composition or therapeutic composition includes one or more pharmaceutical proteins or therapeutic proteins.

The terms "polynucleotide," "oligonucleotide," and "nucleic acid sequence" are used interchangeably herein and include, but are not limited to, coding sequences (polynucleotide(s) or nucleic acid sequence(s) which are transcribed and translated into polypeptide in vitro or in vivo when placed under the control of appropriate regulatory or control sequences); control sequences (e.g., translational start and stop codons, promoter sequences, ribosome binding sites, polyadenylation signals, transcription factor binding sites, transcription termination sequences, upstream and

downstream regulatory domains, enhancers, silencers, and the like); and regulatory sequences (DNA sequences to which a transcription factor(s) binds and alters the activity of a gene's promoter either positively (induction) or negatively (repression). No limitation as to length or to synthetic origin are suggested by the terms described above.

As used herein the terms "peptide," "polypeptide" and "protein" refer to a polymer of amino acids in a serial array, linked through peptide bonds. A "peptide" typically is a polymer of at least two to about 30 amino acids linked in a serial array by peptide bonds. The term "polypeptide" includes proteins, protein fragments, protein analogues, oligopeptides and the like. The term "polypeptides" contemplates polypeptides as defined above that are encoded by nucleic acids, produced through recombinant technology (isolated from an appropriate source such as a bird), or synthesized. The term "polypeptides" further contemplates polypeptides as defined above that include chemically modified amino acids or amino acids covalently or noncovalently linked to labeling moieties.

The terms "percent sequence identity" or "percent sequence similarity" as used herein refer to the degree of sequence identity between two nucleic acid sequences or two amino acid sequences as determined using the algorithm of Karlin & Attschul, Proc. Natl. Acad. Sci. 87: 2264-2268 (1990), modified as in Karlin & Attschul, Proc. Natl. Acad. Sci. 90: 5873-5877 (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Attschul et al, 1990, T. Mol. Biol. 215: 403-410. BLAST nucleotide searches are performed with the NBLAST program, score = 100, word length = 12, to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches are performed with the XBLAST program, score = 50, word length = 3, to obtain amino acid sequences homologous to a reference polypeptide. To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized as described in Attschul et al, Nucl. Acids Res. 25: 3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g. XBLAST and NBLAST) are used. Other algorithms, programs and default settings may also be suitable such as, but not only, the GCG-

Sequence Analysis Package of the U.K. Human Genome Mapping Project Resource Centre that includes programs for nucleotide or amino acid sequence comparisons. Examples of useful algorithms are FASTA and BESTFIT.

5 The term "polyclonal antibodies" as used herein refers to a population of antibodies each of which recognize the same antigen or each of which recognize an antigen of a substance which contains one or more antigens.

10 The term "promoter" as used herein refers to the DNA sequence that determines the site of transcription initiation by an RNA polymerase. A "promoter-proximal element" is a regulatory sequence generally within about 200 base pairs of the transcription start site.

15 The term "pseudo-recombination site" as used herein refers to a site at which an integrase can facilitate recombination even though the site may not have a sequence identical to the sequence of its wild-type recombination site. For example, a phiC31 integrase and vector carrying a phiC31 wild-type recombination site can be placed into an avian cell. The wild-type recombination sequence aligns itself with a sequence in the avian cell genome and the integrase facilitates a recombination event. When the sequence from the genomic site in the avian cell, where the integration of the vector took place, is examined, the sequence at the genomic site typically has some identity to, but may not be identical with, the wild-type bacterial genome recombination site.

20 The recombination site in the avian cell genome is considered to be a pseudo-recombination site (c.g., a pseudo-attP site) at least because the avian cell is heterologous to the normal phiC31 phage/bacterial cell system. The size of the pseudo-recombination site can be determined through the use of a variety of methods including, but not limited to, (i) sequence alignment comparisons, (ii) secondary

25 structural comparisons, (iii) deletion or point mutation analysis to find the functional limits of the pseudo-recombination site, and (iv) combinations of the foregoing.

30 The terms "recombinant cell" and "genetically transformed cell" refer to a cell comprising a combination of nucleic acid segments not found in a single cell with each other in nature. A new combination of nucleic acid segments can be introduced into an organism using a wide array of nucleic acid manipulation techniques available to

those skilled in the art. The recombinant cell may harbor a vector that is extragenomic, i.e. that does not covalently insert into the cellular genome, including a non-nuclear (e.g. mitochondrial) genome(s). A recombinant cell may further harbor a vector or a portion thereof that is intragenomic, i.e. covalently incorporated within the genome of the recombinant cell.

The term "recombination site" as used herein refers to a polynucleotide stretch comprising a recombination site normally recognized and used by an integrase. For example, λ phage is a temperate bacteriophage that infects *E. coli*. The phage has one attachment site for recombination (attP) and the *E. coli* bacterial genome has an attachment site for recombination (attB). Both of these sites are recombination sites for λ integrase. Recombination sites recognized by a particular integrase can be derived from a homologous system and associated with heterologous sequences, for example, the attP site can be placed in other systems to act as a substrate for the integrase.

The terms "recombinant nucleic acid" and "recombinant DNA" as used herein refer to combinations of at least two nucleic acid sequences that are not naturally found in a eukaryotic or prokaryotic cell. The nucleic acid sequences may include, but are not limited to, nucleic acid vectors, gene expression regulatory elements, origins of replication, suitable gene sequences that when expressed confer antibiotic resistance, protein-encoding sequences and the like. The term "recombinant polypeptide" is meant to include a polypeptide produced by recombinant DNA techniques. A recombinant polypeptide may be distinct from a naturally occurring polypeptide either in its location, purity or structure. Generally, a recombinant polypeptide will be present in a cell in an amount different from that normally observed in nature.

As used herein, the term "satellite DNA-based artificial chromosome (SATAC)" (e.g., ACE) is a type of artificial chromosome. These artificial chromosomes are substantially all neutral non-coding sequences (heterochromatin) except for foreign heterologous, typically gene-encoding nucleic acid, that is present within (see U.S. Pat. Nos. 6,025,155, issued February 15, 2000 and 6,077,697, issued

June 20, 2000 and International PCT application No. WO 97/40183, published October 30, 1997).

The term "source of integrase activity" as used herein refers to a polypeptide or multimeric protein having serine recombinase (integrase) activity in an avian cell. The
5 term may further refer to a polynucleotide encoding the serine recombinase, such as an mRNA, an expression vector, a gene or isolated gene that may be expressed as the recombinase-specific polypeptide or protein.

As used herein the term "therapeutic substance" refers to a component that comprises a substance which can provide for a therapeutic effect, for example, a
10 therapeutic protein.

"Transchromosomic avian" means an avian which contains an artificial chromosome in some or all of its cells. A transchromosomic avian can include the artificial chromosome in its germ cells.

The term "transcription regulatory sequences" as used herein refers to
15 nucleotide sequences that are associated with a gene nucleic acid sequence and which regulate the transcriptional expression of the gene. Exemplary transcription regulatory sequences include enhancer elements, hormone response elements, steroid response elements, negative regulatory elements, and the like.

The term "transfection" as used herein refers to the process of inserting a
20 nucleic acid into a host cell. Many techniques are well known to those skilled in the art to facilitate transfection of a nucleic acid into an eukaryotic cell. These methods include, for instance, treating the cells with high concentrations of salt such as a calcium or magnesium salt, an electric field, detergent, or liposome mediated transfection, to render the host cell competent for the uptake of the nucleic acid
25 molecules, and by such methods as micro-injection into a pro-nucleus, sperm-mediated and restriction-mediated integration.

The term "transformed" as used herein refers to a heritable alteration in a cell resulting from the uptake of a heterologous DNA.

As used herein, the term "transgene" means a nucleic acid sequence that is
30 partly or entirely heterologous, i.e., foreign, to the transgenic animal or cell into which

it is introduced, or, is homologous to an endogenous gene of the transgenic animal or cell into which it is introduced, but which is designed to be inserted, or is inserted, into the animal's genome in such a way as to alter the genome of the cell into which it is inserted (e.g., it is inserted at a location which differs from that of the natural gene or its insertion results in a knockout).

As used herein, a "transgenic avian" is any avian, as defined herein, in which one or more of the cells of the avian contain heterologous nucleic acid introduced by manipulation, such as by transgenic techniques. The nucleic acid may be introduced into a cell, directly or indirectly, by introduction into a precursor of the cell by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. Genetic manipulation also includes classical cross-breeding, or in vitro fertilization. The heterologous nucleic acid may be an artificial chromosome or may be integrated within a chromosome of the avian, or it may be extrachromosomally replicating DNA.

The term "trisomic" as used herein refers to a cell or animal, such as an avian cell or bird that has a $2n+1$ chromosomal complement, where n is the haploid number of chromosomes, for the animal species concerned.

The terms "vector" or "nucleic acid vector" as used herein refer to a natural or synthetic single or double stranded plasmid or viral nucleic acid molecule (RNA or DNA) that can be transfected or transformed into cells and replicate independently of, or within, the host cell genome. The term "expression vector" as used herein refers to a nucleic acid vector that comprises a transcription regulatory region operably linked to a site wherein is, or can be, inserted, a nucleotide sequence to be transcribed and, optionally, to be expressed, for instance, but not limited to, a sequence coding at least one polypeptide.

Detailed Description

The present invention provides for the production of polyclonal antibodies, for example, human polyclonal antibodies, in avians and isolated avian cells. Such avians or avian cells may produce any useful type of antibody including, but not limited to,

one or more of IgG, IgM, IgA, IgE, and IgD including each of the subtypes of these antibodies. For example, subtypes of IgG include IgG1, IgG2, IgG3 and IgG4.

In one particularly useful embodiment, the invention provides for the production of polyclonal antibodies which are deposited in the eggs of avians, such as chickens. It has been shown that active deposition of chicken IgG into the egg is mediated by specific sequences on the Fc portion of the antibody (Morrison et al (2002) *Mol Immunol.* 8:619-25). The IgG Fc antibody portion has also been shown to mediate the deposition into an egg of either intravenously injected human IgG₅ or human monoclonal antibody produced in vivo from a transplanted chicken B-cell line, with high efficiency (Mohammed et al (1998) *Immunotechnology*, 4(2):115-25). Chicken IgY does not bind to protein A or G and therefore human IgG can be easily affinity purified from other proteins including chicken immunoglobulins using protein A and/or protein G based purification methodologies as is known in the art. In a particular aspect, the antibody is deposited in the yolk of the egg of the avian.

The invention provides for the insertion of large DNA segments into the germline of avians or avian cells. In one particular aspect, the large DNA segments include regions encoding components necessary for the production of human polyclonal antibodies. In one embodiment, the DNA segments include one or more Ig loci. The Ig loci may include one or more of human Ig λ , Ig κ , IgH, and portions thereof. The Ig loci may be modified to include additional components, such as additional variable or constant regions, or they may be in their native form. Certain Ig loci and other disclosure which may be useful in accordance with the present invention are disclosed in, for example, US patent application publication No. 2002/0132373, published September 19, 2002; US patent application publication No. 2002/0088016, published July 4, 2002; US patent application publication No. 2004/0231012, published November 18, 2004; US Patent No. 6,348,349, issued February 19, 2002; US Patent No. 5,545,807, issued August 13, 1996; and Popov et al (1999) *J. Exp. Med.* 189: 1611-1619. The disclosures of each of these three published patent applications and two issued patents and journal article are incorporated in their entirety herein by reference. In one useful embodiment, the Ig loci shown in Fig. 27A and 27B

are used to produce polyclonal antibodies in accordance with the present invention. These loci are disclosed in Nicholson, et al (1999) J. Immunology 163(12):6898-6906.

The DNA segments comprising regions encoding components necessary for the production of human polyclonal antibodies may be employed in the invention in any useful form. For example, the DNA may be linear or circular. Typically, the DNA segments are present in a cloning vehicle which will facilitate the germline transmission of the DNA encoding the polyclonal antibodies. In one embodiment, artificial chromosomes which include one or more transgenes comprising components necessary for the production of human polyclonal antibodies are contemplated for use to produce germline transgenic avians of the invention. Typically, in this embodiment, a germline chimeric avian is obtained from embryos or germline cells of avians, such as chickens, into which one or more artificial chromosomes comprising the polyclonal antibody transgenes have been introduced as disclosed herein. Subsequently, a transgenic or fully transgenic (e.g., transchromosomal) G1 bird can be obtained from the germline chimera.

In one useful embodiment, one or more Ig loci are included in an artificial chromosome. The artificial chromosome is introduced into an avian genome as disclosed herein.

In one embodiment two artificial chromosomes are used, one having an Ig heavy chain locus and the other having an Ig light chain locus. In one embodiment, the two artificial chromosomes are co-introduced into an avian embryo to produce a germline-transgenic or transchromosomal avian which contains both chromosomes in its genome.

In another embodiment, one or more DNA segments comprising regions necessary for the production of human polyclonal antibodies (e.g., Ig loci) may be used to produce chimeric and germline transgenic avians by incorporation into the genome of an avian by employing integrase mediated transgenesis as disclosed herein.

The invention also contemplates one or more transgenes comprising components necessary for the production of human polyclonal antibodies such as Ig loci being introduced into an immortalized avian cell line, the cells of which may be

capable of secreting the polyclonal antibodies into growth medium. In one particular embodiment, immortalized cell lines are derived from tumor cells of an avian oviduct or tumor cells from other cells of an avian, for example, cell lines disclosed in US Patent No 10/926,707, filed August 25, 2004.

5 The invention also provides for the production and isolation of cell lines capable of producing monoclonal antibodies. By using standard methodologies well known in the art such as those disclosed in Michael et al (1998) Proc Natl Acad Sci USA 95:1166-1171, the disclosure of which is incorporated in its entirety herein by reference, cells of transgenic avians which contain human Ig heavy chain and human
10 light chain producing loci in their genomes can be used to produce cell lines capable of producing human monoclonal antibodies. For example, the transgenic or transchromosomal chicken is immunized with an antigen and hybridomas are produced by fusing cells (e.g., spleen cells) of the transgenic bird to an immortalized cell line to produce hybridomas. Antibody produced by individual hybridoma clones
15 is screened to identify antibody with binding specificity for the antigen. The exon DNA (e.g., cDNA) encoding the antibody is cloned into mammalian Ig expression vectors which are co-transfected into mammalian myeloma cells to produce antigen specific antibody.

 Further, it is contemplated that immunoglobulin genes and other useful
20 products can be provided by the invention. For example, genes encoding monoclonal antibodies can be obtained from monoclonal antibody producing cell lines produced in accordance with the present invention.

 The present invention contemplates the production of artificial chromosomes containing large transgenes. In one specific embodiment, the invention provides for
25 the production of artificial chromosomes containing yeast artificial chromosomes (YACs) which contain a large DNA insert such as an Ig locus.

 In one embodiment, the present invention provides for the production of artificial chromosomes which contain transgenes wherein the transgene is introduced into the artificial chromosome during the de novo construction of the artificial
30 chromosome. In one particularly useful embodiment of the invention, production of

artificial chromosomes which contain large transgenes (e.g., one or more Ig locus) is provided for. Large transgenes as disclosed herein can refer to transgenes greater in size than, for example, about 8 kb or about 10 kb or about 20 kb (e.g., about 8 kb to about 100 mb in size or about 10 kb to about 100 mb in size).

5 In one embodiment, the invention provides for the introduction of transgene DNA into a cell in which the artificial chromosome is produced at the time of production or assembly of the artificial chromosome. For example, components useful for the production of an artificial chromosome and one or more transgenes are introduced into the cell at about the same time leading to the production of an artificial
10 chromosome containing the transgene or transgenes. In one embodiment, the transgene will include a cloning vector such as a BAC or YAC to which the transgene is linked when introduced into the cell. In another embodiment, the transgene is removed from the cloning vehicle before introduction into the cell. For example, a cloning vehicle containing the transgene can be digested with a nuclease such as a
15 restriction enzyme to release from the cloning vehicle the DNA sequence (transgene) to be included in the artificial chromosome during its assembly. In one embodiment, after restriction nuclease digestion the DNA sequence of interest is separated from the cloning vehicle (e.g., by electrophoresis) prior to introduction of the DNA sequence into the cell. Without wishing to limit the invention to any theory or mechanism of
20 operation, it is believed that as the artificial chromosome is assembled in the cell the transgene(s) is incorporated into the artificial chromosome during the assembly.

 In one embodiment, for artificial chromosome assembly, cells may be cotransfected with the transgene DNA and ribosomal RNA encoding DNA (rDNA). In one embodiment, the rDNA is included in a cloning vehicle such as a plasmid or a
25 cosmid. In one useful aspect of the invention, the cell in which the artificial chromosome is produced provides for certain components which will make up the new artificial chromosome such as telomeric nucleotide sequences. The cells which contain the new transgene containing artificial chromosome are identified and isolated. In one embodiment, the transgene carries a selectable marker such as a drug resistant
30 gene providing for the selection of cells containing the new artificial chromosome.

Spontaneous generation of artificial chromosomes may be accomplished by the introduction of heterologous DNA and a marker gene into a cell such as a fibroblast cell, for example, DF-I cells (US Patent No. 5,672,485, issued September 30, 1997) or chicken embryo fibroblast cells. However, the methods are not limited to use of a
5 fibroblast cell and the invention contemplates the employment of any useful cell. For example, cell lines such as CHO cells, HeLa cells and other animal cell lines, for example, mammalian cell lines, are contemplated for use as disclosed herein. In one embodiment, the present invention contemplates the introduction of a desired transgene into a cell in combination with a marker and heterologous DNA thereby
10 providing for the spontaneous generation of artificial chromosomes containing the desired transgene. The desired transgene typically includes a pharmaceutical protein coding sequence, such as a coding sequence for a pharmaceutical protein disclosed herein, and/or a promoter which functions in the avian oviduct or an active portion thereof. In one useful embodiment, the desired transgene comprises one or more
15 human Ig locus or a portion thereof.

Any useful method for the spontaneous assembly or production of artificial chromosomes is contemplated for use in accordance with the present invention. That is, incorporation of a nucleotide sequence of interest such as a promoter (e.g., ovalbumin promoter, ovomucoid promoter, lysozyme promoter or other promoters
20 which function in the avian oviduct) and/or a coding sequence for a pharmaceutical protein during assembly of the chromosome (e.g., spontaneous assembly) is contemplated. For example, spontaneous assembly of artificial chromosomes (e.g., dicentric chromosomes minichromosomes, satellite artificial chromosomes or megachromosomes) as disclosed in, for example, US Patent No. 6,743,967, issued
25 June 1, 2004; US Patent No. 5,288,625, issued February 22, 1994; and WO97/40183, the disclosures of which are incorporated in their entirety herein by reference, is contemplated for use in conjunction with the present invention.

A selectable marker may be included in one or more vectors which are used in artificial chromosome construction (e.g., transgene containing vectors and/or other
30 vectors containing DNA useful in production of the artificial chromosomes, for

example, and without limitation, rDNA). In the case where multiple vectors are introduced into a cell to produce an artificial chromosome, some or all of the vectors may have a selectable marker. In such a case, the selectable markers may be different selectable markers. Examples of useful selectable markers include, without limitation, genes which provide for resistance to hygromycin, zeomycin, neomycin and blastomycin. In one embodiment, vectors, for example, linearized vectors, when present in a cell that is producing a chromosome of the invention, may incorporate efficiently into the new chromosome, thereby precluding the need for one or more markers.

One advantage of introducing large DNA molecules into an artificial chromosome during its assembly is that large DNA molecules can be gel purified and directly transfected as a linear molecule into the cell line in which the chromosome is being assembled. Gel purification is important for isolating DNA molecules such as YACs from the other components of the host cells including the native cellular chromosomal DNA. Large, linear YACs are routinely purified in intact form by gel purification methods. Large circular YACs (cYAC) are not able to migrate through agarose in pulsed field gel electrophoresis (PFGE) (i.e., the cYACs remain in the wells) and therefore cannot be gel purified.

The present methods are contemplated for the production of artificial chromosomes which contain any useful transgene. In one embodiment, artificial chromosomes which contain immunoglobulin genes (e.g., coding sequences for immunoglobulins and/or certain native gene expression controlling regions for immunoglobulins), such as human immunoglobulin loci or loci portions, are produced. In one particularly useful embodiment, the Ig loci include coding sequences for the immunoglobulins and certain native gene expression controlling regions of immunoglobulins. The human Ig containing artificial chromosome may be introduced into an avian such as a chicken such that the chicken produces human antibodies in its serum and the antibodies localize to the egg. In one useful embodiment, the antibodies are polyclonal in nature and are produced by immunization of the transgenic animal with an antigen. In the case of such transgenic avians, such as chickens, the invention

contemplates the polyclonal human antibodies being deposited in the yolk of laying hens through a native transport system that has been shown to transfer antibodies, including human antibodies, from the blood serum to the yolk of forming eggs. In one embodiment, the invention contemplates the deposition of an amount between about 0.1 μ g and about 1 gram of polyclonal antibody per egg.

Human Ig genes are encoded on separate loci. Human heavy chain (IgH) is believed to be encoded by a single locus that is ~1.5 mb in size. There are believed to be two loci for the human light chain, Igl ϵ and Igl λ , either of which may be used for production of functional antibodies. The Igi ζ locus is believed to be ~1.1 mb and the Igl λ locus is believed to be ~3 mb. The invention contemplates the production of transgenic avians that carry either the light or the heavy chain or both the light and the heavy chain in their genome. For example, the loci may be present on one or more artificial chromosomes introduced into an avian's cells or may be introduced into the avian's genome by integrase mediated recombination as disclosed herein.

In one embodiment, two artificial chromosomes are produced, one containing the light chain and one containing the heavy chain. In one embodiment, each artificial chromosome may be used to produce a separate line of animal (e.g., two lines of chickens). The two lines are crossed and offspring are selected that carry heavy and light chain artificial chromosomes. In another embodiment, the two artificial chromosomes are co-introduced into the avian, e.g., co-injected into a germinal disc.

In another embodiment, an artificial chromosome may be created that carries both the heavy locus and light chain locus allowing generation of a single line of animals capable of producing antibodies.

In one embodiment of the invention, it is contemplated that the Ig gene(s) includes one or more additional variable region genes and/or one or more constant region genes which are not normally present in the Ig gene(s).

Ig genes are polymorphic, particularly in the variable coding regions. Therefore, Ig-artificial chromosomes can be produced that are capable of creating polyclonal antibodies that are specifically enhanced for a particular target antigen. For example, it is found that a human family is particularly resistant to the development of

cancer, for example, a certain type of cancer such as breast cancer. The resistance trait is traced to their heavy and light chain genes, suggesting that this combination of heavy and light chain alleles can produce a mixture of antibodies that are exceptionally able to target and destroy cancer cells such as breast cancer cells. The heavy and light chain genes can be cloned from DNA extracted from a family member and inserted into an artificial chromosome. Therefore, in one embodiment of the invention, a transgenic animal such as a chicken carrying an artificial chromosome will produce polyclonal antibodies such that when immunized with cancer cells, or antigens thereof, such as breast cancer cells, or antigens thereof, polyclonal antibodies will be produced that can be used to treat cancer patients, for example, breast cancer patients.

The present invention provides for recombinant vertebrate cells (e.g., transgenic or transchromosomal avian cells) and transgenic vertebrate animals (e.g., transgenic or transchromosomal avians) and methods of making the cells and the animals. For example, the invention provides for methods of inserting nucleotide sequences into the genome of vertebrate animals or into the cells of vertebrate animals in a site specific manner. Examples of vertebrates include, without limitation, birds, mammals, fish, reptiles and amphibians. Examples of mammals include sheep, goats and cows. In one certain embodiment of the invention, the vertebrate animals are birds or avians. Examples of birds include, without limitation, chickens, turkeys, ducks, geese, quail, pheasants, parrots, finches, hawks, crows and ratites including ostriches, emu and cassowary. Methods disclosed herein for producing transgenic and transchromosomal avians are generally applicable for all avians. For example, though the size of the hard shell egg laid by avians may vary substantially (e.g., hummingbird eggs compared to ostrich eggs), the size and structure of the germinal disc is substantially the same among avians. Therefore, since the present invention, in large part, relies on the injection of large DNA molecules (e.g., artificial chromosomes) into a germinal disc, a practitioner in the art would expect that the invention will function universally among avians.

In one embodiment, the present invention provides for methods of inserting nucleotide sequences into the genome of an animal using methods of transgenesis

based on site specific integration, for example, site specific integrase mediated-transgenesis. The present invention contemplates any useful method of integrase mediated transgenesis including but not limited to, transgenesis mediated by serine recombinases and tyrosine recombinases. Serine recombinases are well known in the art and include without limitation, EcoYBCK₅ ΦC31, SCH10.38c, SCC88.14, SC8F4.15c, SCD12A.23, Bxb1, WwK₅ Sau CcrB, Bsu CisB, TP901-1, Φ370.1, Φ105, ΦFC1, A118, Cae1956, Cae1951, Sau CcrA, Spn, TnpX, TndX, SPBc2, SC3C8.24, SC2E1.37, SCD78.04c, R4₅ ΦRv1, Y4bA₅ BJa₅ SsoISC1904b, SsoISC1904a₅ Aam, MjaMJ1004, Pab₅ SsoISC1913, HpyIS607₅ MceRvO921, MtuRvO921, MtuRv2979c, MtuRv2792c, MtuISY349₅ MtuRv3828c, SauSK1, Spy, EcoTn21, Mlo92₅ EcoTn3, LIa₅ Cpe, SauSK41, BmeTn5083, SfaTn917, Bme53₅ Ran, RmzY4CG₅ SarpNLI, Pje, Xan, ISXc5, Pae, Xca, Req, Mlo90, PpsTn5501, pMER05, CgI₅ MuGin, StyHin, Xfa911₅ Xfa910, Rrh, SauTn552 and Aac serine recombinases. Tyrosine recombinases well known in the art include without limitation, BS codV, BS ripX, BS ydcL, CB tnpA, ColID₅ CP4, Crc, D29, DLP12, DN int, EC FimB, EC FimE, EC orf, EC xerC, EC xerD, Φ11, Φ13₅ Φ80, Φadh, ΦCTX, ΦLC3, FLP, ΦR73, Hlorf, HI rci, HI xerC, HI xerD, HK22, HPI, L2, L5, L54, λ, LL orf, LL xerC, LO L5, MJ orf, ML orf, MP int, MT int, MT orf, MV4, P186, P2, P21, P22, P4, P434, PA sss, PM fmiB, pAEI, pCLI, pKDI, pMEA, pSAM2, pSB2, pSB3, pSDL2, pSEIOI, pSE211, pSMI, pSRI, pWS58, R721, Rci, SF6, SLPI, SM orf, SsrA, SSVI, T12, Tn21, Tn4430, Tn554a, Tn554b, Tn7, Tn916, Tuc, WZ int, XisA and XisC. Other enzymes which may be useful for mediation of transgenesis in accordance with the present invention include, certain transposases, invertases and resolvases.

In certain instances, integration host factors (IHF) may be necessary for the integration of nucleotide sequences of the invention into the genome of cells as disclosed herein. In such a case, the integration host factors may be delivered to the cells directly or they may be delivered to the cells in the form of a nucleic acid which, in the case of RNA, is translated to produce the IHF or, in the case of DNA, is transcribed and translated to produce the IHF.

The present invention contemplates the use of any system capable of site specifically inserting a nucleotide sequence of interest into the genome of a cell, for example, to produce a transgenic vertebrate animal. Typically, although not exclusively, these systems require at least three components: 1) a sequence in the genome which specifies the site of insertion; 2) a nucleotide sequence which is directed to the site of insertion and an enzyme which catalyzes the insertion of the nucleotide sequence into the genome at the site of insertion. Many enzymes, including integrases, which are capable of site specifically inserting nucleotide sequences into the genome have been characterized. Examples of these enzymes are disclosed in for example, Esposito et al (1997) *Nucleic Acids Research*, 25;3605-3614 and Nunes-Düby et al (1998) *Nucleic Acids Research*, 26; 391-406. The disclosure of each of these references is incorporated herein in their entirety.

In one embodiment of the present invention, a serine recombinase is employed. Serine recombinase integrase mediates recombination between an attB site on a transgene vector and an attP or a pseudo attP site on a chromosome. In the method of the invention for integrase-mediated transgenesis, a heterologous wild-type attP site can be integrated into a nuclear genome to create a transgenic cell line or a transgenic vertebrate animal, such as an avian. A serine recombinase (integrase) and an attB-bearing transgene vector are then introduced into cells harboring the heterologous attP site, or into embryos derived from animals which bear the attP recombination site. The locations of attP and attB may be reversed such that the attB site is inserted into a chromosome and the attP sequence resides in an incoming transgene vector. In either case, the att site of the introduced vector would then preferentially recombine with the integrated heterologous att site in the genome of the recipient cell.

The methods of the invention are based, in part, on the discovery that there exists in vertebrate animal genomes, such as avian genomes, a number of specific nucleic acid sequences, termed pseudo-recombination sites, the sequences of which may be distinct from wild-type recombination sites but which can be recognized by a site-specific integrase and used to promote the efficient insertion of heterologous genes or polynucleotides into the targeted nuclear genome. The inventors have

identified pseudo-recombination sites in avian cells capable of recombining with a recombination site, such as an attB site within a recombinant nucleic acid molecule introduced into the target avian cell. The invention is also based on the prior integration of a heterologous att recombination site, typically isolated from a bacteriophage or a modification thereof, into the genome of the target avian cell.

Integration into a predicted chromosomal site is useful to improve the predictability of expression, which is particularly advantageous when creating transgenic avians. Transgenesis by methods that result in insertion of the transgene into random positions of the avian genome is unpredictable since the transgene may not express at the expected levels or in the predicted tissues.

The invention as disclosed herein, therefore, provides methods for site-specifically genetically transforming an avian nuclear genome. In general, an avian cell having a first recombination site in the nuclear genome is transformed with a site-specific polynucleotide construct comprising a second recombination sequence and one or more polynucleotides of interest. Into the same cell, integrase activity may be introduced that specifically recognizes the first and second recombination sites under conditions such that the polynucleotide sequence of interest is inserted into the nuclear genome via an integrase-mediated recombination event between the first and second recombination sites.

The integrase activity, or a source thereof, can be introduced into the cell prior to, or concurrent with, the introduction of the site-specific construct. The integrase can be delivered to a cell as a polypeptide, or by expressing the integrase from a source polynucleotide such as an mRNA or from an expression vector that encodes the integrase, either of which can be delivered to the target cell before, during or after delivery of the polynucleotide of interest. Any integrase that has activity in a cell may be useful in the present invention, including HK022 (Kolot et al, (2003) Biotechnol. Bioeng. 84: 56-60). In one embodiment, the integrase is a serine recombinase as described, for example, by Smith & Thorpe, in Mol. Microbiol., 44: 299-307 (2002). For example, the integrase may be TP901-1 (Stoll et al, J. Bact., 184: 3657-3663 (2002); Olivares et al, Gene, 278:167-176 (2001) or the integrase from the phage

phiC31.

The nucleotide sequence of the junctions between an integrated transgene into the attP (or attB site) would be known. Thus, a PCR assay can be designed by one of skill in the art to detect when the integration event has occurred. The PCR assay for
5 integration into a heterologous wild-type attB or attP site can also be readily incorporated into a quantitative PCR assay using TAQMAN™ or related technology so that the efficiency of integration can be measured.

In one embodiment, the minimal attB and attP sites able to catalyze recombination mediated by the phiC31 integrase are 34 and 39 bp, respectively. In
10 cell lines that harbor a heterologous integrated attP site, however, integrase may have a preference for the inserted attP over any pseudo-attP sites of similar length, because pseudo-attP sites have very low sequence identity (for example, between 10 to 50% identity) compared to the more efficient wild-type attP sequence. It is within the scope of the methods of the invention, however, for the recombination site within the target
15 genome to be a pseudo-att site such as a pseudo-attP site or an attP introduced into a genome.

The sites used for recognition and recombination of phage and bacterial DNAs (the native host system) are generally non-identical, although they typically have a common core region of nucleic acids. In one embodiment, the bacterial sequence is
20 called the attB sequence (bacterial attachment) and the phage sequence is called the attP sequence (phage attachment). Because they are different sequences, recombination can result in a stretch of nucleic acids (for example, attL or attR for left and right) that is neither an attB sequence or an attP sequence, and likely is functionally unrecognizable as a recombination site to the relevant enzyme, thus
25 removing the possibility that the enzyme will catalyze a second recombination reaction that would reverse the first.

The integrase may recognize a recombination site where sequence of the 5' region of the recombination site can differ from the sequence of the 3' region of the recombination sequence. For example, for the phage phiC31 attP (the phage
30 attachment site), the core region is 5'-TTG-3' the flanking sequences on either side are

represented here as attP5' and attP3', the structure of the attP recombination site is, accordingly, attP5'-TTG-attP3'. Correspondingly, for the native bacterial genomic target site (attB) the core region is 5'-TTG-3', and the flanking sequences on either side are represented here as attB5' and attB3', the structure of the attB recombination site is, accordingly, attB5'-TTG-attB3'. After a single-site, phiC31 integrase-mediated recombination event takes place between the phiC31 phage and the bacterial genome, the result is the following recombination product: attB5'-TTG-attP3'{phiC31 vector sequences}-attP5'-TTG-attB3'. In the method of invention, the attB site will be within a recombinant nucleic acid molecule that may be delivered to a target cell. The corresponding attP (or pseudo-attP) site will be within the cell nuclear genome. Consequently, after phiC31 integrase mediated recombination, the recombination product, the nuclear genome with the integrated heterologous polynucleotide will have the sequence attP5'-TTG-attB3'{heterologous polynucleotide}-attB5'-TTG-attP3'. Typically, after recombination the post-recombination recombination sites are no longer able to act as substrate for the phiC31 integrase. This results in stable integration with little or no integrase mediated excision.

While the one useful recombination site to be included in the recombinant nucleic acid molecules and modified chromosomes of the present invention is the attP site, it is contemplated that any attP-like site may be used if compatible with the attB site. For instance, any pseudo-attP site of the chicken genome may be identified according to the methods of Example 7 herein and used as a heterologous att recombination site. For example, such attP-like sites may have a sequence that is greater than at least 25% identical to SEQ ID NO: 11 as shown in Fig. 19, such as described in Groth et al, Proc. Natl. Acad. Sci. U.S.A. 97: 5995-6000 (2000) incorporated herein by reference in its entirety. In one embodiment, the selected site will have a similar degree of efficiency of recombination, for example, at least the same degree of efficiency of recombination as the attP site (SEQ ID NO: 11) itself.

In the present invention, the recipient cell population may be an isolated cell line such as, for example, DF-I chicken fibroblasts, chicken DT40 cells or a cell population derived from an early stage embryo, such as a chicken stage I embryo or

mid stage or late stage (e.g., stage X) embryos. One useful avian cell population is blastodermal cells isolated from a stage X avian embryo. The methods of the present invention, therefore, include steps for the isolation of blastodermal cells that are then suspended in a cell culture medium or buffer for maintaining the cells in a viable state, and which allows the cell suspension to contact the nucleic acids of the present invention. It is also within the scope of the invention for the nucleic acid construct and the source of integrase activity to be delivered directly to an avian embryo such as a blastodermal layer, or to a tissue layer of an adult bird such as the lining of an oviduct.

When the recipient cell population is isolated from an early stage avian embryo, the embryos must first be isolated. For stage I avian embryos from, for example, a chicken, a fertilized ovum is surgically removed from a bird before the deposition of the outer hard shell has occurred. The nucleic acids for integrating a heterologous nucleic acid into a recipient cell genome may then be delivered to isolated embryos by lipofection, microinjection (as described in Example 6 below) or electroporation and the like. After delivery of the nucleic acid, the transfected embryo and its yolk may be deposited into the infundibulum of a recipient hen for the deposition of egg white proteins and a hard shell, and laying of the egg. Stage X avian embryos are obtained from freshly laid fertilized eggs and the blastodermal cells isolated as a suspension of cells in a medium, as described in Example 4 below. Isolated stage X blastodermal cell populations, once transfected, may be injected into recipient stage X embryos and the hard shell eggs resealed according to the methods described in U.S. Patent No. 6,397,777, issued June 4, 2002, the disclosure of which is incorporated in its entirety by reference herein.

In one embodiment of the invention, once a heterologous nucleic acid is delivered to the recipient cell, integrase activity is expressed. The expressed integrase (or injected integrase polypeptide) then mediates recombination between the att site of the heterologous nucleic acid molecule, and the att (or pseudo att) site within the genomic DNA of the recipient avian cell.

It is within the scope of the present invention for the integrase-encoding sequence and a promoter operably linked thereto to be included in the delivered

nucleic acid molecule and that expression of the integrase activity occurs before integration of the heterologous nucleic acid into the cell genome. In one embodiment, an integrase-encoding nucleic acid sequence and associated promoter are in an expression vector that may be co-delivered to the recipient cell with the heterologous
5 nucleic acid molecule to be integrated into the recipient genome.

One suitable integrase expressing expression vector for use in the present invention is pCMV-C3 lint (SEQ ID NO: 1) as shown in Fig. 9, and described in Groth et al, Proc. Natl. Acad. Sci. U.S.A. 97: 5995-6000 (2000), incorporated herein by reference in its entirety. In pCMV-C3 lint, expression of the integrase-encoding
10 sequence is driven by the CMV promoter. However, any promoter may be used that will give expression of the integrase in a recipient cell, including operably linked avian-specific gene expression control regions of the avian ovalbumin, lysozyme, ovomucin, ovomucoid gene loci, viral gene promoters, inducible promoters, the RSV promoter and the like.

15 The recombinant nucleic acid molecules of the present invention for delivery of a heterologous polynucleotide to the genome of a recipient cell may comprise a nucleotide sequence encoding the attB attachment site of *Streptomyces ambifaciens* as described in Thorpe & Smith, Proc. Natl. Acad. Sci. U.S.A. 95: 5505-5510 (1998). The nucleic acid molecule of the present invention may further comprise an expression
20 cassette for the expression in a recipient cell of a heterologous nucleic acid encoding a desired heterologous polypeptide. Optionally, the nucleic acid molecules may also comprise a marker such as, but not limited to, a puromycin resistance gene, a luciferase gene, EGFP, and the like.

It is contemplated that the expression cassette, for introducing a desired
25 heterologous polypeptide, comprises a promoter operably linked to a nucleic acid encoding the desired polypeptide and, optionally, a polyadenylation signal sequence. Exemplary nucleic acids suitable for use in the present invention are more fully described in the examples below.

In one embodiment of the present invention, following delivery of the nucleic
30 acid molecule and a source of integrase activity into a cell population, for example, an

avian cell population, the cells are maintained under culture conditions suitable for the expression of the integrase and/or for the integrase to mediate recombination between the recombination site of the nucleic acid and recombination site in the genome of a recipient cell. When the recipient cell is cultured in vitro, such cells may be incubated
5 at 37° Celsius. For example, chicken early stage blastodermal cells may be incubated at 37° Celsius. They may then be injected into an embryo within a hard shell, which is resealed for incubation until hatching. Alternatively, the transfected cells may be maintained in in vitro culture.

In one embodiment, the present invention provides methods for the site-
10 specific insertion of a heterologous nucleic acid molecule into the nuclear genome of a cell by delivering to a target cell that has a recombination site in its nuclear genome, a source of integrase activity, a site-specific construct that has another recombination site and a polynucleotide of interest, and allowing the integrase activity to facilitate a recombination event between the two recombination sites, thereby integrating the
15 polynucleotide of interest into the nuclear genome.

(a) Expression vector nucleic acid molecules: A variety of recombinant nucleic acid expression vectors are suitable for use in the practice of the present invention. The site-specific constructs described herein can be constructed utilizing methodologies well known in the art of molecular biology (see, for example, Ausubel or Maniatis) in
20 view of the teachings of the specification. As described above, the constructs are assembled by inserting into a suitable vector backbone a recombination site such as an attP or an attB site, a polynucleotide of interest operably linked to a gene expression control region of interest and, optionally a sequence encoding a positive selection marker. Polynucleotides of interest can include, but are not limited to, expression
25 cassettes encoding a polypeptide to be expressed in the transformed cell or in a transgenic vertebrate animal derived therefrom. The site-specific constructs are typically, though not exclusively, circular and may also contain selectable markers, an origin of replication, and other elements.

Any of the vectors of the present invention may also optionally include a
30 sequence encoding a signal peptide that directs secretion of the polypeptide expressed

by the vector from the transgenic cells, for instance, from tubular gland cells of the oviduct of an avian. In one embodiment, this aspect of the invention effectively broadens the spectrum of exogenous proteins that may be deposited in the whites of avian eggs using the methods of the invention. Where an exogenous polypeptide would not otherwise be secreted, the vector bearing the coding sequence can be modified to comprise, for instance, about 60 bp encoding a signal peptide. The DNA sequence encoding the signal peptide may be inserted in the vector such that the signal peptide is located at the N-terminus of the polypeptide encoded by the vector.

The expression vectors of the present invention can comprise a transcriptional regulatory region, for example, an avian transcriptional regulatory region, for directing expression of either fusion or non-fusion proteins. With fusion vectors, a number of amino acids are usually added to the desired expressed target gene sequence such as, but not limited to, a polypeptide sequence for thioredoxin. A proteolytic cleavage site may further be introduced at a site between the target recombinant protein and the fusion sequence. Additionally, a region of amino acids such as a polymeric histidine region may be introduced to allow binding of the fusion protein to metallic ions such as nickel bonded to a solid support, for purification of the fusion protein. Once the fusion protein has been purified, the cleavage site allows the target recombinant protein to be separated from the fusion sequence. Enzymes suitable for use in cleaving the proteolytic cleavage site include, but are not limited to, Factor Xa and thrombin. Fusion expression vectors that may be useful in the present invention include pGex (Amrad Corp., Melbourne, Australia), pRIT5 (Pharmacia, Piscataway, NJ) and pMAL (New England Biolabs, Beverly, MA), that fuse glutathione S-transferase, protein A, or maltose E binding protein, respectively, to a desired target recombinant protein.

Epitope tags are short peptide sequences that are recognized by epitope specific antibodies. A fusion protein comprising a recombinant protein and an epitope tag can be simply and easily purified using an antibody bound to a chromatography resin, for example. The presence of the epitope tag furthermore allows the recombinant protein to be detected in subsequent assays, such as Western blots, without having to produce an antibody specific for the recombinant protein itself. Examples of commonly used

epitope tags include V5, glutathione-S-transferase (GST), hemagglutinin (IIA), the peptide Phe-His-His-Thr-Thr, chitin binding domain, and the like.

Exemplary gene expression control regions for use in cells such as avian cells (e.g., chicken cells) include, but are not limited to, avian specific promoters such as the chicken lysozyme, ovalbumin, or ovomucoid promoters, and the like. Particularly useful in avian systems are tissue-specific promoters such as avian oviduct promoters that allow for expression and delivery of a heterologous polypeptide to an egg white.

Viral promoters serve the same function as bacterial or eukaryotic promoters and either provide a specific RNA polymerase in trans (bacteriophage T7) or recruit cellular factors and RNA polymerase (SV40, RSV, CMV). Viral promoters can be useful as they are generally particularly strong promoters. One useful promoter for employment in avian cells is the RSV promoter.

Selection markers are valuable elements in expression vectors as they provide a means to select for growth of only those cells that contain a vector. Common selectable marker genes include those for resistance to antibiotics such as ampicillin, puromycin, tetracycline, kanamycin, bleomycin, streptomycin, hygromycin, neomycin, ZEOCIN™, and the like.

Another element useful in an expression vector is an origin of replication. Replication origins are unique DNA segments that contain multiple short repeated sequences that are recognized by multimeric origin-binding proteins and that play a key role in assembling DNA replication enzymes at the origin site. Suitable origins of replication for use in expression vectors employed herein include E. coli oriC, colEI plasmid origin, and the like.

A further useful element in an expression vector is a multiple cloning site or polylinker. Synthetic DNA encoding a series of restriction endonuclease recognition sites is inserted into a vector, for example, downstream of the promoter element. These sites are engineered for convenient cloning of DNA into the vector at a specific position.

Elements such as the foregoing can be combined to produce expression vectors suitable for use in the methods of the invention. Those of skill in the art will be able

to select and combine the elements suitable for use in their particular system in view of the teachings of the present specification.

Provided for is the stable introduction of a large DNA molecule into the cell of an avian. In one particularly useful embodiment, the large DNA molecule is a
5 chromosome. The chromosomes to be introduced into cells of an avian may be referred to herein as "artificial chromosomes"; however, the term "artificial chromosome" is not a limiting term and any useful large DNA molecule or chromosome may be employed in the present invention.

The present invention provides modified chromosomes, which are either
10 isolated chromosomes or artificial chromosomes, which function as useful vectors to shuttle transgenes or gene clusters into the genome. By delivering the modified or artificial chromosome to an isolated recipient cell, the target cell, and progeny thereof, become trisomic or transchromosomic. Typically, an additional or trisomic chromosome will not affect the subsequent development of the recipient cell and/or an
15 embryo, nor interfere with the reproductive capacity of an adult developed from such cells or embryos. The chromosome also should be stable within chicken cells. An effective method is also required to isolate a population of chromosomes for delivery into chicken embryos or early cells.

Chickens that are trisomic for microchromosome 16 have been described
20 (Miller et al, Proc. Natl. Acad. Sci. U.S.A. 93: 3958-3962 (1996); Muscarella et al, J. Cell Biol. 101: 1749-1756 (1985). In these cases, triploidy and trisomy occurred naturally, and illustrate that an extra copy of one or more of the chicken chromosomes is compatible with normal development and reproductive capacity.

The transchromosomic avians resulting from the cellular introduction of an
25 artificial chromosome typically will comprise cells which include the normal complement of chromosomes plus at least one additional chromosome. In one embodiment, about 0.001% to 100% of the cells of the avian will include an additional chromosome. In another embodiment, about 0.1% to 100% of the cells of the avian will include an additional chromosome. In another embodiment, about 5% to 100% of
30 the cells of the avian will include an additional chromosome. In another embodiment,

about 10% to 100% of the cells of the avian will include an additional chromosome. In another embodiment, about 50% to 100% of the cells of the avian will include an additional chromosome. In one particularly useful embodiment, the additional chromosome is transmitted through the germ-line of the transchromosomic avian and
5 many, for example, most (i.e., more than 50%) of the cells of the offspring avians will include the additional chromosome. The invention contemplates the introduction and propagation of any useful number of chromosomes into the cell(s) of a transgenic avian or isolated avian cells. For example, the invention contemplates one artificial chromosome or two artificial chromosomes or three artificial chromosomes stably
10 incorporated into the genome of the cell(s) of a transchromosomal avian or isolated avian cells.

Any or all tissues of the transchromosomic avian can include the artificial chromosome. In one useful embodiment, one or more cells of the oviduct of the avians include the additional chromosome. For example, tubular gland cells of the
15 oviduct may include the additional chromosome.

A number of artificial chromosomes are useful in the methods of the invention, including, for instance, a human chromosome modified to work as an artificial chromosome in a heterologous species as described, for example, for mice (Tomizuka et al, Proc. Natl. Acad. Sci. U.S.A. 97: 722-727 (2000); for cattle (Kuroiwa et al, Nat.
20 Biotechnol. 20: 889-894 (2002); a mammalian artificial chromosome used in mice (Co et al, Chromosome Res. 8: 183-191 (2000).

Examples of large nucleic acid molecules include, but are not limited to, natural chromosomes and fragments thereof, for example, chromosomes (e.g., mammalian chromosomes) and fragments thereof which retain a centromere, artificial
25 chromosome expression systems (satellite DNA-based artificial chromosomes (SATACs); see U.S. Pat. Nos. 6,025,155, issued February 15, 2000 and 6,077,697 issued June 20, 2000, the disclosures of which are incorporated herein in their entirety by reference), mammalian artificial chromosomes (MACs) (e.g., HACs), plant artificial chromosomes, insect artificial chromosomes, avian artificial chromosomes
30 and minichromosomes (see, e.g., U.S. Pat. Nos. 5,712,134 issued January 27, 1998;

5,891,691, issued April 6, 1999; 5,288,625, issued February 22, 1994; 6,743,967 issued June 1, 2004; and U.S. Patent Application Nos. 10/235,119, published June 19, 2003, the disclosure of each of these six patents and the patent application are incorporated herein in their entirety by reference). Also contemplated for use herein are YACs, BACs, bacteriophage-derived artificial chromosomes (BBPACs), cosmid or P1 derived artificial chromosomes (PACs).

As used herein, a large nucleic acid molecule such as artificial chromosomes can stably replicate and segregate alongside endogenous chromosomes in a cell. It has the capacity to act as a gene delivery vehicle by accommodating and expressing foreign genes contained therein. A mammalian artificial chromosome (MAC) refers to chromosomes that have an active mammalian centromere(s). Plant artificial chromosomes, insect artificial chromosomes and avian artificial chromosomes refer to chromosomes that include plant, insect and avian centromeres, respectively. A human artificial chromosome (HAC,) refers to chromosomes that include human centromeres. For exemplary artificial chromosomes, see, e.g., U.S. Pat. Nos. 6,025,155, issued February 15, 2000; 6,077,697, issued June 20, 2000; 5,288,625, issued February 22, 1994; 5,712,134, issued January 27, 1998; 5,695,967, issued December 9, 1997; 5,869,294, issued February 9, 1999; 5,891,691, issued April 6, 1999 and 5,721,118, issued February 24, 1998 and published International PCT application Nos., WO 97/40183, published October 30, 1997 and WO 98/08964, published March 5, 1998, the disclosure of each of these eight patents and two PCT applications are incorporated in their entirety herein by reference.

The large nucleic acid molecules (e.g., chromosomes) can include a single copy of a desired nucleic acid fragment encoding a particular nucleotide sequence, such as a gene of interest (e.g., transgene of interest), or can carry multiple copies thereof or multiple genes, different heterologous nucleotide sequences or expression cassettes or may encode one or more heterologous transcripts each encoding more than one useful protein product (for example, the transcript(s) may comprise an IRES). Any useful IRES may be employed in the invention. See, for example, US Patent No. 4,937,190, issued January 26, 1990; Nature (1988) 334:320-325; J Virol (1988)

62:3068-3072; Ccell (1992) 68:119-131; J Virol (1990) 64:4625-4631; and J Virol (1992) 66:1476-1483, the disclosures of which are incorporated in their entirety herein by reference, which disclose useful IRESs. For example, the nucleic acid molecules can carry 40 or even more copies of genes of interest. The large nucleic acid molecules can be associated with proteins, for example, chromosomal proteins, that typically function to regulate gene expression and/or participate in determining overall structure (e.g., nucleosomes).

Certain useful artificial chromosomes, such as satellite DNA-based artificial chromosomes, can include substantially all neutral non-coding sequences (heterochromatin) except for foreign heterologous, typically gene-encoding, nucleic acid (see U.S. Pat. Nos. 6,025,155, issued February 15, 2000 and 6,077,697, issued June 20, 2000 and International PCT application No. WO 97/40183, published October 30, 1997 and Lindenbaum et al Nucleic Acids Res (2004) vol 32 no. 21 e172, the disclosures of these two patents, the PCT application and the publication are incorporated in their entirety herein by reference). Foreign genes (i.e., nucleotide sequences of interest) contained in these artificial chromosomes can include, but are not limited to, nucleic acid that encodes therapeutically effective substances (e.g., therapeutic proteins such as those disclosed elsewhere herein and traceable marker proteins (reporter genes), such as fluorescent proteins, such as green, blue or red fluorescent proteins (GFP, BFP and RFP, respectively), other reporter genes, such as beta-galactosidase and proteins that confer drug resistance, such as a gene encoding hygromycin-resistance.

Preferably, the artificial chromosomes employed herein do not interfere with the host cells' processes and can be easily purified by useful purification methods such as large-scale by high-speed flow cytometry. See, for example, de Jong, G, et al. Cytometry 35: 129-33, 1999, the disclosure of which is incorporated herein in its entirety by reference. In one embodiment, flow cytometry is employed to purify chromosomes according to de Jong supra, with the exception that the Hoechst 33258 used to stain the chromosome suspension prior to flow cytometric sorting is diluted to a concentration of about 0.125 µg/ml opposed to 2.5 µg/ml. Such artificial

chromosomes are useful for the production of transchromosomic chickens produced by introduction of the chromosomes into certain cells, for example, the germline cells, of an avian. In one particularly useful embodiment of the present invention, the transchromosomic chickens are produced by microinjection of the chromosomes, for example, cytoplasmic injection of the chromosomes into avian embryos, for example, early stage embryos such as a Stage I embryos, see, for example, US Patent Application No. 10/679,034, filed October 2, 2003, the disclosure of which is incorporated in its entirety herein by reference.

In one embodiment, heterologous nucleic acid is introduced into an artificial chromosome. Any useful method to introduce the nucleic acid into the chromosome may be employed in the invention. Thereafter, the artificial chromosomes are isolated in a mixture substantially free of other chromosomes or cellular material. For example, artificial chromosomes may be isolated by flow cytometry (e.g., dual laser high-speed flow cytometer as described previously (de Jong, G, et al. Cytometry 35: 129-33, 1999). See, for example, US Patent Application Publication No. 20030113917, published June 19, 2003, the disclosure of which is incorporated in its entirety herein by reference.

In accordance with the present invention, any useful number of artificial chromosomes may be introduced into an avian cell (e.g., injected), for example, an avian germinal cell such as a cell of an ova, an embryo or a germinal disc of an avian egg. Any useful method of introducing the chromosomes into the avian cell is contemplated for use in the present invention. In addition, the invention contemplates the introduction of any useful number of chromosomes into an avian cell. For example, and without limitation, the invention contemplates the introduction of 1 to about 1,000,000 chromosomes injected per egg. In one embodiment, 1 to about 100,000 chromosomes are injected per egg. In another embodiment about 5 to about 100,000 artificial chromosomes are injected per egg. For example, about 10 to about 50,000 chromosomes may be injected per egg.

In one embodiment, there is a lower hatch rate for eggs injected with more than a certain number of chromosomes. In one embodiment, an injection of over 100,000

chromosomes reduces or brings the hatch rate to zero. In another embodiment, an injection of over 20,000 chromosomes reduces or brings the hatch rate to zero. In another embodiment, an injection of over 5,000 chromosomes reduces or brings the hatch rate to zero. In another embodiment, an injection of over 2,000 chromosomes
5 reduces or brings the hatch rate to zero. For example, an injection of over 1,000 (e.g., 550) chromosomes reduces or brings the hatch rate to zero.

For injection, any useful volume of injection buffer may be used for each injection. For example, about 1 nl to about 1 μ l may be injected, in addition, any useful concentration of chromosomes may be employed in the injection buffer. For
10 example, and without limitation, 1 to about 100,000 chromosomes per microliter may be used. In addition, any useful number of injections may be performed on each egg.

In one embodiment, a concentration of 7000-11,500 chromosomes is used per μ l of injection buffer (Monteith, D, et al. Methods Mol Biol 240: 227-242, 2004). In one embodiment, 25-100 nanoliters (nl) of injection buffer is used per injection.

Any useful avian embryos may be employed in the present invention. For example, the embryos may be collected from 24-36 week-old hens (e.g., commercial White Leghorn variety of *G. gallus*). In one embodiment, a germinal disc is injected with the chromosomes. In one embodiment, the embryo donor hens are inseminated weekly using pooled semen from roosters to produce eggs for injection. Any useful
15 method, such as methods known to those skilled in the art, may be employed to collect fertilized eggs.

Cytoplasmic injection of artificial chromosomes can be achieved by employing certain microinjection systems or assemblies. In one particularly useful embodiment, the microinjection assembly or microinjection system disclosed in US Patent
25 Application No. 09/919,143, filed July 31, 2001 (the '143 application), the disclosure of which is incorporated herein in its entirety, is employed. Use of such a cytoplasmic injection device allows for the precise delivery of chromosomes into the cytoplasm of avian embryos, for example, early stage avian embryos, e.g., Stage I embryos.

Typically, following microinjection, the embryos are transferred to the oviduct
30 of recipient hens utilizing any useful technique, such as that disclosed in Olsen, M and

Neher, B. (1948) J Exp Zool 109: 355-66 followed by incubation and hatching of the birds.

Any useful method, such as PCR, may be used to test for the production of transchromosomic avians. Typically, the identification of a transchromosomic offspring is confirmed by fluorescence in-situ hybridization (FISH) and/or DNA analysis such as Southern blot or the like. In one useful embodiment, artificial chromosomes can be used as vectors to introduce large DNA payloads, such as nucleotide sequences to be expressed heterologously in the avian to yield a desired biomolecule, of stably maintained genetic information into transgenic chickens.

10 Production of germline transchromosomic avians is confirmed by the production of transchromosomic offspring from the GObirds.

The present invention provides for the introduction of desired nucleotide sequences into a chromosome, the chromosome of which can subsequently be isolated/purified and thereafter introduced into an avian as disclosed herein.

15 A useful chromosome isolation protocol can comprise the steps of inserting a lac-operator sequence (Robinett et al J. Cell Biol. 135: 1685-1700 (1996) into an isolated chromosome and, optionally, inserting a desired transgene sequence within the same chromosome. In one embodiment, the lac operator region is a concatamer of a plurality of lac operators for the binding of multiple lac repressor molecules. Insertion

20 can be accomplished, for instance, by identifying a region of known nucleotide sequence associated with a particular avian chromosome. A recombinant DNA molecule may be constructed that comprises the identified region, a recombination site such as attB or attP and a lac-operator concatamer. The recombinant molecule is delivered to an isolated avian cell, for example, but not limited to, chicken DT40 cells

25 that have elevated homologous recombination activity compared to other avian cell lines, whereupon homologous recombination will integrate the heterologous recombination site and the lac-operator concatamer into the targeted chromosome as shown in the schema illustrated in Fig. 20. A tag-polypeptide comprising a label domain and a lac repressor domain is also delivered to the cell, for example, by

30 expression from a suitable expression vector. The nucleotide sequence coding for a

GFP-lac-repressor fusion protein (Robinett et al, J. Cell Biol. 135: 1685-1700 (1996)) may be inserted into the same chromosome as the lac-operator insert. The lac repressor sequence, however, can also be within a different chromosome. An inducible promoter may also be used to allow the expression of the GFP-lac-repressor only after chromosome is to be isolated.

Induced expression of the GFP-lac-repressor fusion protein will result in specific binding of the tag fusion polypeptide to the lac-operator sequence for identification and isolation of the genetically modified chromosome. The tagged mitotic chromosome can be isolated using, for instance, flow cytometry as described in de Jong et al Cytometry 35: 129-133 (1999) and Griffin et al Cytogenet. Cell Genet. 87: 278-281 (1999).

A tagged chromosome can also be isolated using microcell technology requiring treatment of cells with the mitotic inhibitor colcemid to induce the formation of micronuclei containing intact isolated chromosomes within the cell. Final separation of the micronuclei is then accomplished by centrifugation in cytochalasin as described by Killary & Founder in Methods Enzymol. 254: 133-152 (1995). Further purification of microcells containing only the desired tagged chromosome could be done by flow cytometry. It is contemplated, however, that alternative methods to isolate the mitotic chromosomes or microcells, including mechanical isolation or the use of laser scissors and tweezers, and the like.

The present invention envisions the employment of any useful protein-DNA binding or interaction to assist in isolating/purifying chromosomes of the invention. Such other methods in which a desired chromosome can be labeled for purposes of isolation/purification, are well known in the art including but not limited to, steroid receptor (such as the glucocorticoid receptor); site specific response element systems, see, for example, McNally et al (2000) Science 287:1262-1265; the bacteriophage lambda repressor system; and human homeobox genes. In addition, certain mutant forms of proteins which are employed in these systems (e.g., mutant proteins which bind there substrate with greater affinity than the non-mutant form of the protein) can be particularly useful for chromosome tagging (i.e., association of the chromosome

with a marker that allows distinction of the chromosome, for example, distinction from cellular components such as other chromosomes) from other and subsequent isolation/purification of the chromosomes. Furthermore the invention contemplates the use of one or more selectable markers to identify cells which contain chromosomes
5 comprising an introduced sequence of interest.

One specific embodiment of the making of a recombinant artificial chromosome can be seen in FIG. 25. In this embodiment the artificial chromosome includes a promoter which expresses a marker. Shown in FIG. 25 is an SV40 promoter, however, any useful promoter may be employed. For example, any
10 promoter which will facilitate transcription in a cell line in which the artificial chromosome is present may be employed. For example, and without limitation, the following promoters may be useful: Pol III promoters (including type 1, type 2 and type 3 Pol III promoters) such as H1 promoters, U6 promoters, tRNA promoters, RNase MPR promoters and functional portions of each of these promoters. Other
15 promoters that may be useful include, without limitation, Pol I promoters, Pol II promoters, cytomegalovirus (CMV) promoters, rous-sarcoma virus (RSV) promoters, murine leukemia virus (MLV) promoters, mouse mammary tumor virus (MMTV) promoters, ovalbumin promoters, lysozyme promoters, conalbumin promoters, ovomucoid promoters, ovomucin promoters, ovotransferrin promoters and functional
20 portions of each of these promoters.

The schematic of FIG. 25 shows a vector which includes an OMC24-IRES-EPO nucleotide sequence of interest and a marker coding sequence both contained on a vector which integrates into the artificial chromosome. FIG. 25 shows a hygromycin resistance marker being employed. However, any useful marker (e.g., antibiotic
25 resistant marker) may be used. For example, and without limitation, zeomycin resistance, neomycin resistance and blastomycin resistance markers can be used. Also shown is the use of an attP present on the artificial chromosome and an attB site present on the vector. However, any useful recombination sites and integrase may be employed such as those disclosed elsewhere herein.

In one embodiment, a useful cell line such as LMTK- containing the chromosome (A) in FIG. 25 is transfected with the vector B by standard methodologies such as lipofection. After introduction of the vector (B) into the artificial chromosome containing cell line, integration occurs, for example, between
5 integration sites such as lambda attB and attP sites, wherein the hygromycin marker is expressed in the cells which contain the recombined artificial chromosome allowing for selection of the cells. For the employment of such integration sites, integrase or an integrase encoding gene is typically also introduced into the cell. In one useful embodiment, a lambda integrase gene is used which produces an integrase protein with
10 a substitution mutation at the glutamine residue at position 174 to a lysine. This mutation removes the requirement for host factors allowing the integrase to function in cell lines. A practitioner of skill in the art will recognize that many variations to this basic recombination methodology may be employed.

It is contemplated that more than one, for example, between 1 and 100 rounds
15 of integration of a nucleotide sequence of interest into the artificial chromosome may be performed. For example, one, two, three, four or more rounds of integration may be performed. In certain useful instances of multiple insertions of nucleotide sequences of interest into the artificial chromosome, it can be advantageous to employ different selectable markers. Any useful selectable markers can be employed in the
20 case of multiple insertions, for example, and without limitation, genes which provide for resistance to hygromycin, zeomycin, neomycin and blastomycin can be used.

In one embodiment, multiple integration sites (e.g., multiple attP sites) are present in the artificial chromosome. Multiple rounds of integration can be performed to obtain insertions of more than one nucleotide sequence of interest in an artificial
25 chromosome or to obtain an artificial chromosome with multiple copies of the same nucleotide sequence of interest. After each round of integration, a different marker can be used for each round of integration. For example, the nucleotide sequence to be inserted can include a hygromycin resistance coding sequence in the first round of integration, in the second round of integration the nucleotide to be inserted can include
30 a zeomycin resistance marker coding sequence, in the third round of integration the

nucleotide sequence to be inserted can include a neomycin resistance marker coding sequence and , in the fourth round of integration the nucleotide sequence to be inserted can include a blastomycin resistance marker coding sequence. A round of integration is where a nucleotide sequence of interest is introduced into the cell containing the
5 artificial chromosome in which integration of the nucleotide sequence takes place as disclosed herein (e.g., integration into a site having a promoter proximal to the integration site which is operable to express the marker). This is merely an example of a method for integration of multiple nucleotide sequences of interest into an artificial chromosome for use as disclosed herein. The employment of other useful
10 methodologies for integration of multiple nucleotide sequences into an artificial chromosome, as are understood by a practitioner of skill in the art, is included within the scope of the invention.

In certain instances it can be useful to rescue an artificial chromosome from an avian or cultured avian cell which contains the artificial chromosome in its genome.
15 For example, certain artificial chromosomes may fragment after being introduced into an avian. During fragmentation, the artificial chromosome can be reduced substantially in size, (e.g., reduced by about 50% or about 60% or about 70% or about 80% or about 90% or more in size). However, after fragmentation the artificial chromosome can stabilize and as such is no longer susceptible to significant
20 fragmentation or degradation in avian cells. These stabilized artificial chromosomes can be particularly useful for efficiently producing transchromosomal avians because, since the chromosomes are stabilized, they do not further fragment upon introduction into an avian or avian cell.

After recovery of the artificial chromosome from cells of the transchromosomal
25 avian in which the artificial chromosome was originally introduced, the chromosome can be re-introduced into a cell line such as a CHIO cell line in which a nucleotide sequence of interest is introduced into an integration site present in the artificial chromosome, as disclosed herein, producing a recombinant stabilized artificial chromosome. In one embodiment, multiple integration sites of the same type are
30 present on the artificial chromosome thereby increasing the chance that one or more of

the integration sites remain present on the artificial chromosome after the fragmentation occurs. The recombinant stabilized chromosome can then be introduced into an avian embryo to produce a line of avians containing in their genome a stabilized artificial chromosome comprising a transgene.

5 Any useful method may be used to recover the stabilized artificial chromosomes from the bird. For example, intact chromosomes can be prepared from blood cells of the birds as is known in the art, for example, essentially as disclosed in de Jong et al. Cytometry 35: 129-133 (1999) and Griffin et al. Cytogenet. Cell Genet. 87: 278-281 (1999). After preparation, the chromosomes are flow sorted to isolate the
10 stabilized chromosome. In one embodiment, the stabilized artificial chromosome is of a size that does not allow it to be easily distinguished from the debris field in the cytometry histogram. In this instance sequence specific marker dyes can be used to tag the stabilized artificial chromosome thereby facilitating its purification. In one useful embodiment, polyamide probes can be used to tag the stabilized artificial chromosome
15 as disclosed herein.

Typically, the artificial chromosomes introduced into avians are stably maintained in the avians and are passed to offspring through the germline. In addition, artificial chromosomes can be stably maintained in avian cell lines such as chicken cell line (DT-40).

20 The invention is also useful for visualizing gene activity in avian cells as is understood by a practitioner of ordinary skill in the art (See, for example, Tsukamoto, et al (2000) Nature Cell Biology, 2:871-878).

Most non-viral methods of gene transfer rely on normal mechanisms used by eukaryotic cells for the uptake and intracellular transport of macromolecules. In
25 certain useful embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the subject transcriptional regulatory region and operably linked polypeptide-encoding nucleic acid by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, poly-lysine conjugates, and artificial viral envelopes. Modified chromosomes as described

above may be delivered to isolated avian embryonic cells for subsequent introduction to an embryo.

In a representative embodiment, a nucleic acid molecule can be entrapped in liposomes bearing positive charges on their surface (e.g., lipofectins) and (optionally) which are tagged with antibodies against cell surface antigens of the target tissue (Mizuno et al, 1992, NO Shinkei Geka 20: 547-551; PCT publication WO91/06309, published May 16, 1991; Japanese patent application 1047381, published February 21, 1989; and European patent publication EP-A-43075, published January 6, 1982, all of which are incorporated herein by reference in their entireties).

In similar fashion, the gene delivery system can comprise an antibody or cell surface ligand that is cross-linked with a gene binding agent such as polylysine (see, for example, PCT publications WO93/04701, published March 18, 1993; WO92/22635, published December 23, 1992; WO92/20316, published November 26, 1992; WO92/19749, published November 12, 1992; and WO92/06180, published April 16, 1992, the disclosures of which are incorporated herein by reference in their entireties). It will also be appreciated that effective delivery of the subject nucleic acid constructs via receptor-mediated endocytosis can be improved using agents which enhance escape of genes from the endosomal structures. For instance, whole adenovirus or fusogenic peptides of the influenza HA gene product can be used as part of the delivery system to induce efficient disruption of DNA-containing endosomes (Mulligan et al, 1993, Science 260:926-932; Wagner et al, 1992, Proc. Natl. Acad. Sci. 89:7934-7938; and Christiano et al, 1993, Proc. Natl. Acad. Sci. 90:2122-2126, all of which are incorporated herein by reference in their entireties). It is further contemplated that a recombinant nucleic acid molecule of the present invention may be delivered to a target host cell by other non-viral methods including by gene gun, microinjection, sperm-mediated transfer, or the like.

In one embodiment of the invention, an expression vector that comprises a recombination site, such as an attB site, and a region encoding a polypeptide deposited into an egg white are delivered to oviduct cells by in vivo electroporation. In this method, the luminal surface of an avian oviduct is surgically exposed. A buffered

solution of the expression vector and a source of integrase activity such as a second expression vector expressing integrase (for example, pCMV-int) is deposited on the luminal surface. Electroporation electrodes are then positioned on either side of the oviduct wall, the luminal electrode contacting the expression vector solution. After
5 electroporation, the surgical incisions are closed. The electroporation will deliver the expression vectors to some, if not all, treated recipient oviduct cells to create a tissue-specific chimeric animal. Expression of the integrase allows for the integration of the heterologous polynucleotide into the genome of recipient oviduct cells. While this method may be used with any bird, a useful recipient is a chicken due to the size of the
10 oviduct. Also useful is a transgenic bird that has a transgenic attP recombinant site in the nuclear genomes of recipient oviduct cells, thus increasing the efficiency of integration of the expression vector.

The attB/P integrase system is useful in the in vivo electroporation method to allow the formation of stable genetically transformed oviduct cells that otherwise
15 progressively lose the heterologous expression vector.

The stably modified oviduct cells will express the heterologous polynucleotide and deposit the resulting polypeptide into the egg white of a laid egg. For this purpose, the expression vector will further comprise an oviduct-specific promoter such as ovalbumin or ovomucoid operably linked to the desired heterologous
20 polynucleotide.

Another aspect of the invention is the generation of a trisomic or transchromosomic avian cell comprising a genetically modified extra chromosome. The extra chromosome may be an artificial chromosome or an isolated avian chromosome that has been genetically modified. Introduction of the extra
25 chromosome to an avian cell will generate a trisomic or transchromosomic cell with $2n+1$ chromosomes, where n is the haploid number of chromosomes of a normal avian cell.

Delivery of an isolated chromosome into an isolated avian cell or embryo can be accomplished in several ways. Isolated mitotic chromosomes or a micronucleus
30 containing an interphase chromosome can be injected into early stage I embryos by

cytoplasmic injection. The injected zygote would then be surgically transferred to a recipient hen for the production and laying of a hard shell egg. This hard shell egg would then be incubated until hatching of a chick.

5 In one embodiment, isolated microcells which contain the artificial chromosome can be fused to primordial germ cells (PGCs) isolated from the blood stream of late stage 15 embryos as described by Killary & Fournier in *Methods Enzymol.* 254: 133-152 (1995). The PGC/microcell hybrids can then be transplanted into the blood stream of a recipient embryo to produce germline chimeric chickens. (See Naito et al (1994) *Mol. Reprod. Dev.* 39: 153-161). The manipulated eggs would
10 then be incubated until hatching of the bird.

Blastodermal cells isolated from stage X embryos can be transfected with isolated mitotic chromosomes. Following in vitro transfection, the cells are transplanted back into stage X embryos as described, for example, in Etches et al, *Poult. Sci.*, 72: 882-889 (1993), and the manipulated eggs are incubated to hatching.

15 Stage X blastodermal cells can also be fused with isolated microcells and then transplanted back into to stage X embryos or fused to somatic cells to be used as nuclear donors for nuclear transfer as described by Kuroiwa et al, *Nat. Biotechnol.* 20: 889-894 (2002).

Chromosomal vectors, as described above, may be delivered to a recipient
20 avian cell by, for example, microinjection, liposomal delivery or microcell fusion.

In the methods of the invention, a site-specific integrase is introduced into an avian cell whose genome is to be modified. Methods of introducing functional proteins into cells are well known in the art. Introduction of purified integrase protein can ensure a transient presence of the protein and its activity. Thus, the lack of
25 permanence associated with most expression vectors is not expected to be detrimental.

The integrase used in the practice of the present invention can be introduced into a target cell before, concurrently with, or after the introduction of a site-specific vector. The integrase can be directly introduced into a cell as a protein, for example, by using liposomes, coated particles, or microinjection, or into the blastodermal layer
30 of an early stage avian embryo by microinjection. A source of the integrase can also

be delivered to an avian cell by introducing to the cell an mRNA encoding the integrase and which can be expressed in the recipient cell as an integrase polypeptide. Alternately, a DNA molecule encoding the integrase can be introduced into the cell using a suitable expression vector.

5 The present invention provides novel nucleic acid vectors and methods of use that allow integrases, such as phiC31 integrase, to efficiently integrate a heterologous nucleic acid into a vertebrate animal genome, for example, an avian genome. A novel finding is that the phiC31 integrase is remarkably efficient in avian cells and increases the rate of integration of heterologous nucleic acid at least 30-fold over that of random
10 integration. Furthermore, the phiC31 integrase works equally well at 37°C and 41°C, indicating that it will function in the environment of the developing avian embryo, as shown in Example 1.

 It is important to note that the present invention is not bound by any mechanism or theory of operation. For example, the mechanism by which integrase,
15 or any other substance described herein, facilitates transgenesis is unimportant. Integrase, for example, may facilitate transgenesis by mediating the integration of DNA into the genome of a recipient cell or integrase may facilitate transgenesis by facilitating the entry of the DNA into the cell or integrase may facilitate transgenesis by some other mechanism.

20 The site-specific vector components described above are useful in the construction of expression cassettes containing sequences encoding an integrase. One integrase-expressing vector useful in the methods of the invention is pCMV-C31int (SEQ ID NO: 1 as shown in Fig. 9) where the phiC31 integrase is encoded by a region under the expression control of the strong CMV promoter. Another useful promoter is
25 the RSV promoter as used in SEQ ID NO: 9 shown in Fig. 17. Expression of the integrase is typically desired to be transient. Accordingly, vectors providing transient expression of the integrase are useful. However, expression of the integrase can be regulated in other ways, for example, by placing the expression of the integrase under the control of a regulatable promoter (i.e., a promoter whose expression can be
30 selectively induced or repressed).

Delivery of the nucleic acids introduced into cells, for example, embryonic cells (e.g., avian cells), using methods of the invention may also be enhanced by mixing the nucleic acid to be introduced with a nuclear localization signal (NLS) peptide prior to introduction, for example, microinjection, of the nucleic acid. Nuclear
5 localization signal (NLS) sequences are a class of short amino acid sequences which may be exploited for cellular import of linked cargo into a nucleus. The present invention envisions the use of any useful NLS peptide, including but not limited to, the NLS peptide of SV40 virus T-antigen.

An NLS of the invention is an amino acid sequence which mediates nuclear
10 transport into the nucleus, wherein deletion of the NLS reduces transport into the nucleus. In certain embodiments, an NLS is a cationic peptide, for example, a highly cationic peptide. The present invention includes the use of any NLS sequence, including but not limited to, SV40 virus T-antigen. NLSs known in the art include, but are not limited to those discussed in Cokol et al, 2000, EMBO Reports, 1(5):411-
15 415, Boulikas, T., 1993, Crit. Rev. Eukaryot. Gene Expr., 3:193-227, Collas, P. et al, 1996, Transgenic Research, 5: 451-458, Collas and Alestrom, 1997, Biochem. Cell Biol. 75: 633-640, Collas and Alestrom, 1998, Transgenic Research, 7: 303-309, Collas and Alestrom, Mol. Reprod. Devel., 1996, 45:431-438. The disclosure of each of these references is incorporated by reference herein in its entirety.

Not to be bound by any mechanism of operation, DNA is protected and hence
20 stabilized by cationic polymers. The stability of DNA molecules in the cytoplasm of cells may be increased by mixing the DNA to be introduced, for example, microinjected with cationic polymers (for example, branched cationic polymers), such as polyethylenimine (PEI), polylysine, DEAE-dextran, starburst dendrimers, starburst
25 polyamidoamine dendrimers, and other materials that package and condense the DNA molecules (Kukowska-Latallo et al, 1996, Proc. Natl. Acad. Sci. USA 93:4897-4902).

Once the DNA molecules are delivered to the cytoplasm of cells, they migrate into the cell's endocytotic vesicles. Furthermore, migration into the cell's endosome is followed by fast inactivation of DNA within the endolysosomal compartment in
30 transfected or injected cells, both in vitro and in vivo (Godbey, W, et al 1999, Proc

Natl Acad Sci U S A 96: 5177-5181; and Lechardeur, D₅ et al 1999, Gene Ther 6: 482-497; and references cited therein). Accordingly, in certain embodiments, DNA uptake is enhanced by the receptor-mediated endocytosis pathway using transferrin-polylysine conjugates or adenoviral-mediated vesicle disruption to effect the release of DNA
5 from endosomes. However, the invention is not limited to this or any other theory or mechanism of operation referred to herein.

Buffering the endosomal pH using endosomal-scaping elements also protects DNA from degradation (Kircheis, R, et al 2001, Adv Drug Deliv Rev 53: 341-358 ; Boussif, O, et al 1995, Proc Natl Acad Sci U S A 92: 7297-7301; and Pollard, H, et al
10 1998, J Biol Chem 273: 7507-7511; and references cited therein). Thus, in certain embodiments, DNA complexes are delivered with polycations or cationic polymers that possess substantial buffering capacity below physiological pH, such as polyethylenimine, lipopolyamines and polyamidoamine polymers. In certain embodiments, DNA condensing compounds, such as the ones described above, are
15 combined with viruses (Curiel, D, et al Proc Natl Acad Sci U S A 88: 8850-8854, 1991; Wagner, E, et al Proc Natl Acad Sci U S A 89: 6099-6103, 1992 and Gotten, M, et al, 1992, Proc Natl Acad Sci U S A 89: 6094-6098), viral peptides (Wagner, E, et al 1992, Proc Natl Acad Sci U S A 89: 7934-7938; Plank, C, et al 1994, J Biol Chem 269: 12918-12924) and subunits of toxins (Uherek, C, et al, 1998, J Biol Chem 273:
20 8835-48). These materials significantly enhance the release of DNA from endosomes. In certain embodiments, viruses, viral peptides, toxins or subunits of toxins may be coupled to DNA/polylysine complexes via biochemical means or specifically by a streptavidin-biotin bridge (Wagner et al, 1992, Proc. Natl. Acad. Sci. USA 89:6099-6103; Plank et al, 1994, J. Biol Chem. 269(17):12918-12924). In other certain
25 embodiments, the virus that is complexed with the DNA may be adenovirus, retrovirus, vaccinia virus, or parvovirus. The viruses may be linked to PEI or another cationic polymer associated with the nucleic acid. In certain embodiments, the virus may be alphavirus, orthomyxovirus, or picornavirus. In certain embodiments, the virus is defective or chemically inactivated. The virus may be inactivated by short-
30 wave UV radiation or the DNA intercalator psoralen plus long-wave UV. The

adenovirus may be coupled to polylysine, either enzymatically through the action of transglutaminase or biochemically by biotinylating adenovirus and streptavidinylating the polylysine moiety. Transferrin may also be useful in combination with cationic polymers, adenoviruses and/or other materials disclosed herein to produce transgenic avians. For example, DNA complexes containing PEI₅ PEI-modified transferrin, and PEI-bound influenza peptides may be used to enhance transgenic avian production.

In other certain embodiments, complexes containing plasmid DNA, transferrin-PEI conjugates, and PEI-conjugated peptides derived from the N-terminal sequence of the influenza virus hemagglutinin subunit HA-2 may be used to produce transgenic chickens. In certain embodiments, the PEI-conjugated peptide may be an amino-terminal amino acid sequence of influenza virus hemagglutinin which may be elongated by an amphipathic helix or by carboxyl-terminal dimerization.

The present invention provides for methods of dispersing or distributing nucleic acid in a cell, for example, in an avian cell. The avian cell may be, for example, and without limitation, a cell of a stage I avian embryo, a cell of a stage II avian embryo, a cell of a stage III avian embryo, a cell of a stage IV avian embryo, a cell of a stage V avian embryo, a cell of a stage VI avian embryo, a cell of a stage VII avian embryo, a cell of a stage VIII avian embryo, a cell of a stage IX avian embryo, a cell of a stage X avian embryo, a cell of a stage XI avian embryo or a cell of a stage XII avian embryo. In one particularly useful embodiment, the avian cell is a cell of a stage X avian embryo.

In one aspect of the present invention, cationic polymers are useful to distribute, for example, homogeneously distribute, nucleic acid introduced into a cell, for example, an embryonic avian cell. The present invention contemplates the use of cationic polymers including, but not limited to, those disclosed herein.

However, substances other than cationic polymers also capable of distributing or dispersing nucleic acids in a cell are included within the scope of the present invention.

The concentration of cationic polymer used is not critical though, in one useful embodiment, enough cationic polymer is present to coat the nucleic acid to be

introduced into the avian cell. The cationic polymer may be present in an aqueous mixture with the nucleic acid to be introduced into the cell at a concentration in a range of an amount equal to about the weight of the nucleic acid to a concentration wherein the solution is saturated with cationic polymer. In one useful embodiment, the cationic polymer is present in an amount in a range of about 0.01% to about 50 %, for example, about 0.1% to about 20% (e.g., about 5%). The molecular weights of the cationic polymers can range from a molecular weight of about 1,000 to a molecular weight of about 1,000,000. In one embodiment, the molecular weight of the cationic polymers range from about 5,000 to about 100,000 for example, about 20,000 to about 30,000.

In one particularly useful aspect of the invention, procedures that are effective to facilitate the production of a transgenic avian may be combined to provide for an enhanced production of a transgenic avian wherein the enhanced production is an improved production of a transgenic avian relative to the production of a transgenic avian by only one of the procedures employed in the combination. For example, one or more of integrase activity, NLS, cationic polymer or other technique useful to enhance transgenic avian production disclosed herein can be used in the same procedure to provide for an enhanced production of transgenic avians relative to an identical procedure which does not employ all of the same techniques useful to enhance transgenic avian production.

Another aspect of the present invention is a vertebrate animal cell which has been genetically modified with a transgene vector according to the present invention and as described herein. For example, in one embodiment, the transformed cell can be a chicken early stage blastodermal cell or a genetically transformed cell line, including a sustainable cell line. The transfected cell according to the present invention may comprise a transgene stably integrated into the nuclear genome of the recipient cell, thereby replicating with the cell so that each progeny cell receives a copy of the transfected nucleic acid. A particularly useful cell line for the delivery and integration of a transgene comprises a heterologous attP site that can increase the efficiency of

integration of a polynucleotide by phiC31 integrase and, optionally, a region for expressing the integrase.

5 A retroviral vector can be used to deliver a recombination site such as an att site into the cellular genomes, such as avian genomes, since an attP or attB site is less than 300 bp. For example, the attP site can be inserted into the NLB retroviral vector, which is based on the avian leukosis virus genome. A lentiviral vector is a particularly suitable vector because lentiviral vectors can transduce non-dividing cells, so that a higher percentage of cells will have an integrated attP site.

10 The lacZ region of NLB is replaced by the attP sequence. A producer cell line would be created by transformation of, for example, the Isolde cell line capable of producing a packaged recombinant NLB-attP virus pseudo-typed with the envA envelope protein. Supernatant from the Isolde NLB-attP line is concentrated by centrifugation to produce high titer preparations of the retroviral vector that can then be used to deliver the attP site to the genome of a cell, for example, as described in
15 Example 9 below.

In one embodiment, an attP-containing line of transgenic birds are a source of attP transgenic embryos and embryonic cells. Fertile zygotes and oocytes bearing a heterologous attP site in either the maternal, paternal, or both, genomes can be used for transgenic insertion of a desired heterologous polynucleotide. A transgene vector
20 bearing an attB site, for example, would be injected into the cytoplasm along with either an integrase expression plasmid, mRNA encoding the integrase or the purified integrase protein. The oocyte or zygote is then cultured to hatch by ex ovo methods or reintroduced into a recipient hen such that the hen lays a hard shell egg the next day containing the injected egg.

25 In another example, fertile stage I to XII embryos, for example, stage VII to XII embryos, hemizygous or homozygous for the heterologous integration site, for example, the attP sequence, may be used as a source of blastodermal cells. The cells are harvested and then transfected with a transgene vector bearing a second recombination site, such as an attB site, plus a nucleotide sequence of interest along
30 with a source of integrase. The transfected cells are then injected into the subgerminal

cavity of windowed fertile eggs. The chicks that hatch will bear the nucleotide sequence of interest and the second integration site integrated into the attP site in a percentage of their somatic and germ cells. To obtain fully transgenic birds, chicks are raised to sexual maturity and those that are positive for the transgene in their semen are bred to non-transgenic mates. As disclosed herein, in certain embodiments, the cells of the invention, e.g., embryos, may include an integrase which specifically recognizes recombination sites and which is introduced into cells containing a nucleic acid construct of the invention under conditions such that the nucleic acid sequence(s) of interest will be inserted into the nuclear genome. Methods for introducing such an integrase into a cell are described herein. In some embodiments, the site-specific integrase is introduced into the cell as a polypeptide. In alternative embodiments, the site-specific integrase is introduced into the transgenic cell as a polynucleotide encoding the integrase, such as an expression cassette optionally carried on a transient expression vector, and comprising a polynucleotide encoding the recombinase.

In one embodiment, the invention is directed to methods of using a vector for site-specific integration of a heterologous nucleotide sequence into the genome of a cell, the vector comprising a circular backbone vector, a polynucleotide of interest operably linked to a promoter, and a first recombination site, wherein the genome of the cell comprises a second recombination site and recombination between the first and second recombination sites is facilitated by an integrase. In certain embodiments, the integrase facilitates recombination between a bacterial genomic recombination site (attB) and a phage genomic recombination site (attP).

In another embodiment, the invention is directed to a cell having a transformed genome comprising an integrated heterologous polynucleotide of interest whose integration, mediated by an integrase, was into a recombination site native to the cell genome and the integration created a recombination-product site comprising the polynucleotide sequence. In yet another embodiment, integration of the polynucleotide was into a recombination site not native to the cell genome, but instead into a heterologous recombination site engineered into the cell genome.

In further embodiments, the invention is directed to transgenic vertebrate

animals, such as transgenic birds, comprising a modified cell and progeny thereof as described above, as well as methods of producing the same.

For example, cells genetically modified to carry a heterologous attB or attP site by the methods of the present invention can be maintained under conditions that, for example, keep them alive but do not promote growth and/or cause the cells to differentiate or dedifferentiate. Cell culture conditions may be permissive for the action of the integrase in the cells, although regulation of the activity of the integrase may also be modulated by culture conditions (e.g., raising or lowering the temperature at which the cells are cultured).

The present invention also provides for methods of purifying artificial chromosomes. In one significant embodiment of the invention, the purified artificial chromosomes are used to produce transchromosomic animals including, but not limited to, transchromosomic avians (e.g., transchromosomic chickens). Any useful type of artificial chromosome is contemplated for use in the present invention.

In one aspect, the present invention is directed to purifying artificial chromosomes useful in producing transgenic avians (e.g., chickens) by tagging the chromosomes with a marker dye, for example, and without limitation, a fluorescent marker dye. In one particularly useful aspect of the invention, sequence specific polyamide probes are used in the tagging process. The tagged chromosomes may be purified by methods which provide for the discrimination of the tagged chromosomes over untagged chromosomes, such as flow cytometry.

For example, the method of Gygi et al (2002) *Nucleic Acids Res.* 30: 2790-2799, the disclosure of which is incorporated by reference herein in its entirety, is contemplated for use in the present invention. Briefly, the protocol provides for the use of synthetic polyamide probes to fluorescently label regions on the artificial chromosomes (e.g., heterochromatin in the case of SATACs) which are then isolated by flow cytometry. The polyamides may bind to the minor groove of DNA of the chromosomes in a sequence specific manner without the need to disrupt the chromosome (e.g., denature the DNA).

Any useful region (e.g., nucleotide sequence or sequences) of the artificial chromosome to be purified can be tagged using probes as disclosed herein. For example, any sequence present in the artificial chromosome to be purified and not present, or present to a lesser degree, in one or more chromosomes naturally occurring in the host cell may be tagged using a probe. For example, telomeric regions, centromeric regions, non-coding regions and/or coding regions of the artificial chromosome may be targeted by the probes thereby tagging the artificial chromosome. For example, the heterochromatic region of SATACs can be targeted for tagging since SATACs (or megachromosomes) (see, for example, US Patent No. 6,077,697, issued June 20, 2000) are comprised primarily of heterochromatic DNA (e.g., repeat sequences of the mouse major satellite DNA sequences). For example, fluorescent in situ hybridization utilizing probes designed to recognize mouse major satellite sequences produce an intense fluorescent signal throughout the length of the heterochromatic region of SATACs. The signal has been shown not to be present in background chromosomes of non-mouse cell lines such as ChY1 cells showing that cell lines such as ChY1 are useful to host the artificial chromosome when targeting the artificial chromosome for tagging (e.g., tagging with labeled probes).

The invention contemplates the purification of any artificial chromosome useful for the production of transchromosomic animals (e.g., transchromosomic avians) as disclosed herein. For example, artificial chromosomes and methods related thereto such as those disclosed in U S Patent No. 6,025,155 issued February 15, 2000; U S Patent No. 6,743,967 issued June 1, 2004; U S Patent No. 6,077,697 issued June 20, 2000; U S Patent No. 5,288,625 issued February 22, 1994; U S Patent No. 5,721,118 issued February 24, 1998; U S Patent No. 6,133,503 issued October 17, 2000; US patent publication 2003/0113917 published June 19, 2003; US patent publication 2003/0003435 published January 2, 2003; WO 95/32297, International Publication Date November 30, 1995 are contemplated for use in accordance with the present invention. The disclosures of each of these six US patents, two published US patent applications and one published WO patent application are incorporated in their entirety herein by reference. In addition, the following publications disclose artificial

chromosomes or methods related thereto which are contemplated for use in the present invention. Each of these publications is incorporated herein in its entirety by reference: Bower, "Constructing a fully defined human minichromosome: Cloning a centromere" (1987) *Proc. 4th Eur. Congress Biotechnol.* 3:571; Carine, et al, "Chinese hamster cells with a minichromosome containing centromere region of human chromosome 1"(1986) *Somatic Cell Molec.Genet.* 12:479—491; Carine, et al., "Molecular characterization of human minichromosomes with centromere from chromosome 1 in hamster—human hybrids"(1989) *Somatic Cell Molec. Genet.* 15(15):445—460; Fair, et al. "Generation of a human X—derived minichromosome using telomere—associated chromosome fragmentation" (1995) *EMBO J.* 14:5444—5454; Hadlaczky, et al, Centromere formation in mouse cells cotransformed with human DNA and a dominant marker gene (1991) *Proc. Natl. Acad. Sci.USA*, 88:8106-8110; Hadlaczky, et al, Satellite DNA-based artificial chromosomes for use in gene therapy (2001) *Curr. Opin. Mol. Ther.*, Apr. 3(2):125-32; Hadlaczky and Szalay, "Mammalian artificial chromosomes: Potential vectors for gene therapy" (1996) Abstract from International Symposium on Gene Therapy of Cancer, AIDS and Genetic Disorders, Trieste (Italy) (Apr, 10—13); Hadlaczky and Szalay, "Mammalian artificial chromosomes: Introduction of novel genes into mammalian artificial chromosomes"(1996) Abstract from International Symposium on Gene Therapy of Cancers AIDS and Genetic Disorders, Trieste (Italy) (Apr. 10—13,); Harrington et al., Formation of de novo centromeres and construction of first-generation human artificial microchromosomes (1997) *Nature Genetics*, 15: (4) 345-355; Heller et al. Mini—chromosomes derived from the human Y chromosome by telomere directed chromosome breakage (1996) *Proc. Natl. Acad. Sci. USA*, 93:7125—7130; Huxley, "Mammalian artificial chromosomes: a new tool for gene therapy" (1994) *Gene Therapy*, 1:7—12; Huxley, C , Mammalian artificial chromosomes and chromosome transgenics (1997) *Trends Genet.*, Sep;13(9):345-7; Keresδ, et al, De novo chromosome formations by large-scale amplification of the centromeric region of mouse chromosomes, *Chromosome Res.* (1996) Apr; 4(3):226-239; Katoh, et al, Construction of a novel human artificial chromosome vector for gene delivery (2004)

Biochem. Biophys. Res. Commun. 203:280-290; Larin, et al, Advances in human artificial chromosome technology (2002) Trends Genet., Jun;18 (6):313-9; Lipps, et al, Chromosome-based vectors for gene therapy (2003) Gene, 304:23-33; Mills, et al, Generation of an -2.4 Mb human X centromere-based minichromosome by targeted telomere-associated chromosome fragmentation in DT40 (1999) Human Molecular Genetics, 8(5) 751-761; Murray, et al., "Construction of artificial chromosomes in yeast" (1983) Nature 305:189—193; Masumoto, Structural and functional analyses of the centromere of human chromosome 21: construction of human artificial chromosomes (2001) Tanpakushitsu Kakusan Koso., Dec;46(16 Suppl):2375-8; Praznovszky, et al, De novo chromosome formation in rodent cells (1991) Proc. Natl. Acad. Sci.USA, 88:11042-11046; Raimondi, et al., "X—ray mediated size reduction, molecular characterization and transfer in model systems of a human artificial minichromosome" (1996) Abstract from International Symposium on Gene Therapy of Cancer, AIDS and Genetic Disorders, Trieste (Italy) Apr. 10—13; Robi, et al., Artificial chromosome vectors and expression of complex proteins in transgenic animals (2003) Theriogenology, Jan 1;59(1):107-13; Telenius, et al, Stability of a functional murine satellite DNA-based artificial chromosome across mammalian species (1999) Chromosome Research, 7:3-7; and Wang, et al, Expression of a Reporter Gene After Microinjection of Mammalian Artificial Chromosomes into Pronuclei of Bovine Zygotes (2001) Mol. Reprod. and Dev. 60:433-438.

In one useful embodiment of the invention, labeled (e.g., fluorescently labeled) polyamide probes are employed to tag the artificial chromosomes to facilitate purification (e.g., flow cytometry based purification) of the artificial chromosome. Polyamide probes may be prepared by any useful method known in the art such as those methods disclosed in: PCT/US97/12733; PCT/US97/03332; PCT/US97/12722; PCT/US98/06997; PCT/US98/02444; PCT/US98/02684; PCT/US98/01006; PCT/US98/03829; PCT/US98/0714 and US Patent No. 6,673,940, issued January 6, 2004. The disclosures of each of these nine PCT applications and one issued patent are incorporated in their entirety herein by reference. Other useful methods included in the following references are contemplated for use in accordance with the present

invention: US Patent No. 6,673,940, issued January 6, 2004; US Patent No. 6,555,692, issued April 29, 2003; US Patent No. 6,506,906, issued January 14, 2003; US Patent No. 6,472,537, issued October 29, 2002; US Patent No. 6,432,638, issued August 13, 2002; US Patent No. 6,303,312, issued October 16, 2001; and US Patent No. 6,143,901, issued November 7, 2000. The disclosures of each of these seven issued patents are incorporated in their entirety herein by reference. In addition, the following five publications, the disclosures of which are incorporated in their entirety herein by reference, disclose compositions and methods which are contemplated for use in the present invention: Dervan, Molecular Recognition of DNA by Small Molecules (2001) Bioorganic & Medicinal Chem. 9; 2215-2235; Gygi, et al., Use of Fluorescent Sequence-Specific Polyamides to Discriminate Human Chromosomes by Microscopy and Flow Cytometry (2002) Nucleic Acids Research; 30: (13) 2790-2799; Rucker, et al., Sequence Specific Fluorescence Detection of Double Strand DNA (2003); J. Am.Chem. Soc; 125: 1195-1202; Recognition of the DNA minor groove by pyrrole-imidazole polyamides (2003) Curr Opin Struct Biol 13: 284-99; and J Am Chem Soc (2003) 125: 1195-202.

Wade, et al. (J. Am. Chem. Soc, vol. 114:8783-8794 (1992)) reported the design of polyamides that bind in the minor groove of dsDNA at 5'-(A,T)G(A,T)C(A,T)-3' sequences by a dimeric, side-by-side motif. Mrksich, et al. (Proc. Natl. Acad. Sci. USA, vol. 89:7586-7590 (1992)), reported an antiparallel, side-by-side polyamide motif for sequence-specific recognition in the minor groove of dsDNA by the designed peptide 1-methylimidazole-2-carboxamide netropsin. Trauger, et al. (Nature, vol. 382:559-561 (1996)) reports the recognition of a targeted dsDNA by a polyamide at subnanomolar concentrations. The disclosure of each of these three publications is incorporated by reference herein in its entirety.

In one embodiment, the particular order of amino acids in polyamides useful for making labeled (e.g., fluorescently labeled) polyamide probes, and their pairing in dimeric, antiparallel complexes formed by association of two polyamide polymers can be used to determine the sequence of nucleotides in dsDNA with which the polymers preferably associate.

The development of pairing rules for minor groove binding polyamides derived from N-methylpyrrole (Py) and N-methylimidazole (Im) amino acids can provide a useful code to control target nucleotide base pair sequence specificity. Specifically, in one embodiment an Im/Py pair in adjacent polymers can distinguish G-C from C-G and both of these from A-T or T-A base pairs. A Py/Py pair can specify A-T from G-C but cannot distinguish A-T from T-A. White, et al. (Biochemistry, vol. 35:12532-12537 (1996), the disclosure of which is incorporated in its entirety herein by reference) reported the effects of the A-T/T-A degeneracy of Py/Im polyamide recognition in the minor groove of dsDNA. White, et al. (Chem. & Biol. vol. 4:569-578 (1997), the disclosure of which is incorporated in its entirety herein by reference) reported the pairing rules for recognition in the minor groove of dsDNA by Py/Im polyamides and the 5' to 3', N to C orientation preference for polyamide binding in the minor groove of dsDNA.

The inclusion of an aromatic amino acid, such as 3-hydroxy-N-methylpyrrole (Hp)(made by replacing a single hydrogen atom in Py with a hydroxy group), in a polyamide and paired opposite Py enables A-T to be discriminated from T-A by an order of magnitude. Utilizing Hp together with Py and Im in polyamides provides a code to distinguish all four Watson-Crick base pairs (i.e., A-T, T-A, G-C, and C-G) in the minor groove of dsDNA, as follows:

20

Pairing Code
for Minor Groove Recognition

	Pair	G-C	C-G	T-A	A-T
25	Im/Py	+	-	-	-
	Py/Im	-	+	-	-
	Hp/Py	-	-	+	-
	Py/Hp	-	-	-	+

30

Favored (+)
Disfavored (-)

It is understood that the method of designing and making probes use as disclosed herein is unimportant and present invention is not limited to any particular

probe or to any particular polyamide probe or method of making such probe which is used to purify artificial chromosomes for the production of transchromosomic animals, such as transchromosomic avians as disclosed herein.

One aspect of the invention are methods for generating a genetically modified cell for example, an avian cell, and progeny thereof, using a tagged chromosome. The methods may include providing an isolated modified chromosome comprising a lac operator region and a first recombination site, delivering the modified chromosome to an avian cell, thereby generating a trisomic or transchromosomic avian cell, delivering to the avian cell a source of a tagged polypeptide comprising a fluorescent domain and a lac repressor domain, delivering a source of integrase activity to the avian cell, delivering a polynucleotide comprising a second recombination site and a region encoding a polypeptide to the avian cell, maintaining the avian cell under conditions suitable for the integrase to mediate recombination between the first and second recombination sites, thereby integrating the polynucleotide into the modified chromosome and generating a genetically modified avian cell, expressing the tag polypeptide by the avian cell, allowing the tag polypeptide to bind to the modified chromosome so as to label the modified chromosome, and isolating the modified chromosome by selecting modified chromosomes having a tag polypeptide bound thereto.

In one embodiment of the invention, the second avian cell is selected from the group consisting of a stage VII-XII blastodermal cell, a stage I embryo, a stage X embryo; an isolated primordial germ cell, an isolated non-embryonic cell, and an oviduct cell.

In various embodiments, the isolated modified chromosome is an avian chromosome or an artificial chromosome.

In other embodiments of the invention, the step of providing an isolated modified chromosome comprising a lac operator region and a first recombination site comprises the steps of generating a trisomic or transchromosomic avian cell by delivering to an isolated avian cell an isolated chromosome and a polynucleotide comprising a lac operator and a second recombination site, maintaining the trisomic or

transchromosomal cell under conditions whereby the heterologous polynucleotide is integrated into the chromosome by homologous recombination, delivering to the avian cell a source of a tag polypeptide to label the chromosome, and isolating the labeled chromosome.

5 In one embodiment of the invention, the lac operator region is a concatamer of lac operators. In other embodiments of the invention, the tag polypeptide is expressed from an expression vector.

 In one embodiment of the invention, the tag polypeptide is microinjected into the cell. In various embodiments of the invention, the method of delivery of a
10 chromosome to an avian cell is selected from the group consisting of liposome delivery, microinjection, microcell, electroporation and gene gun delivery, or a combination thereof.

 In embodiments of the invention, the fluorescent domain of the tag polypeptide is GFP.

15 In one embodiment of the invention, the method further comprises the step of delivering the second avian cell to an avian embryo. The embryo may be maintained under conditions suitable for hatching as a chick.

 In one embodiment of the invention, the second avian cell is maintained under conditions suitable for the proliferation of the cell, and progeny thereof.

20 In various embodiments of the invention, the source of integrase activity is delivered to a first avian cell as a polypeptide or expressed from a polynucleotide, said polynucleotide being selected from an mRNA and an expression vector.

 In one embodiment of the invention, the tag polypeptide activity is delivered to the avian cell as a polypeptide or expressed from a polynucleotide operably linked to a
25 promoter. In another embodiment of the invention, the promoter is an inducible promoter. In yet another embodiment of the invention, the integrase is phiC31 integrase and in various embodiments of the invention, the first and second recombination sites are selected from an attB and an attP site, but wherein the first and second sites are not identical.

Other aspects of the present invention include methods of expressing a heterologous polypeptide in vertebrate cells by stably transfecting cells using site-specific integrase-mediation and a recombinant nucleic acid molecule, as described herein, and culturing the transfected cells under conditions suitable for expression of the heterologous polypeptide. In addition, the present invention includes methods of expressing a heterologous polypeptide in a transgenic vertebrate animal by producing a transgenic vertebrate animal using methods known in the field or described herein in combination with using site-specific integration of nucleic acid molecules as described herein, and exposing the animal to conditions suitable for expression of the heterologous polypeptide.

The protein of the present invention may be produced in purified form by any known conventional techniques. For example, in the case of heterologous protein production in eggs, the egg white may be homogenized and centrifuged. The supernatant may then be subjected to sequential ammonium sulfate precipitation and heat treatment. The fraction containing the protein of the present invention is subjected to gel filtration in an appropriately sized dextran or polyacrylamide column to separate the proteins. If necessary, the protein fraction may be further purified by HPLC or other methods well known in the art of protein purification.

The methods of the invention are useful for expressing nucleic acid sequences that are optimized for expression in the host cells and which encode desired polypeptides or derivatives and fragments thereof. Derivatives include, for instance, polypeptides with conservative amino acid replacements, that is, those within a family of amino acids that are related in their side chains (commonly known as acidic, basic, nonpolar, and uncharged polar amino acids). Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids and other groupings are known in the art (see, for example, "Biochemistry", 2nd ed, L. Stryer, ed., W.H. Freeman & Co., 1981). Peptides in which more than one replacement has taken place can readily be tested for activity in the same manner as derivatives with a single replacement, using conventional polypeptide activity assays (e.g. for enzymatic or ligand binding activities).

Regarding codon optimization, if the recombinant nucleic acid molecules are transfected into a recipient chicken cell, the sequence of the nucleic acid insert to be expressed can be optimized for chicken codon usage. This may be determined from the codon usage of at least one, or more than one, protein expressed in a chicken cell according to well known principles. For example, in the chicken the codon usage could be determined from the nucleic acid sequences encoding the proteins such as lysozyme, ovalbumin, ovomucin and ovotransferrin of chicken. Optimization of the sequence for codon usage can elevate the level of translation in avian eggs.

The present invention provides methods for the production of a protein by cells comprising the steps of maintaining a cell, transfecting with a first expression vector and, optionally, a second expression vector, under conditions suitable for proliferation and/or gene expression and such that an integrase will mediate site specific recombination at art sites. The expression vectors may each have a transcription unit comprising a nucleotide sequence encoding a heterologous polypeptide, wherein one polypeptide is an integrase, a transcription promoter, and a transcriptional terminator. The cells may then be maintained under conditions for the expression and production of the desired heterologous polypeptide(s).

The present invention further relates to methods for gene expression by cells, such as avian cells, from nucleic acid vectors, and transgenes derived therefrom, that include more than one polypeptide-encoding region wherein, for example, a first polypeptide-encoding region can be operatively linked to an avian promoter and a second polypeptide-encoding region is operatively linked to an Internal Ribosome Entry Sequence (IRES). It is contemplated that the first polypeptide-encoding region, the IRES and the second polypeptide-encoding region of a recombinant DNA of the present invention may be arranged linearly, with the IRES operably positioned immediately 5' of the second polypeptide-encoding region. This nucleic acid construct can be used for the production of certain proteins in vertebrate animals or in their cells. For example, when inserted into the genome of an avian cell or a bird and expressed therein, will generate individual polypeptides that may be post-translationally modified and combined in the white of a hard shell bird egg. Alternatively, the expressed

polypeptides may be isolated from an avian egg and combined in vitro.

The invention, therefore, includes methods for producing multimeric proteins including immunoglobulins, such as antibodies, and antigen binding fragments thereof. Thus, in one embodiment of the present invention, the multimeric protein is
5 an immunoglobulin, wherein the first and second heterologous polypeptides are immunoglobulin heavy and light chains respectively. Illustrative examples of this and other aspects of the present invention for the production of heterologous multimeric polypeptides in avian cells are fully disclosed in U.S. Patent Application No. 09/877,374, filed June 8, 2001, and U.S. Patent Application No. 10/251,364, filed
10 September 18, 2002, both of which are incorporated herein by reference in their entirety.

Accordingly, the invention further provides immunoglobulin and other multimeric proteins that have been produced by transgenic vertebrates including avians of the invention.

15 In various embodiments, an immunoglobulin polypeptide encoded by the transcriptional unit of at least one expression vector may be an immunoglobulin heavy chain polypeptide comprising a variable region or a variant thereof, and may further comprise a D region, a J region, a C region, or a combination thereof. An immunoglobulin polypeptide encoded by an expression vector may also be an
20 immunoglobulin light chain polypeptide comprising a variable region or a variant thereof, and may further comprise a J region and a C region. The present invention also contemplates multiple immunoglobulin regions that are derived from the same animal species, or a mixture of species including, but not only, human, mouse, rat, rabbit and chicken. In certain embodiments, the antibodies are human or humanized.

25 In other embodiments, the immunoglobulin polypeptide encoded by at least one expression vector comprises an immunoglobulin heavy chain variable region, an immunoglobulin light chain variable region, and a linker peptide thereby forming a single-chain antibody capable of selectively binding an antigen.

30 Examples of therapeutic antibodies that may be produced in methods of the invention include but are not limited to HERCEPTIN™ (Trastuzumab) (Genentech,

CA) which is a humanized anti-HER2 monoclonal antibody for the treatment of patients with metastatic breast cancer; REOPRO™ (abciximab) (Centocor) which is an anti-glycoprotein IIb/IIIa receptor on the platelets for the prevention of clot formation; ZENAPAX™ (daclizumab) (Roche Pharmaceuticals, Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection; PANOREX™ which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a murine anti-idiotypic (GD3 epitope) IgG antibody (ImClone System); IMC-C225 which is a chimeric anti-EGFR IgG antibody (ImClone System); VITAXIN™ which is a humanized anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Campath 1H/LDP-03 which is a humanized anti CD52 IgG1 antibody (Leukosite); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); RITUXAN™ which is a chimeric anti-CD20 IgG1 antibody (IDEC Pharm/Genentech, Roche/Zetenyaku); LYMPHOCIDE™ which is a humanized anti-CD22 IgG antibody (Immunomedics); ICM3 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primate anti-CD80 antibody (IDEC Pharm/Mitsubishi); ZEVALIN™ is a radiolabeled murine anti-CD20 antibody (IDEC/Schering AG); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); D2E7 is a humanized anti-TNF- α antibody (CATIBASF); CDP870 is a humanized anti-TNF- α Fab fragment (Celltech); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CDP571 is a humanized anti-TNF- α IgG4 antibody (Celltech); LDP-02 is a humanized anti- α 4 β 7 antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA™ is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-VLA-4 IgG antibody (Elan); and CAT-152 is a human anti-TGF- β 2 antibody (Cambridge Ab Tech).

The invention can be used to express, in large yields and at low cost, a wide range of desired proteins including those used as human and animal pharmaceuticals, diagnostics, and livestock feed additives. Proteins such as fusion proteins, growth hormones, cytokines, structural proteins and enzymes including human growth hormone, interferon, lysozyme, and β -casein are examples of proteins which are desirably expressed in the oviduct and deposited in eggs according to the invention. Other possible proteins to be produced include, but are not limited to, albumin, α -1 antitrypsin, antithrombin III, collagen, factors VIII, IX, X (and the like), fibrinogen, hyaluronic acid, insulin, lactoferrin, protein C, erythropoietin (EPO), granulocyte colony-stimulating factor (G-CSF), granulocyte macrophage colony-stimulating factor (GM-CSF), tissue-type plasminogen activator (tPA), feed additive enzymes, somatotropin, and chymotrypsin. Immunoglobulins (shown, for example in Example 10 below) and genetically engineered antibodies, including immunotoxins which bind to surface antigens on human tumor cells and destroy them, can also be expressed for use as pharmaceuticals or diagnostics.

Other specific examples of therapeutic proteins which are contemplated for production as disclosed herein include, with out limitation, factor VIII, b-domain deleted factor VIII, factor Vila, factor IX, anticoagulants; hirudin, alteplase, tpa, reteplase, tpa, tpa - 3 of 5 domains deleted, insulin, insulin lispro, insulin aspart, insulin glargine, long-acting insulin analogs, hgh, glucagons, tsh, follitropin-beta, fsh, gm-csf, pdgh, ifn alpa2a, inf-apha, inf-beta Ib, differs from h protein by cl7 to s, ifn-beta Ia, ifn-gammalb, il-2, il-11, hbsag, ospa, murine mab directed against t-lymphocyte antigen, murine mab directed against tag-72, tumor-associated glycoprotein, fab fragments derived from chimeric mab, directed against platelet surface receptor gpII(b)/III(a), murine mab fragment directed against tumor-associated antigen cal25, murine mab fragment directed against human carcinoembryonic antigen, cea, murine mab fragment directed against human cardiac myosin, murine mab fragment directed against tumor surface antigen psma, murine mab fragments (fab/fab2 mix) directed against hmw-maa, murine mab fragment (fab) directed against carcinoma-associated antigen, mab fragments (fab) directed against nca 90, a surface

granulocyte nonspecific cross reacting antigen, chimeric mab directed against cd20 antigen found on surface of b lymphocytes, humanized mab directed against the alpha chain of the il2 receptor, chimeric mab directed against the alpha chain of the 112 receptor, chimeric mab directed against tnfr-alpha, humanized mab directed against an epitope on the surface of respiratory syncytial virus, humanized mab directed against her 2, i.e., human epidermal growth factor receptor 2, human mab directed against cytokeratin tumor-associated antigen anti-ctla4, chimeric mab directed against cd 20 surface antigen of b lymphocytes dornase-alpha dnase, beta glucocerebrosidase, tnfr-alpha, il-2-diphtheria toxin fusion protein, tnfr-igg fragment fusion protein laronidase, dnaases, alefacept, darbepoetin alfa (colony stimulating factor), tositumomab, murine mab, alemtuzumab, rasburicase, agalsidase beta, teriparatide, parathyroid hormone derivatives, adalimumab (lggl), anakinra, biological modifier, nesiritide, human b-type natriuretic peptide (hbnp), colony stimulating factors, pegvisomant, human growth hormone receptor antagonist, recombinant activated protein c, omalizumab, immunoglobulin e (ige) blocker and lbritumomab tiuxetan.

In various embodiments of the transgenic vertebrate animal of the present invention, the expression of the transgene may be restricted to specific subsets of cells, tissues or developmental stages utilizing, for example, trans-acting factors acting on the transcriptional regulatory region operably linked to the polypeptide-encoding region of interest of the present invention and which control gene expression in the desired pattern. Tissue-specific regulatory sequences and conditional regulatory sequences can be used to control expression of the transgene in certain spatial patterns. Moreover, temporal patterns of expression can be provided by, for example, conditional recombination systems or prokaryotic transcriptional regulatory sequences.

Another aspect of the present invention provides a method for the production of a heterologous protein capable of forming an antibody suitable for selectively binding an antigen. This method comprises a step of producing a transgenic vertebrate animal incorporating at least one transgene, the transgene encoding at least one heterologous polypeptide selected from an immunoglobulin heavy chain variable region, an immunoglobulin heavy chain comprising a variable region and a constant

region, an immunoglobulin light chain variable region, an immunoglobulin light chain comprising a variable region and a constant region, and a single-chain antibody comprising two peptide-linked immunoglobulin variable regions.

5 In one embodiment of this method, the isolated heterologous protein is an antibody capable of selectively binding to an antigen and which may be generated by combining at least one immunoglobulin heavy chain variable region and at least one immunoglobulin light chain variable region, for example, cross-linked by at least one disulfide bridge. The combination of the two variable regions generates a binding site that binds an antigen using methods for antibody reconstitution that are well known in
10 the art.

The present invention also encompasses immunoglobulin heavy and light chains, or variants or derivatives thereof, to be expressed in separate transgenic avians, and thereafter isolated from separate media including serum or eggs, each isolate comprising one or more distinct species of immunoglobulin polypeptide. The method
15 may further comprise the step of combining a plurality of isolated heterologous immunoglobulin polypeptides, thereby producing an antibody capable of selectively binding to an antigen. In this embodiment, for instance, two or more individual transgenic avians may be generated wherein one transgenic produces serum or eggs having an immunoglobulin heavy chain variable region, or a polypeptide comprising such, expressed therein. A second transgenic animal, having a second transgene,
20 produces serum or eggs having an immunoglobulin light chain variable region, or a polypeptide comprising such, expressed therein. The polypeptides from two or more transgenic animals may be isolated from their respective sera and eggs and combined in vitro to generate a binding site capable of binding an antigen.

25 One aspect of the present invention, therefore, concerns transgenic vertebrate animals such as transgenic birds, for example, transgenic chickens, comprising a recombinant nucleic acid molecule and which may (though optionally) expresses a heterologous gene in one or more cells in the animal. Suitable methods for the generation of transgenic animals are known in the art and are described in, for
30 example, WO 99/19472, published April 22, 1999; WO 00/1 1151, published March 2,

2000; and WO 00/56932, published September 28, 2000, the disclosures of which are incorporated herein by reference in their entirety.

Embodiments of the methods for the production of a heterologous polypeptide by avian tissue such as oviduct tissue and the production of eggs which contain
5 heterologous protein involve providing a suitable vector and introducing the vector into embryonic blastodermal cells together with an integrase, for example, a serine recombinase such as phiC3 1 integrase, so that the vector can integrate into the avian genome. A subsequent step involves deriving a mature transgenic avian from the transgenic blastodermal cells produced in the previous steps. Deriving a mature
10 transgenic avian from the blastodermal cells optionally involves transferring the transgenic blastodermal cells to an embryo and allowing that embryo to develop fully, so that the cells become incorporated into the bird as the embryo is allowed to develop.

Another alternative may be to transfer a transfected nucleus to an enucleated
15 recipient cell which may then develop into a zygote and ultimately an adult bird. The resulting chick is then grown to maturity.

In another embodiment, the cells of a blastodermal embryo are transfected or transduced with the vector and integrase directly within the embryo. It is contemplated, for example, that the recombinant nucleic acid molecules of the present
20 invention may be introduced into a blastodermal embryo by direct microinjection of the DNA into a stage X or earlier embryo that has been removed from the oviduct. The egg is then returned to the bird for egg white deposition, shell development and laying. The resulting embryo is allowed to develop and hatch, and the chick allowed to mature.

25 In one embodiment, a transgenic bird of the present invention is produced by introducing into embryonic cells such as, for instance, isolated avian blastodermal cells, a nucleic acid construct comprising an attB recombination site capable of recombining with a pseudo-attP recombination site found within the nuclear genome of the organism from which the cell was derived, and a nucleic acid fragment of
30 interest, in a manner such that the nucleic acid fragment of interest is stably integrated

into the nuclear genome of germline cells of a mature bird and is inherited in normal Mendelian fashion. It is also within the scope of the invention that the targeted cells for receiving the transgene have been engineered to have a heterologous attP recombination site, or other recombination site, integrated into the nuclear genome of
5 the cells, thereby increasing the efficiency of recognition and recombination with a heterologous attB site.

In either case, the transgenic bird produced from the transgenic blastodermal cells is known as a "founder". Some founders can be chimeric or mosaic birds if, for example, microinjection does not deliver nucleic acid molecules to all of the
10 blastodermal cells of an embryo. Some founders will carry the transgene in the tubular gland cells in the magnum of their oviducts and will express the heterologous protein encoded by the transgene in their oviducts. If the heterologous protein contains the appropriate signal sequences, it will be secreted into the lumen of the oviduct and onto the yolk of an egg.

Some founders are germline founders. A germline founder is a founder that carries the transgene in genetic material of its germline tissue, and may also carry the transgene in oviduct magnum tubular gland cells that express the heterologous protein. Therefore, in accordance with the invention, the transgenic bird will have tubular gland cells expressing the heterologous protein and the offspring of the transgenic bird
20 will also have oviduct magnum tubular gland cells that express the selected heterologous protein. (Alternatively, the offspring express a phenotype determined by expression of the exogenous gene in a specific tissue of the avian.)

The stably modified oviduct cells will express the heterologous polynucleotide and deposit the resulting polypeptide into the egg white of a laid egg. For this
25 purpose, the expression vector will further comprise an oviduct-specific promoter such as ovalbumin or ovomucoid operably linked to the desired heterologous polynucleotide.

The invention also relates to methods of screening for cells (e.g., avian cells) in which a nucleotide sequence has been inserted. The invention provides for the
30 isolation of such cells by employing the expression of a marker coding sequence.

Cells that are contemplated for use as disclosed herein include, without limitation, germline cells which may include sperm cells, ova cells, and embryo cells. The embryos may be for example, stage I, stage II, stage III, stage IV₅ stage V, stage VI, stage VII, stage VIII, stage IX, stage X, stage XI or stage XII embryos. In one particularly useful embodiment, the cells contemplated for use include blastodermal cells.

In one embodiment, a first nucleotide sequence comprising a first recombination site, such as recombination sites disclosed elsewhere herein (e.g., an attP site), also includes a functional transcription initiation site. Any useful functional transcription initiation site may be employed. In one embodiment, a U3 promoter is employed. In one embodiment, a long terminal repeat (LTR) region of a retrovirus is employed as the transcription initiation site. For example, a LTR which includes a U3 promoter may be employed.

Examples of other useful transcription initiation sites may include, without limitation, Pol III promoters (including type 1, type 2 and type 3 Pol III promoters) such as H1 promoters, U6 promoters, tRNA promoters, RNase MPR promoters and functional portions of each of these promoters. Other promoters that may be useful in the present invention include, without limitation, Pol I promoters, Pol II promoters, cytomegalovirus (CMV) promoters, rous-sarcoma virus (RSV) promoters, avian leukemia virus (ALV) promoters, actin promoters such as beta actin promoters, murine leukemia virus (MLV) promoters, mouse mammary tumor virus (MMTV) promoters, SV40 promoters, ovalbumin promoters, lysozyme promoters, conalbumin promoters, ovomucoid promoters, ovomucin promoters, ovotransferrin promoters and functional portions of each of these promoters.

In accordance with the present methods, the first nucleotide sequence comprising the first recombination site and transcription initiation site is inserted into a genome of a cell by any useful method. For example, the first nucleotide sequence may be inserted into the genome as part of a retrovirus construct (e.g., ALV). For example, a retrovirus comprising an attP site may be transduced into the genome of the cell (Fig. 26).

The invention provides for the introduction of a second nucleotide sequence, which includes a second recombination site such as recombination sites disclosed elsewhere herein (e.g., an attB site) a nucleotide sequence of interest (denote as "transgene" in Fig. 26) and a promoterless marker coding sequence, into one or more
5 cells which include the first nucleotide sequence in their genome.

Any useful method for the introduction of the nucleotide sequences into the cells is contemplated for use herein. Exemplary delivery systems for the nucleic acids include, without limitation, liposomal derived systems, poly-lysine conjugates, protoplast fusion, microinjection and electroporation.

10 Any useful marker coding sequence may be employed in the present screening methods. For example, a bioluminescent protein coding sequence may serve as the marker coding sequence for use as disclosed herein. In one embodiment, the present invention contemplates the use of a green fluorescent protein (GFP) marker gene coding sequence. In one embodiment, antibiotic resistance is the marker.

15 In one embodiment, the marker coding sequence is positioned such that when integration occurs between the first and second recombination sites, the marker expression will be under the control of the transcription initiation site of the first nucleotide sequence and will be expressed. Cells in which integration has occurred can be identified by expression of the marker coding sequence.

20 The present invention provides for the isolation of one or more cells in which the marker coding sequence is expressed. In the case of bioluminescent markers such as GFP, the cells may be sorted and thereafter isolated using flow cytometry by methods well known in the art such as those methods disclosed in de Jong et al. Cytometry 35: 129-133 (1999) and Griffin et al. Cytogenet. Cell Genet. 87: 278-281
25 (1999). Any useful methods of cell separation or isolation are contemplated for use herein including mechanical isolation or the use of laser scissors and tweezers, and the like.

In one useful embodiment, the second nucleotide sequence is introduced into blastodermal cells which include the first nucleotide sequence in their genome. For
30 example, the blastodermal cells may comprise avian blastodermal cells isolated from

fertile embryos, such as stage VII to stage XII embryos. Blastodermal cells in which the marker coding sequence is expressed are isolated and introduced into the subgerminal cavity of fertile eggs. Suitable methods for the manipulation of avian eggs, including opening and resealing hard shell eggs are described in U.S. Patent
5 Serial Nos. 5,897,998 and 6,397,777 the disclosures of which are incorporated herein by reference in their entireties. The eggs are hatched and the chicks raised to maturity by methods well known in the field.

This description uses gene nomenclature accepted by the Cucurbit Genetics Cooperative as it appears in the Cucurbit Genetics Cooperative Report 18:85 (1995),
10 which are incorporated herein by reference in its entirety.

The disclosures of publications such as journal articles, patents, and published patent applications referred to in this application are hereby incorporated by reference in their entirety into the present application.

It will be apparent to those skilled in the art that various modifications,
15 combinations, additions, deletions and variations can be made in the present invention without departing from the scope or spirit of the invention. For instance, features illustrated or described as part of one embodiment can be used in another embodiment to yield a still further embodiment. It is intended that the present invention covers such modifications, combinations, additions, deletions and variations as come within
20 the scope of the appended claims and their equivalents.

The present invention is further illustrated by the following examples, which are provided by way of illustration and should not be construed as limiting. The contents of all references, published patents and patents cited throughout the present application are hereby incorporated by reference in their entireties.

25

Example 1: Phage phiC31 Integrase Functions in Avian Cells

(a) A luciferase vector bearing either an attB (SEQ ID NO: 2 shown in Fig. 10) or attP (SEQ ID NO: 3 shown in Fig. 11) site was cotransfected with an integrase expression vector CMV-C31int (SEQ ID NO: 1) into DF-I cells, a chicken fibroblast cell line.

The cells were passaged several times and the luciferase levels were assayed at each passage.

Cells were passaged every 3-4 days and one third of the cells were harvested and assayed for luciferase. The expression of luciferase was plotted as a percentage of the expression measured 4 days after transfection. A luciferase expression vector bearing an attP site as a control was also included.

As can be seen in Fig. 2, in the absence of integrase, luciferase expression from a vector bearing attP or attB decreased to very low levels after several days. However, luciferase levels were persistent when the luciferase vector bearing attB was cotransfected with the integrase expression vector, indicating that the luciferase vector had stably integrated into the avian genome.

(b) A drug-resistance colony formation assay was used to quantitate integration efficiency. The puromycin resistance expression vector pCMV-pur was outfitted with an attB (SEQ ID NO: 4 shown in Fig. 12) or an attP (SEQ ID NO: 5 shown in Fig. 13) sites. Puromycin resistance vectors bearing attB sites were cotransfected with phiC31 integrase or a control vector into DF-I cells. One day after transfection, puromycin was added. Puromycin resistant colonies were counted 12 days post-transfection.

In the absence of cotransfected integrase expression, few DF-I cell colonies were observed after survival selection. When integrase was co-expressed, multiple DF-I cell colonies were observed, as shown in Fig. 3. Similar to the luciferase expression experiment, the attB sequence (but not the attP sequence) was able to facilitate integration of the plasmid into the genome. Fig. 3 also shows that phiC31 integrase functions at both 37° Celsius and 41° Celsius. Integrase also functions in quail cells using the puromycin resistance assay, as shown in Fig. 4.

(c) The CMV-pur-attB vector (SEQ ID NO: 4) was also cotransfected with an enhanced green fluorescent protein (EGFP) expression vector bearing an attB site (SEQ ID NO: 6 shown in Fig. 14) into DF-I cells and the phiC31 integrase expression vector CMV-C31int (SEQ ID NO: 1). After puromycin selection for 12 days, the colonies were viewed with UV light to determine the percentage of cells that expressed EGFP. Approximately 20% of puromycin resistant colonies expressed

EGFP in all of the cells of the colony, as shown in Fig. 5, indicating that the integrase can mediate multiple integrations per cell.

(d) PhiC31 integrase promoted the integration of large transgenes into avian cells. A puromycin expression cassette comprising a CMV promoter, puromycin resistance gene, polyadenylation sequence and the attB sequence was inserted into a vector
5 containing a 12.0 kb lysozyme promoter and the human interferon $\alpha 2b$ gene (SEQ ID NO: 7 shown in Fig. 15) and into a vector containing a 10.0 kb ovomucoid promoter and the human interferon $\alpha 2b$ gene (SEQ ID NO: 8) as shown in Fig. 16.

DF-I cells were transfected with donor plasmids of varying lengths bearing a
10 puromycin resistance gene and an attB sequence in the absence or presence of an integrase expression plasmid. Puromycin was added to the culture media to kill those cells which did not contain a stably integrated copy of the puromycin resistance gene. Cells with an integrated gene formed colonies in the presence of puromycin in 7-12 days. The colonies were visualized by staining with methylene blue and the entire 60
15 mm culture dish was imaged.

PhiC3 1 integrase mediated the efficient integration of both vectors as shown in Fig. 7.

Example 2: Cell Culture Methods

20 DF-I cells were cultured in DMEM with high glucose, 10% fetal bovine serum, 2 mM L-glutamine, 100 units/ml penicillin and 100 μ g/ml streptomycin at 37° Celsius and 5% CO₂. A separate population of DF-I cells was grown at 41° Celsius. These cells were adapted to the higher temperature for one week before they were used for experiments.

25 Quail QT6 cells were cultured in FIO medium (Gibco) with 5% newborn calf serum, 1% chicken serum heat inactivated (at 55° Celsius for 45 mins), 10 units/ ml penicillin and 10 μ g/ml streptomycin at 37° Celsius and 5% CO₂.

30

Example 3: Selection and Assay Methods

(a) Puromycin selection assay: About 0.8×10^6 DF-I (chicken) or QT6 (quail) cells were plated in 60 mm dishes. The next day, the cells were transfected as follows:

10 to 50 ng of a donor plasmid and 1 to 10 μg of an Integrase-expressing
5 plasmid DNA were mixed with 150 μl of OptiMEM. 15 μl of DMRIE-C was mixed with 150 μl of OptiMEM in a separate tube, and the mixtures combined and incubated for 15 mins. at room temperature.

While the liposome/DNA complexes were forming, the cells were washed with OptiMEM and 2.5 ml of OptiMEM was added. After 15 minutes, 300 μl of the DNA-lipid mixture was added drop wise to the 2.5 ml of OptiMEM covering the cell layers.
10 The cells were incubated for 4-5 hours at either 37° Celsius or 41° Celsius, 5% CO₂. The transfection mix was replaced with 3 ml of culture media. The next day, puromycin was added to the media at a final concentration of 1 $\mu\text{g}/\text{ml}$, and the media replaced every 2 to 4 days. Puromycin resistant colonies were counted or imaged 10-
15 12 days after the addition of puromycin.

(b) Luciferase assay: Chicken DF-I or quail QT6 cells (0.8×10^6) were plated in 60 mm dishes. Cells were transfected as described above. The cells from a plate were transferred to a new 100 mm plate when the plate became confluent, typically on day 3-4, and re-passaged every 3-4 days.

20 At each time point, one-third of the cells from a plate were replated, and one-third were harvested for the luciferase assay. The cells were pelleted in an eppendorf tube and frozen at -70°C.

The cell pellet was lysed in 200 μl of lysis buffer (25 mM Tris-acetate, pH7.8, 2mM EDTA, 0.5% Triton X-100, 5% glycerol). Sample (5 μl) was assayed using the
25 Promega BrightGlo reagent system.

(c) Visualization of EGFP: EGFP expression was visualized with an inverted microscope with FITC illumination [Olympus 1X70, 100 W mercury lamp, IIQ-FITC Band Pass Emission filter cube, exciter 480/40 nm, emission 535/50 nm, 20X phase contrast objective (total magnification was $2.5 \times 10 \times 20$)].

(d) Staining of cell colonies: After colonies had formed, typically after 7-12 days of culture in puromycin medium, the cells were fixed in 2% formaldehyde, 0.2% glutaraldehyde for 15 mins, and stained in 0.2% methylene blue for 30 mins. followed by several washes with water. The plates were imaged using a standard CCD camera
5 in visible light.

Example 4: Production of Genetically Transformed Avian Cells

Avian stage X blastodermal cells are used as the cellular vector for the transgenes. Stage X embryos are collected and the cells dispersed and mixed with
10 plasmid DNA. The transgenes are then introduced to blastodermal cells via electroporation. The cells are immediately injected back into recipient embryos.

The cells are not cultured for any time period to ensure that they remain capable of contributing to the germline of resulting chimeric embryos. However, because there is no culture step, cells that bear the transgene cannot be identified.
15 Typically, only a small percentage of cells introduced to an embryo will bear a stably integrated transgene (0.01 to 1%). To increase the percentage of cells bearing a transgene, therefore, the transgene vector bears an attB site and is co-electroporated with a vector bearing the CMV promoter driving expression of the phiC31 transgene (CMV-C31int (SEQ ID NO: 1). The integrase then drives integration of the transgene
20 vector into the nuclear genome of the avian cell and increases the percentage of cells bearing a stable transgene.

(a) Preparation of avian stage X blastodermal cells:

- i) Collect fertilized eggs from Barred Rock or White leghorn chickens (Gallus gallus) or quail (Japonica coturnix) within 48 hrs. of laying;
- 25 ii) Use 70% ethanol to clean the shells;
- iii) Crack the shells and open the eggs;
- iv) Remove egg whites by transferring yolks to opposite halves of shells, repeating to remove most of the egg whites;
- v) Put egg yolks with embryo discs facing up into a 10cm petri dish;
- 30 vi) Use an absorbent tissue to gently remove egg white from the embryo discs;

- vii) Place a Whatman filter paper 1 ring over the embryos;
- viii) Use scissors to cut the membranes along the outside edge of the paper ring while gently lifting the ring/embryos with a pair of tweezers;
- ix) Insert the paper ring with the embryos at a 45 degree angle into a petri dish containing PBS-G solution at room temperature;
- 5 x) After ten embryo discs are collected, gently wash the yolks from the blastoderm discs using a Pasteur pipette under a stereo microscope;
- xi) Cut the discs by a hair ring cutter (a short piece of human hair is bent into a small loop and fastened to the narrow end of a Pasteur pipette with Parafilm);
- 10 xii) Transfer the discs to a 15 ml sterile centrifuge tube on ice;
- xiii) Place 10 to 15 embryos per tube and allow to settle to the bottom (about 5 mins.);
- xiv) Aspirate the supernatant from the tube;
- 15 xv) Add 5 mis of ice-cold PBS without Ca^{++} and Mg^{++} , and gently pipette 4 to 5 times using a 5 mis pipette;
- xvi) Incubate in ice for 5-7 mins. to allow the blastoderms to settle, and aspirate the supernatant;
- xvii) Add 3 mis of ice cold 0.05% trypsin/0.02% EDTA to each tube and gently pipette 3 to 5 times using a 5 ml pipette;
- 20 xviii) Put the tube in ice for 5 mins. and then flick the tube by finger 40 times. Repeat;
- xix) Add 0.5 mis FBS and 3-5 mis BDC medium to each tube and gently pipette 5-7 times using a 5 ml pipette;
- 25 xx) Spin at 500 rpm (RCF 57 x g) at 4° Celsius for 5 mins;
- xxi) Remove the supernatant and add 2 mis ice cold BDC medium into each tube; and
- xxii) Resuspend the cells by gently pipetting 20-25 times; and
- xxiii) Determine the cell titer by hemacytometer and ensure that about 95% of all
- 30 BDCs are single cells, and not clumped.

(b) Transfection of linearized plasmids into blastodermal cells by small scale electroporation:

- i) Centrifuge the blastodermal cell suspension from step (xxiii) above at RCF 57 x g, 4° Celsius, for 5 mins;
- 5 ii) Resuspend cells to a density of $1-3 \times 10^6$ per ml with PBS without Ca^{2+} and Mg^{2+} ;
- iii) Add linearized DNA, 1-30 μg per $1-3 \times 10^5$ blastodermal cells in an eppendorf tube at room temperature. Add equimolar molar amounts of the non-linearized transgene plasmid bearing an attB site, and an integrase expression plasmid;
- 10 iv) Incubate at room temperature for 10 mins;
- v) Aliquot 100 μl of the DNA-cell mixture to a 0.1 cm cuvette at room temperature;
- vi) Electroporate at 240 V and 25 μFD (or 100 V and 125 μFD for quail cells) using, for example, a Gene Pulser II™ (BIO-RAD).
- 15 vii) Incubate the cuvette at room temperature for 1-10 mins.
- viii) Before the electroporated cells are injected into a recipient embryo, they are transferred to a eppendorf tube at room temperature. The cuvette is washed with 350 μl of media, which is transferred to the eppendorf, spun at room temperature and re-suspended in 0.01-0.3 ml medium;
- 20 ix) Inject 1-10 μl of cell suspension into the subgerminal cavity of a non-irradiated or, for example, an irradiated (c.g., with 300-900 rads) stage X egg. Shell and shell membrane are removed and, after injection, resealed according to U.S. Patent No. 6,397,777, issued June 6, 2002, the disclosure of which is incorporated herein by reference in its entirety; and
- 25 x) The egg is then incubated to hatching.

(c) Blastodermal Cell Culture Medium:

- i) 409.5 mis DMEM with high glucose, L-glutamine, sodium pyruvate, pyridoxine hydrochloride;
- 30 ii) 5 mis Men non-essential amino acids solution, 10 mM;

- iii) 5 mis Penicillin-streptomycin 5000 U/ml each;
- iv) 5 mis L-glutamine, 200 mM;
- v) 75 mis fetal bovine serum; and
- vi) 0.5 mis β -mercaptoethanol, 11.2mM.

5

Example 5; Transfection of Stage X Embryos with attB Plasmids

(a) DNA-PEI: Twenty-five μ g of a phage phiC31 integrase expression plasmid (pCMV-int), and 25 μ g of a luciferase-expressing plasmid (p β -actin-GFP-attB) are combined in 200 μ l of 28 mM Hepes (pH 7.4). The DNA/Hepes is mixed with an
10 equal volume of PEI which has been diluted 10-fold with water. The DNA/Hepes/PEI is incubated at room temperature for 15 mins. Three to seven μ l of the complex are injected into the subgerminal cavity of windowed stage X white leghorn eggs which are then sealed and incubated as described in U.S. Patents No. 6,397,777, issued June 6, 2002. The complexes will also be incubated with blastodermal cells isolated from
15 stage X embryos which are subsequently injected into the subgerminal cavity of windowed irradiated stage X white leghorn eggs. Injected eggs are sealed and incubated as described above.

(b) Adenovirus-PEI:

Two μ g of a phage phiC31 integrase expression plasmid (pCMV-int), 2 μ g of a GFP
20 expressing plasmid (p β -actin-GFP-attB) and 2 μ g of a luciferase expressing plasmid (pGLB) were incubated with 1.2 μ l of JetPEI™ in 50 μ l of 20 mM Hepes buffer (pH7.4). After 10 mins at 25°C, 3×10^9 adenovirus particles (Ad5-Null, Qbiogene) were added and the incubation continued for an additional 10 mins. Embryos are transfected in ovo or ex ovo as described above.

25

Example 6: Stage I Cytoplasmic Injection

Production of transgenic chickens by cytoplasmic DNA injection using DNA injection directly into the germinal disk as described in Sang et al, Mol. Reprod. Dev.,
1: 98-106 (1989); Love et al, Biotechnology, 12: 60-63 (1994) incorporated herein by
30 reference in their entireties.

In the method of the present invention, fertilized ova, or stage I embryos, are isolated from euthanized hens 45 mins. to 4 hrs. after oviposition of the previous egg. Alternatively, eggs were isolated from hens whose oviducts have been fistulated according to the techniques of Gilbert & Wood-Gush, J. Reprod. Fertil., 5: 451-453 (1963) and Pancer et al, Br. Poult. Sci., 30: 953-7 (1989) incorporated herein in their entirety.

An isolated ovum was placed in dish with the germinal disk upwards. Ringer's buffer medium was then added to prevent drying of the ovum. Any suitable microinjection assembly and methods for microinjecting and reimplanting avian eggs are useful in the method of cytoplasmic injection of the present invention. A particularly suitable apparatus and method for use in the present invention is described in U.S. Patent Application Serial No: 09/919,143, published July 31, 2001, the disclosure of which is incorporated in its entirety herein by reference. The avian microinjection system described in the '143 Application allowed the loading of a DNA solution into a micropipette, followed by prompt positioning of the germinal disk under the microscope and guided injection of the DNA solution into the germinal disk. Injected embryos could then be surgically transferred to a recipient hen as described, for example, in Olsen & Neher, J. Exp. Zool., 109: 355-66 (1948) and Tanaka et al, J. Reprod. Fertil., 100: 447-449 (1994). The embryo was allowed to proceed through the natural in vivo cycle of albumin deposition and hard-shell formation. The transgenic embryo is then laid as a hard-shell egg which was incubated until hatching of the chick. Injected embryos were surgically transferred to recipient hens via the ovum transfer method of Christmann et al in PCT/US01/26723, published August 27, 2001, the disclosure of which is incorporated herein by reference in its entirety, and hard shell eggs were incubated and hatched.

Approximately 25 nl of DNA solution (about 60ng/ μ l) with either integrase mRNA or protein were injected into a germinal disc of stage I White Leghorn embryos obtained 90 minutes after oviposition of the preceding egg. Typically the concentration of integrase mRNA used was 100 ng/ μ l, and the concentration of integrase protein was 66 ng/ μ l.

To synthesize the integrase mRNA, a plasmid template encoding the integrase protein was linearized at the 3' end of the transcription unit. mRNA was synthesized, capped and a polyadenine tract added using the mMESSAGE mMACHINE T7 Ultra Kit™ (Ambion, Austin, TX). The mRNA was purified by extraction with phenol and chloroform and precipitated with isopropanol. The integrase protein was expressed in E. coli and purified as described by Thorpe et al, *Mol. Microbiol.*, 38: 232-241 (2000).

A plasmid encoding for the integrase protein is transfected into the target cells. However, since the early avian embryo transcriptionally silent until it reaches about 22,000 cells, injection of the integrase mRNA or protein was expected to result in better rates of transgenesis, as shown in the Table 1 below.

The chicks produced by this procedure were screened for the presence of the injected transgene using a high throughput PCR-based screening procedure as described in Harvey et al, *Nature Biotech.*, 20: 396-399 (2002).

Table 1: Summary of cytoplasmic injection results using different integrase strategies

Experimental group	Ovum transfers	Hard shells produced (%)	Chicks hatched (%) *	Transgenic chicks (%) ‡
No Integrase	5164	3634 (70%)	500 (14%)	58 (11.6%)
Integrase mRNA	1109	833 (75%)	115 (13.8%)	19 (16.5%)
Integrase protein	374	264 (70.6%)	47(17.8%)	16 (34%)

* : Percentages based on the number of hard shells

‡ : Percentages based on the number of hatched birds

Example 7: Characterization of phiC31 Integrase-Mediated Integration Sites in the Chicken Genome

To characterize phiC31-mediated integration into the chicken genome, a plasmid rescue method was used to isolate integrated plasmids from transfected and selected chicken fibroblasts. Plasmid pCR-XL-TOPO-CMV-pur-attB (SEQ ID NO:

10, shown in Fig. 18) does not have BamH I or Bgl II restriction sites. Genomic DNA from cells transformed with pCR-XL-TOPO-CMV-pur-attB was cut with BamH I or Bgl II (either or both of which would cut in the flanking genomic regions) and religated so that the genomic DNA surrounding the integrated plasmid would be captured into the circularized plasmid. The flanking DNA of a number of plasmids were then sequenced.

DF-I cells (chicken fibroblasts), 4×10^5 were transfected with 50 ng of pCR-XL-TOPO-CMV-pur-attB and 1 μ g of pCMV-int. The following day, the culture medium was replaced with fresh media supplemented with 1 μ g/ml puromycin. After 10 days of selection, several hundred puromycin-resistant colonies were evident. These were harvested by trypsinization, pooled, replated on 10 cm plates and grown to confluence. DNA was then extracted.

Isolated DNA was digested with BamH I and Bgl II for 2-3 hrs, extracted with phenol:chloroform:isoamyl alcohol, chloroform:isoamyl alcohol and ethanol precipitated. T4 DNA ligase was added and the reaction incubated for 1 hr at room temperature, extracted with phenol:chloroform:isoamyl alcohol and chloroform:isoamyl alcohol, and precipitated with ethanol. 5 μ l of the DNA suspended in 10 μ l of water was electroporated into 25 μ l of GeneHogs™ (Invitrogen) in an 0.1 cm cuvette using a GenePulser II (Biorad) set at 1.6 kV, 100 ohms, 25 μ F and plated on Luria Broth (LB) plates with 5 μ g/ml phleomycin (or 25 μ g/ml zeocin) and 20 μ g/ml kanamycin. Approximately 100 individual colonies were cultured, the plasmids extracted by standard miniprep techniques and digested with Xba I to identify clones with unique restriction fragments.

Thirty two plasmids were sequenced with the primer attB-for (5'-TACCGTCGACGATGTAGGTCACGGTC-3') (SEQ ID NO: 12) which allows sequencing across the crossover site of attB and into the flanking genomic sequence. All of plasmids sequenced had novel sequences inserted into the crossover site of attB, indicating that the clones were derived from plasmid that had integrated into the chicken genome via phiC31 integrase-mediated recombination.

The sequences were compared with sequences at GenBank using Basic Local Alignment Search Tool (BLAST). Most of the clones harbored sequences homologous to Gallus genomic sequences in the TRACE database.

5 **Example 8: Insertion of a Wild-Type attP Site into the Avian Genome Augments
Integrase-Mediated Integration and Transgenesis**

10 The chicken B-cell line DT40 cells (Buerstedde et al (1990) E.M.B.O. J., 9: 921-927) are useful for studying DNA integration and recombination processes (Buerstedde & Takeda (1991) Cell, 67:179-88). DT40 cells were engineered to harbor
15 a wild-type attP site isolated from the Streptomyces phage phiC31. Two independent cell lines were created by transfection of a linearized plasmid bearing an attP site linked to a CMV promoter driving the resistance gene to G418 (DT40-NLB-attP) or bearing an attP site linked to a CMV promoter driving the resistance gene for puromycin (DT40-pur-attP). The transfected cells were cultured in the presence of
20 G418 or puromycin to enrich for cells bearing an attP sequence stably integrated into the genome.

25 A super-coiled luciferase vector bearing an attB (SEQ ID NO: 2 shown in Fig. 10) was cotransfected, together with an integrase expression vector CMV-C31int (SEQ ID NO: 1) or a control, non-integrase expressing vector (CMV-BL) into wild-type DT40 cells and the stably transformed lines DT40-NLB-attP and DT40-pur-attP.

30 Cells were passaged at 5, 7 and 14 days post-transfection and about one third of the cells were harvested and assayed for luciferase. The expression of luciferase was plotted as a percentage of the expression measured 5 days after transfection. As can be seen in Fig. 21, in the absence of integrase, or in the presence of integrase but in
35 the DT40 cells lacking an inserted wild-type attP site, luciferase expression from a vector bearing attB progressively decreased to very low levels. However, luciferase levels were persistent when the luciferase vector bearing attB was cotransfected with the integrase expression vector into the attP bearing cell lines DT40-NLB-attP and DT40-pur-attP. Inclusion of an attP sequence in the avian genome augments the level

of integration efficiency beyond that afforded by the utilization of endogenous pseudo-attP sites.

Example 9: Generation of attP Transgenic Cell Line and Birds Using an NLB

5 **Vector**

The NLB-attP retroviral vector is injected into stage X chicken embryos laid by pathogen-free hens. A small hole is drilled into the egg shell of a freshly laid egg, the shell membrane is cut away and the embryo visualized by eye. With a drawn needle attached to a syringe, 1 to 10 μ l of concentrated retrovirus, approximately 2.5×10^5 IU₅ is injected into the subgerminal cavity of the embryo. The egg shell is resealed with a hot glue gun. Suitable methods for the manipulation of avian eggs, including opening and resealing hard shell eggs are described in U.S. Patent Serial Nos: 5,897,998, issued May 27, 1999 and 6,397,777, issued June 4, 2002, the disclosures of which are herein incorporated by reference in their entireties.

15 Typically, 25% of embryos hatch 21 days later. The chicks are raised to sexual maturity and semen samples are taken. Birds that have a significant level of the transgene in sperm DNA will be identified, typically by a PCR-based assay. Ten to 25% of the hatched roosters will be able to give rise to G1 transgenic offspring, 1 to 20% of which may be transgenic. DNA extracted from the blood of G1 offspring is analyzed by PCR and Southern analysis to confirm the presence of the intact transgene. Several lines of transgenic roosters, each with a unique site of attP integration, are then bred to non-transgenic hens, giving 50% of G2 transgenic offspring. Transgenic G2 hens and roosters from the same line can be bred to produce G3 offspring homozygous for the transgene. Homozygous offspring will be distinguished from hemizygous offspring by quantitative PCR. The same procedure can be used to integrate an attB or attP site into transgenic birds.

Example 10: Expression of Immunoglobulin Chain Polypeptides by Transgenic Chickens

30 Bacterial artificial chromosomes (BACs) containing a **70 kb** segment of the

chicken ovomucoid gene with the light and heavy chain cDNAs for a human monoclonal antibody inserted along with an internal ribosome entry site into the 3' untranslated region of the ovomucoid gene were equipped with the attB sequence. The heavy and light chain cDNAs were inserted into separate ovomucoid BACs such that expression of an intact monoclonal antibody requires the presence of both BACs in the nucleus.

Several hens produced by coinjection of the attB-bearing ovomucoid BACs and integrase-encoding mRNA into stage I embryos produced intact monoclonal antibodies in their egg white. One hen, which had a high level of the light chain ovomucoid BAC in her blood DNA as determined by quantitative PCR particularly expressed the light chain portion of the monoclonal antibody in the egg white at a concentration of 350 nanograms per ml, or approximately 12 µg per egg.

Example 11; Stage I Cytoplasmic Injection with Integrase Activity and PEI

Production of transgenic chickens by cytoplasmic DNA injection directly into the germinal disk was done as described in Example 6.

DNA (about 60ng/µl) which includes a transgene was placed in approximately 25 nl of aqueous solution with integrase mRNA or integrase protein and was mixed with an equal volume of PEI that had been diluted ten fold. The mixture was injected into a germinal disc of stage I White Leghorn embryos obtained about 90 minutes after oviposition of the preceding egg. Typically the concentration of integrase mRNA used was about 100 ng/µl, and the concentration of integrase protein was about 66 ng/µl. The integrase mRNA was synthesized according to Example 6.

Transgenic chicks produced by this procedure using: integrase mRNA/PEI and integrase protein/PEI showed positive results for the presence of heterologously expressed protein in the blood, semen and egg white.

Example 12; Stage I Cytoplasmic Injection with Integrase Activity and NLS

Production of transgenic chickens by cytoplasmic DNA injection directly into the germinal disk was done as described in Example 6.

DNA which includes a transgene was suspended in 0.25 M KCl and SV40 T antigen nuclear localization signal peptide (NLS peptide, amino acid sequence CGGPKKKRKVG (SEQ ID NO: 13)) was added to achieve a peptide DNA molar ratio of 100:1. The DNA (about 60ng/μl) was allowed to associate with the SV40 T antigen NLS peptide by incubating at 25 degrees C for about 15 minutes.

Integrase mRNA or integrase protein was added to approximately 25 nl of an aqueous DNA/NLS solution, typically, to produce a final concentration of integrase mRNA of about 50 ng/μl, or an integrase protein concentration of about 33 ng/μl. The mixture was injected into a germinal disc of stage I White Leghorn embryos obtained about 90 minutes after oviposition of the preceding egg. The integrase mRNA was synthesized as according to Example 6.

Transgenic chicks produced by this procedure using: integrase mRNA/NLS and integrase protein/NLS showed positive results for the presence of heterologously expressed protein in blood, semen and egg white.

15

Example 13: Dispersing of Plasmid DNA in Avian Stage I Embryos

DNA samples are Cy3 labeled with a Cy3 ULS labeling kit (Amersham Pharmacia Biotech). Briefly, plasmid DNA (1 μg) was sheared to approximately 100 to 500 bp fragments by sonication. Resulting DNA was incubated at 65°C for 15 min in Cy3 ULS labeling solution and unincorporated Cy3 dye was removed by spin column chromatography (CentriSep, Princeton Separations). The distribution of the DNA in stage I avian embryos was visualized after introduction into the stage I avian embryo. Enough high molecular weight or low molecular weight PEI was added to the DNA to coat the DNA. Typically, PEI was added to the DNA to a concentration of about 5%. Any useful volume of DNA/PEI can be used, for example about 25 nl.

Figure 22 shows an avian stage one embryo containing Cy3 labeled naked DNA. In Figure 22 it can be seen that the DNA is localized to certain areas of the embryo. Figure 23 and Figure 24 show an avian stage one embryo containing Cy3 labeled DNA coated with low molecular (22 kD) weight PEI (Figure 23) and high

molecular weight (25 kD) PEI (Figure 24). In Figures 23 and 24, it can be seen that the DNA is dispersed throughout the embryos.

These experiments show that DNA/PEI conjugates are distributed more uniformly in the cytoplasm of injected embryos when compared with naked DNA.

5

Example 14: Production of an attP Transgenic Chicken

GO transgenic chickens have been produced as described in Example 9. Several hundred stage X White Leghorn eggs were injected with the NLB-attP vector and about 50 chicks hatched. Sperm from approximately 30% of the hatched roosters
10 has been shown to be positive for the attP site. These hemizygotic chickens are used to generate transgenic G2 chickens homozygotic for the attP site.

Example 15: Cytoplasmic Injection of attP Stage I Embryos with OMC24-attB-IRES-CTLA4

15 Transgenic chickens are produced by cytoplasmic DNA injection directly into the germinal disk of eggs laid by transgenic homozygous attP chickens and fertilized with sperm from the same line of homozygous attP roosters, the line produced as described in Example 14. The cytoplasmic injections are carried out as described in U.S. Patent Application Serial No. 09/919,143, filed July 31, 2001, ('143 Application)
20 and U.S. Patent Application Serial No. 10/251,364, filed September 18, 2002. The disclosures of each of these two patent applications are incorporated herein by reference in their entirety.

Stage I embryos are isolated 45 mins. to 4 hrs. after oviposition of the previous egg. An isolated embryo is placed in a dish with the germinal disk upwards. Ringer's
25 buffer medium is added to prevent drying of the ovum. The avian microinjection system described in the '143 Application allows for the loading of DNA solution into a micropipette, followed by prompt positioning of the germinal disk under the microscope and guided injection of the DNA solution into the germinal disk.

Approximately 25 nl of a DNA solution (about 60ng/ μ l) of the 77 kb OMC24-
30 attB-IRES-CTLA4, disclosed in US Patent Application No. 10/856,218, filed May 28,

2004, the disclosure of which is incorporated in its entirety herein by reference, with either integrase mRNA or protein are injected into a germinal disc of the isolated stage I embryos. Typically, the concentration of integrase mRNA used is 100 ng/μl or the concentration of integrase protein is 66 ng/μl.

5 To synthesize the integrase mRNA, a plasmid template encoding the integrase protein is linearized at the 3' end of the transcription unit. mRNA is synthesized, capped and a polyadenine tract added using the mMESSAGING mMACHINE T7 Ultra Kit™ (Ambion, Austin, TX). The mRNA is purified by extraction with phenol and chloroform and precipitated with isopropanol. The integrase protein is expressed in E.
10 coli and purified as described by Thorpe et al, *Mol. Microbiol.*, 38: 232-241 (2000).

Injected embryos are surgically transferred to a recipient hen as described in Olsen & Neher, *J. Exp. Zool.*, 109: 355-66 (1948) and Tanaka et al, *J. Reprod. Fertil.*, 100: 447-449 (1994). The embryo is allowed to proceed through the natural in vivo cycle of albumin deposition and hard-shell formation. The transgenic embryo is then
15 laid as a hard-shell egg which is incubated until hatching of the chick. Injected embryos are surgically transferred to recipient hens via the ovum transfer method of Christmann et al in PCT/US01/26723, published August 27, 2001, the disclosure of which is incorporated by reference in its entirety, and hard shell eggs are incubated and hatched.

20 The chicks produced by this procedure are screened for the presence of the injected transgene using a high throughput PCR-based screening procedure as described in Harvey et al, *Nature Biotech.*, 20: 396-399 (2002). Approximately 20% of the chicks are positive for the transgene. Eggs from each of the mature hens carrying the transgene are positive for CTLA4.

25

Example 16: Cytoplasmic Injection of attP Stage I Chicken Embryos with OMIO-attB-CTLA4

Transgenic chickens are produced by cytoplasmic DNA injection directly into the germinal disk of eggs laid by transgenic homozygous attP chickens and fertilized

with sperm from the same line of homozygous attP roosters essentially as described in Example 15.

Approximately 25 nl of a 60ng/ μ l DNA solution of the OMC24-attB-IRES-CTLA4 construct of Example 15 with the OMC24 70 kb ovomucoid gene expression controlling region and IRES of the construct replaced with the 10 kb ovomucoid gene expression controlling region of pBS-OVMUP-10, also disclosed in US Patent Application No. 10/856,218, filed May 28, 2004, is injected into a fertilized germinal disc of stage I embryos along with and integrase protein. The concentration of integrase protein used is 66 ng/ μ l.

Injected embryos are then surgically transferred to a recipient hen, hard shell eggs are produced, incubated and hatched. Approximately 30% of the chicks are positive for the transgene. Eggs from each of the mature hens carrying the transgene are positive for CTLA4.

15 **Example 17: Production of attP Transgenic Quail Using an NLB vector**

The NLB-attP retroviral vector is injected into stage X quail embryos laid by pathogen-free quail. A small hole is drilled into the egg shell of a freshly laid egg, the shell membrane cut away and the embryo visualized by eye. With a drawn needle attached to a syringe, 1 to 10 μ l of concentrated retrovirus, approximately 1.0×10^5 IU, is injected into the subgerminal cavity of the embryo. The egg shell is resealed with a hot glue gun.

Typically, 25% of embryos hatch. The chicks are raised to sexual maturity and semen samples are taken. Birds that have a significant level of the transgene in their sperm DNA will be identified, typically by a PCR-based assay. Of the hatched G0 male quail, about 1% to about 20% are transgenic. The transgenic G0 quail are bred to nontransgenic quail to produce hemizygotic G1 offspring. DNA extracted from the blood of G1 offspring is analyzed by PCR and Southern analysis to confirm the presence of the intact transgene. Several lines of hemizygotic transgenic male quail, each with a unique site of attP integration, are then bred to non-transgenic quail giving G2 offspring, 50% of which are transgenic. Transgenic G2 male and female from the

same line are then bred to produce G3 offspring homozygous for the transgene. Homozygous offspring are distinguished from hemizygous offspring by quantitative PCR.

5 **Example 18: Cytoplasmic Injection of attP Stage I Quail Embryos with OMC24-attB-IRES-G-CSF**

Transgenic quail are produced by cytoplasmic DNA injection directly into the germinal disk of eggs laid by fully transgenic homozygous attP quail produced as described in Example 17. The cytoplasmic injections are carried out essentially as
10 described in the '143 Application and U.S. Patent Application Serial No. 10/251,364, filed September 18, 2002.

Stage I embryos from homozygous attP quail fertilized with sperm from a homozygous attP quail are isolated approximately 90 minutes after oviposition of the previous egg. An isolated embryo is placed in a dish with the germinal disk upwards.
15 Ringer's buffer medium is added to prevent drying of the ovum. The avian microinjection system described in the '143 Application is used to inject approximately 25 nl of a DNA solution (about 60ng/μl) of OMC24-attB-IRES-CTLA4, with the CTLA coding sequence replaced with the coding sequence for a human-granulocyte colony stimulating factor, and integrase protein into the germinal
20 disc of the stage I quail embryos. The concentration of integrase protein used is 66 ng/μl.

Injected embryos are surgically transferred to a recipient quail essentially as described in Olsen & Neher, J. Exp. Zool., 109: 355-66 (1948) and Tanaka et al, J. Reprod. Fertil., 100: 447-449 (1994). The embryo is allowed to proceed through the
25 natural in vivo cycle of albumin deposition and hard-shell formation. The transgenic embryo is then laid as a hard-shell egg which is incubated until hatching of the chick.

The chicks produced by this procedure are screened for the presence of the injected transgene using a high throughput PCR-based screening procedure as described in Harvey et al, Nature Biotech., 20: 396-399 (2002). Approximately 20%

of the chicks are positive for the transgene. Eggs from each of the mature female quail carrying the transgene are positive for G-CSF.

Example 19: Generation of attP Transgenic Duck Using an NLB vector

5 The NLB-attP retroviral vector is injected into stage X Duck embryos laid by pathogen-free Ducks. A small hole is drilled into the egg shell of a freshly laid egg, the shell membrane cut away and the embryo visualized by eye. About 1 to 10 µl of concentrated retrovirus, approximately 2.5×10^5 IU, is injected into the subgerminal cavity of the embryo. The egg shell is resealed with a hot glue gun.

10 Homozygous G3 offspring are obtained essentially as described in Example 17 for quail.

Example 20: Stage I Cytoplasmic Injection of attP Stage I Duck Embryos with OM24-attB-IRES-CTLA4

15 Transgenic ducks are produced by cytoplasmic DNA injection directly into the germinal disk of eggs laid by homozygous attP ducks fertilized with sperm from homozygous attP ducks. The injection of the stage I embryos is carried out essentially as described in the '143 Application and U.S. Patent Application Serial No. 10/251,364, filed September 18, 2002. Approximately 25 nl of a DNA solution (about
20 60ng/µl) of OMC24-attB-IRES-CTLA4, with the CTLA4 coding region replaced with a coding sequence for human erythropoietin, and integrase encoding mRNA and protein is injected into the germinal disc of the stage I embryos. The concentration of integrase mRNA used is 100 ng/µl. The injected embryos are surgically transferred to a recipient duck and the embryo is allowed to proceed through the natural in vivo cycle
25 of albumin deposition and hard-shell formation. The transgenic embryo is laid as a hard-shell egg which is incubated until hatching and the chicks are screened for the presence of the injected transgene. Approximately 20% of the chicks are positive for the transgene. Eggs from each of the mature female ducks carrying the transgene are positive for erythropoietin.

30

Example 21: Production of Transchromosomic Chickens Using Satellite DNA-Based Artificial Chromosomes

Satellite DNA-based artificial chromosomes (ACEs, as described in Lindenbaum et al Nucleic Acids Res (2004) vol 32 no. 21 e172) were isolated by a
5 dual laser high-speed flow cytometer as described previously (de Jong, G, et al. Cytometry 35: 129-133, 1999).

The flow-sorted chromosomes were pelleted by centrifugation of a 750µl sample containing approximately 10^6 chromosomes at 2500 x g for 30 min at 4°C. The supernatant, except the bottom 30 microliters (µl) containing the chromosomes,
10 was removed resulting in a concentration of about 7000 to 11,500 chromosomes per µl of injection buffer (Monteith, et al. Methods Mol Biol 240: 227-242, 2004). Depending on the number of chromosomes to be injected, 25-100 nanoliters (nl) of injection buffer was injected per embryo.

Embryos for this study were collected from 24-36 week-old hens from
15 commercial White Leghorn variety of *G. gallus*. Embryo donor hens were inseminated weekly using pooled semen from roosters of the same breed to produce eggs for injection.

On the day of egg collection, fertile hens were euthanized 2h post oviposition by cervical dislocation. Typically, oviposition is followed by ovulation of the next egg
20 after about 24 minutes (Morris, Poultry Science 52: 423-445, 1973). The recently ovulated and fertilized eggs were collected from the upper magnum region of the oviduct under sterile conditions and placed in a glass well and covered with Ringers' Medium (Tanaka, et al. J Reprod Fertil 100: 447-449, 1994) and maintained at 41°C until microinjection.

25 Cytoplasmic injection of artificial chromosomes was achieved using the microinjection apparatus disclosed in US Patent Application No. 09/919,143, filed July 31, 2001. Chromosomes were injected into the Stage I embryos at a single site. Each embryo was cytoplasmically injected with approximately: 175, 250, 350, 450, 550, 800 or >1000 chromosomes. The chromosomes were injected in a suspension of
30 25-100 nanoliters (nl) of injection buffer.

Following microinjection, the embryos were transferred to the oviduct of recipient hens using an optimized ovum transfer (OT) procedure (Olsen, M and Neher, B. J Exp Zool 109: 355-66, 1948), with the exception that the hens were anesthetized by Isofluorane gas. Typically, about 26h after OT, the recipient hens lay a hard shell egg containing the manipulated ovum. Eggs were incubated for 21 days in a regular incubator until hatching of the birds.

The chromosomes were injected into the embryos over a 9 day period. The chromosomes were divided into three batches for delivery to the embryos each batch being injected over a three day period. Chromosomes were introduced into the embryos by a single injection using the microinjection assembly disclosed in the '143 patent application. Following injection, each egg was transferred to a recipient hen. A total of 301 transfers were performed, resulting in 226 (75%) hard shells and 87 hatched chicks (38%, see Table 2).

Table 2: Hatching of embryos microinjected with satellite DNA-based artificial chromosomes.

	Ovum transfers	Hard shells produced	hatched birds
1 st batch	71	53	15
2 nd batch	113	80	33
3 rd batch	117	93	39
Totals	301	226 (75%)	87 (38%)

Previous experiments have determined that hatching is not significantly affected when embryos were injected with up to 100nl of injection buffer. Satellite DNA-based artificial chromosomes were injected in suspensions of between 25-100nl of injection buffer.

As discussed, the embryos were injected with one of seven different numbers of artificial chromosomes. There was shown to be a correlation between the number of chromosomes injected per egg and the hatch rate. AU transchromosomic birds in the present study were obtained from embryos injected with 550 chromosomes or less

(see Table 3). There was no significant difference in the hatching rates observed between the experimental groups (batches I₅2 and 3).

Six transchromosomic founders were produced based on two separate PCR analysis (6.8%, see Table 3) using primers which anneal to the puromycin resistance gene (about 75 copies of the pur^R gene are present on the chromosome. All positive birds appear normal.

Table 3: Effect of the number of Chromosomes injected per embryo on hatching and number of transchromosomic birds produced.

# chromosomes injected per embryo	# of hard shells	# chicks hatched	# of positive birds (bird tag #)
175	31	11 (35%)	3 (BB7478, BB7483, BB7515)
250	51	25 (49%)	1 (BB 7499)
350	15	6 (40%)	0
450	31	11 (35%)	0
550	39	17 (43%)	2 (BB7477, BB7523)
800	26	5 (19%) *	0
1000	33	10 (30%) *	0
Totals	226	87 (38%)	6 (6.8%)

10 *: hatching rates of embryos injected with >550 chromosomes was significantly lower (p < 0.025)

To confirm the PCR results, erythrocytes from all PCR-positive birds as well as fibroblast cells derived from skin biopsies of 5 PCR-positive birds were analyzed by interphase and metaphase FISH using a mouse-specific major satellite DNA probe (Co, et al. Chromosome Res 8: 183-191, 2000). Five of the six chicks (5.3% out of total number of chicks analyzed) tested by FISH were positive in at least one cell type (see Table 4) at 3 weeks of age. FISH analysis of erythrocytes was repeated when the birds reached 8 weeks of age and had tripled their body weight. Similar numbers of artificial chromosome-positive cells found in each bird were observed in this second FISH analysis.

25

Table 4: Summary of FISH analysis of Red Blood Cells (RBCs) and fibroblast cells derived from transchromosomic birds. Fibroblast cells from hen # 7515 were not available for analysis.

Bird #	Sex of Bird	% of artificial chromosome positive RBCs by FISH	% of artificial chromosome positive fibroblasts by FISH
BB7499	Female	77%	87%
BB7483	Female	0.8%	0%
BB7477	Male	3 %	2.8%
BB7478	Male	15%	3%
BB7515	Female	1.3%	NA
BB7523	Male	0 %	0%
Neg. control	-	0 %	0%

5

To verify the chromosomes were intact, metaphase spreads from fibroblast cells derived from founders were made as described previously (Garside and Hillman (1985) *Experientia* 41: 1183-1184). FISH analysis of metaphase spreads using the major satellite DNA probe showed the artificial chromosomes appear intact, with no apparent fragmentation or translocation onto the chicken's chromosomes. FISH analysis using a mouse minor satellite probe, which detects the centromeric region of the introduced chromosomes (Wong and Rattner (1988) *J. Nucleic Acids Res* 16: 11645-11661), demonstrated the centromere of the chromosomes was intact. Furthermore, the percentage of satellite DNA-based artificial chromosomes -positive cells from metaphase spreads agreed closely to those observed in interphase FISH.

Analysis of G1 embryos from test bird BB7499 has shown the artificial chromosome to be transmitted through the germline. In addition, sperm from BB7499 was shown to test positive for the artificial chromosome which will also provide for germline transmission of the artificial chromosome.

Example 22: Production of EPO and G-CSF Vectors for the Production of Transchromosomal Chickens

Two vectors were constructed for introduction into Satellite DNA-based artificial chromosomes. 1OMC24-IRES1-EPO - CliromattB was constructed by inserting an EPO coding sequence into an OMC24-IRES BAC clone disclosed in US Patent Application No. 10/856,218, filed May 28, 2004, the disclosure of which is incorporated in its entirety herein by reference. The EPO coding sequence was inserted in the clone so as to be under the control of the ovomucoid promoter. That is, the EPO coding sequence was inserted in place of the LC portion of OMC-IRES-LC. An attB site and a hygromycin^R coding sequence were also inserted into the vector in such a manner as to facilitate recombination into an attP site in a SATAC artificial chromosome (i.e., ACE), as see in FIG. 25. The attP site in the SATAC is located adjacent to an SV40 promoter which provides for expression of the hygromycin^R coding sequence upon integration of the vector into the attP site allowing for selection of cells containing a recombinant artificial chromosome (see, for example, US Patent No. 6,743,967, issued June 1, 2004; US Patent No. 6,025,155, issued February 15, 2000 and Lindenbaum et al Nucleic Acids Res (2004) vol 32 no. 21 e172 (see FIG. 25), the disclosure of each of these two patents and the publication are incorporated in their entirety herein by reference).

A coding sequence for G-CSF, which was codon optimized for expression in chicken tubular gland cells, was inserted in the 1OMC24-IRES1-EPO - ChromattB construct in place of the EPO coding sequence to produce 1OMC24-IRES-GCSF - ChromattB.

Example 23: Production of Erythropoietin and G-CSF Using Artificial Chromosomes in Chickens

Cells containing the recombinant artificial chromosome are produced and identified as described in Lindenbaum et al Nucleic Acids Res (2004) vol 32 no. 21 e172. Briefly, 2.5 µg of 1OMC24-IRES1-EPO ChromattB and 2.5 µg of an expression vector which contains a lambda integrase gene (int) having a codon

mutation at position 174 to substitute a lysine for a glutamine (pCXLamROK, see Lindenbaum et al Nucleic Acids Res (2004) vol 32 no. 21 e172) are transfected by standard lipofection methodologies into LMTK- cells which contain the platform SATAC (ACE) (A of FIG. 25). DNA comprising the nucleotide sequence of interest, in this case 1OMC24-IRES1-EPO ChromattB, that has been highly purified, for example, utilizing a CsCl gradient centrifugation as is well known in the art, is particularly useful, though not required. Hygromycin resistant cells clones are identified by standard antibiotic selection methodologies.

Recombinant chromosomes are prepared from the cells and isolated by flow cytometry. The substantially purified artificial chromosomes are introduced into chickens by microinjection into stage one embryos as disclosed in US Patent Application Nos. 10/679,034, filed October 2, 2003 and 09/919,143, filed July 31, 2001. Resulting chimeric germline transchromosomal avians can be identified by any useful method such as Southern blot analysis.

Example 24; Production of a Monoclonal Antibody Using Drosophila Artificial Chromosomes in Turkey

Artificial chromosomes comprising a Drosophila chromosome centromere (DAC) are prepared essentially using methods described in US Patent No. 6,025,155, issued February 15, 2000, the disclosure of which is incorporated in its entirety herein by reference.

An attB site and a hygromycin^R coding sequence are inserted into the OMC24-IRES-LC and OMC24-IRES-HC vectors disclosed in US Patent Application No. 10/856,218, filed July 31, 2001, the disclosure of which is incorporated in its entirety herein by reference, which are then each cloned into a DAC essentially as described in Examples 22 and 23. The recombinant DACs are prepared and then isolated by a dual laser high-speed flow cytometer.

The flow-sorted chromosomes are pelleted by centrifugation and are diluted to a concentration of about 7000-12,000 chromosomes per μ l of injection buffer. Approximately 50 nanoliters (nl) of injection buffer is injected per turkey embryo.

Embryos for this study are collected from actively laying commercial turkeys. Embryo donor turkeys are inseminated weekly using pooled semen from male turkeys of the same breed to produce eggs for injection.

On the day of egg collection, fertile hens are euthanized 2h post oviposition by cervical dislocation. The recently ovulated and fertilized eggs are collected from the upper magnum region of the oviduct under sterile conditions and placed in a glass well and covered with Ringers' Medium and maintained at about 40°C until microinjection.

Cytoplasmic injection of artificial chromosomes containing the OMC24-IRES-LC is achieved using the microinjection apparatus disclosed in US Patent Application No. 09/919,143. Approximately 500 chromosomes are injected into the Stage I embryos at a single site.

Following microinjection, the embryos are transferred to the oviduct of recipient turkeys essentially as described in Olsen et al, B. J Exp Zool 109: 355-66, 1948. Typically, about one day after OT, the recipient turkeys lay a hard shell egg containing the manipulated ovum. Eggs are incubated in an incubator until hatching of the birds.

G2 transchromosomal turkeys are obtained which contain the artificial chromosome in their genome. The artificial chromosome containing the OMC24-IRES-HC is introduced into embryos obtained from the G2 turkeys in essentially the same manner as described for the OMC24-IRES-LC.

Eggs from G1 transchromosomal turkeys which contain both the OMC-IRES-LC and OMC24-IRES-HC containing chromosomes in their genome are tested for the presence of intact functional monoclonal antibody. A Costar flat 96-well plate is coated with 100 µl of C Goat-anti-Human kappa at a concentration of 5 µg/ml in PBS. The plate is incubated at 37 °C for two hours. 200 µl of 5% PBA is added to the wells followed by an incubation at 37 °C for about 60-90 minutes followed by a wash. 100 µl of egg white samples (diluted in 1% PBA:LBP) is added to each well and the plate is incubated at 37 °C for about 60-90 min followed by a wash. 100 µl of a 1:2000 dilution of F'2 Goat anti-Human IgG Fc-AP in 1% PBA is added to the wells and the plate is incubated at 37 °C for 60-90 min followed by a wash. The antibody is

detected by placing 75 µl of 1mg/ml PNPP (p-nitrophenyl phosphate) in 5x developing buffer in each well and incubating for about 10-30 mins at room temperature. The detection reaction is stopped using 75ul of 1N NaOH. The egg white tests positive for significant levels of the antibody.

5

Example 25: Production of Interferon Using Avian Artificial Chromosomes in Quail

Artificial chromosomes comprising a chicken (Barred-Rock) chromosome centromere (CAC) are prepared essentially using methods described in US Patent No. 6,743,967, issued June 1, 2004, the disclosure of which is incorporated in its entirety herein by reference.

A coding sequence for interferon alpha 2b disclosed in US Patent Application No. 10/463,980, filed June 17, 2003, the disclosure of which is incorporated in its entirety herein by reference, is inserted in the 1OMC24-IRES1-Epo - ChromattB construct disclosed herein in Example 22 in place of the EPO coding sequence to produce 1OMC24-IRES-INF - ChromattB. The 1OMC24-IRES-INF - ChromattB is cloned into the CACs essentially as described in Example 23. The recombinant CACs are prepared then isolated by a dual laser high-speed flow cytometer.

The flow-sorted chromosomes are pelleted by centrifugation and are diluted to a concentration of about 10,000 chromosomes per µl of injection buffer. Approximately 50 nanoliters (nl) of injection buffer is injected per quail embryo.

Embryos for this study are collected from actively laying quail. Embryo donor quail are inseminated weekly using pooled semen from male quail of the same breed to produce eggs for injection.

On the day of egg collection, fertile quail are euthanized 2h post oviposition by cervical dislocation. The recently ovulated and fertilized eggs are collected from the upper magnum region of the oviduct under sterile conditions and placed in a glass well and covered with Ringers' Medium and maintained at about 40°C until microinjection.

Cytoplasmic injection of artificial chromosomes is achieved using the microinjection apparatus disclosed in US Patent Application No. 09/919,143, filed

July 31, 2001. Chromosomes are injected into the Stage I embryos at a single site in each embryo.

Following microinjection, the embryos are transferred to the oviduct of recipient quail essentially as described in Olsen et al, B. J Exp Zool 109: 355-66, 1948.

5 Typically, about one day after OT, the recipient quail lay a hard shell egg containing the manipulated ovum. Eggs are incubated in an incubator until hatching of the birds.

Eggs from G2 transchromosomal quail test positive for the presence of intact functional interferon alpha 2b.

10 **Example 26: Production of Monoclonal Antibody Using Avian Artificial Chromosomes in Chicken**

An attB site and a hygromycin^R coding sequence are inserted into the OMC24-IRES-LC and OMC24-IRES-HC vectors disclosed in US Patent Application No. 10/856,218, filed July 31, 2001, which are then each cloned into CACs of
15 Example 25 essentially as described in Examples 22 and 23. The CACs are isolated by a dual laser high-speed flow cytometer.

The flow-sorted chromosomes are pelleted by centrifugation and are diluted to a concentration of 7000-12,000 chromosomes per μ l of injection buffer. Approximately 50 nanoliters (nl) of injection buffer is injected per chicken embryo.

20 Embryos for this study are collected from actively laying G. gallus. Embryo donor chickens are inseminated weekly using pooled semen from male chickens of the same breed to produce eggs for injection.

On the day of egg collection, fertile hens are euthanized 2h post oviposition by cervical dislocation. The recently ovulated and fertilized eggs are collected from the
25 upper magnum region of the oviduct under sterile conditions and placed in a glass well and covered with Ringers' Medium and maintained at about 41°C until microinjection.

Cytoplasmic injection of artificial chromosomes containing the OMC24-IRES-LC is achieved using the microinjection apparatus disclosed US Patent Application No. 09/919,143. Approximately 500 chromosomes are injected into the Stage I
30 embryos at a single site.

Following microinjection, the embryos are transferred to the oviduct of recipient chickens essentially as described in Olsen et al, B. J Exp Zool 109: 355-66, 1948. Typically, about one day after OT, the recipient chickens lay a hard shell egg containing the manipulated ovum. Eggs are incubated in an incubator until hatching of the GObirds.

G2 transchromosomal chickens are obtained which contain the artificial chromosome in their genome. The artificial chromosome containing the OMC24-IRES-HC is introduced into embryos obtained from the G2 chickens in essentially the same manner as described for the OMC24-IRES-LC.

Eggs from G1 transchromosomal chickens which contain both the OMC-IRES-LC and OMC24-IRES-HC in their genome are tested for the presence of intact functional monoclonal antibody. A Costar flat 96-well plate is coated with 100 μ l of C Goat-anti-Human kappa at a concentration of 5 μ g/ml in PBS. The plate is incubated at 37 °C for two hours. 200 μ l of 5% PBA is added to the wells followed by an incubation at 37 °C for about 60-90 minutes followed by a wash. 100 μ l of egg white samples (diluted in 1% PBA:LBP) is added to each well and the plate is incubated at 37 °C for about 60-90 min followed by a wash. 100 μ l of a 1:2000 dilution of F2 Goat anti-Human IgG Fc-AP in 1% PBA is added to the wells and the plate is incubated at 37 °C for 60-90 min followed by a wash. The antibody is detected by placing 75 μ l of 1mg/ml PNPP (p-nitrophenyl phosphate) in 5x developing buffer in each well and incubating for about 10-30 mins at room temperature. The detection reaction is stopped using 75 μ l of 1N NaOH. The egg white tests positive for significant levels of the antibody.

Example 27: Cell culture and transfection for the production of an insert containing artificial chromosome and screening for positive clones

pK161 is a cosmid containing a 8.2 kb mouse rDNA insert. The plasmid is produced as disclosed in Csonka et al 2000, Journal of Cell Science 113, 3207-3216. 100 μ g of cosmid pK161 is digested with Cla I, purified by phenol/chloroform extraction and ethanol precipitation then resuspended at approximately 1 μ g/ μ l in TE,

pH 8.0. YAC DNA containing the human light-chain and heavy-chain immunoglobulin loci shown in Fig 27A and 27B are prepared as disclosed in Example 30.

5 LMTK- cells (obtained from ATCC) are cultured at 37°C in 5% CO₂ in a humidified incubator in DMEM (Invitrogen), 10% FBS (Hyclon) (LMTK- media). Prior to the day of transfection, ten 10 cm plates are seeded with approximately 2×10^6 cells per dish. Transfection with ExGen 500 (i.e., sterile 0.01 mM 22 kDa polyethylenimine (PEI), Fermentas Life Sciences) can be performed according to the manufacturers instructions or as follows.

10 On the day of the transfection, LMTK- cells are washed once with 3 ml of Optimem and the media is replaced with 6 ml of Optimem. In an eppendorf tube, 250 µl of HBS (150 mM NaCl, 20 mM HEPES, pH 7.4) is mixed with 3.6 µl of ExGen 500. In a second tube, 250 µl of HBS is mixed with 6 µg of linearized pFK161, 3.0 µg of gel-purified kappa light chain YAC and 3.0 µg of gel-purified heavy chain YAC.
15 The PEI (ExGen) mixture is added dropwise to the DNA mixture, without mixing of the two solutions.

After incubation at RT for 10 min, the solution is gently mixed by pipeting up and down with a wide-bore pipet 3 times. 50 µl of the transfection mix is added to each 10 cm dish of LMTK- cells and the plates are swirled to distribute the DNA/PEI
20 complexes. 4-6 hours post-transfection, the media is replaced with 10 ml of LMTK-media. 48 hours post-transfection, the media is replaced with LMTK- media plus 200 µg/ml G418 (Geneticin, Invitrogen). The selective media is replaced every 2-3 days until colonies are apparent.

Fifty G418-resistant colonies are isolated with cloning cylinders and are
25 transferred to single wells in 24-well tissue culture plates. When the clones are at or near confluency, they trypsinized and split into three 24-well plates.

To determine which clones carry a desired artificial chromosome, metaphase or interphase FISH is performed. Purified light chain YAC DNA is labeled with biotin-14dCTP by random priming (Bioprime DNA labeling system, Invitrogen). The heavy
30 chain YAC DNA is labeled with digoxigenin-1 IdUTP by random priming (Dig High

Prime, Roche Diagnostics). The heavy and light chain YAC probes are mixed and hybridized to metaphase chromosomes or interphase nuclei. The hybridized biotin signals are made visible with fluorescein labeled avidin, and the digoxigenin signals are visualized with rhodamine labeled anti-digoxigenin antibody following standard protocols. The nuclei or chromosomes are counterstained with DAPI and visualized on an Olympus 1X70 microscope configured with DAPI, FITC and rhodamine fluorescent excitation filters.

Two clones are found to have an episomal element indicative of an artificial chromosome. Both clones are positive for the heavy and light chain YACs, indicating that both YACs are incorporated into the artificial chromosomes. The artificial chromosomes are believed to be satellite artificial chromosomes.

Example 28: Copy number determination of Ig loci inserts and determination of structural integrity of the loci in the artificial chromosomes

In order to simplify the interpretation of the analysis of structural integrity of the Ig containing YACs, it is desirable to obtain artificial chromosomes which carry one copy of each YAC. Real time PCR using Taqman® chemistry is utilized to identify clones containing a single copy of the YACs. Several primer/probe sets are designed to detect each YAC. The aniplicon detection probes are labeled using FAM as the dye and TAMRA as the quencher. 10 ng of genomic DNA purified from the positive clones that are identified in Example 30 are assayed in a 30 µl reaction using the TaqMan® Fast Universal PCR Master Mix, No AmpErase® UNG and 7900HT (Applied Biosystems). Amplification curves are compared to standards that are composed of differing amounts of purified YACs in the presence of 10 ng of LMTK-DNA. Both positive clones of Example 27 appear to have a single copy of the light chain YAC as is indicated by overlap of the amplification curves and the Ct value relative to the standard curve. One clone appears to have two or more copies of the heavy chain YAC as the amplification curve had a Ct that is 4 cycles less than the other clone. The other clone appears to have a single copy of the heavy chain YAC. The clone containing one copy of each YAC (clone SC) is selected for further analysis.

PCR primers are designed to amplify 300-500 bp regions of each YAC which are complementary to restriction fragments to be detected in the Southern blot analysis. PCR products are gel purified and quantitated by the Picogreen Assay (Molecular Probes). Radiolabeled probes are generated by random priming using
5 deoxycytidine 5'-[α -³²P] triphosphate and the Rediprime II Random Priming kit (Amersham).

Cells of clone SC are embedded in agarose plugs and subjected to DNA release and restriction digestion according to standard protocols. Several enzymes that cut the YACs into 20 to 150 kb segments are used including Asc I, Pac I and Sbf I. The
10 digested plugs are loaded in multiple lanes such that replicate membranes can be cut from a single membrane. The digested DNAs are separated by PFGE (CHEF) on a 0.8% agarose gel in TAE buffer (switch time 1 = 1 s, switch time 2 = 25 s, 4 V/cm, 15 to 20 h, 14⁰C). The gel is transferred to a UV crosslinker (Stratagene) and exposed to 120 mJ UV radiation. The gel is denatured in 1.5 M NaCl, 0.5 M NaOH for 30
15 minutes at RT and neutralized in 1.5 M NaCl, 1.0 M Tris base, pH 7.4 for 40 minutes at RT. The gel is transferred by capillary action to Genescreen Plus® nylon membrane in 10X SSPE for one to three days. The membrane is briefly rinsed in 2X SSPE and cross-linked with 120 mJ UV radiation (Stratagene). The membrane is cut into replicate pieces and is transferred to roller bottles (Bellco). The membranes are
20 prehybridized in hybridization buffer (1.25X SSPE₃ 0.625% SDS, 40% formamide, IX Denhardt's, 10% dextran sulfate, 0.05 mg/ml denatured salmon sperm DNA) for 2-6 hours at 42°C. The hybridization buffer is changed with new buffer and the appropriate probe is added. The membranes are hybridized overnight at 42°C. The next day the membranes are washed with 0.2X SSPE, 1% SDS or 0.02X SSPE, 1%
25 SDS at 42°C to 65°C until the CPM of each membrane is 400 or less. Membranes are wrapped in Saran Wrap® and exposed one to three days to BioMax MS™ film with a BioMax TranScreen HE™ intensifying screen at -80°C. Clone SC is found to have restriction fragments which demonstrate the structural integrity of both YACs; i.e., no rearrangement of the YACs is apparent.

Example 29: Purification of Ig loci containing artificial chromosome and analysis of human immunoglobulin produced in transgenic avians

Artificial chromosomes are purified from clone SC by flow cytometry and are used for cytoplasmic injections of stage I White leghorn embryos essentially as disclosed in Example 21. 500 embryos are injected with between 100 and 1000 artificial chromosomes. 135 chicks hatch and are analyzed for the presence of the transgene in their blood DNA. DNA is extracted as disclosed in US Patent No. 6,423,488, issued July 23, 2002. 100 ng of DNA is analyzed by real-time PCR using probes to detect the heavy and light chain YACs as disclosed in Example 31. Five birds are found to be positive for the clone SC artificial chromosome at significant levels (> 1 copy of the artificial chromosome for every 100 genomic equivalents).

Serum from hatched birds and eggs from mature hens are analyzed for human Ig λ and IgFc levels by ELISA. Several birds are positive for both human Ig λ and IgFc in their serum, indicating that human IgG is produced in the serum. Eggs from GO hens are collected and the yolks removed. Yolk is diluted and analyzed for human Ig λ and IgFc levels by ELISA. Several hens contain human IgG in the yolk of their eggs.

G1 birds are produced from the GO birds as disclosed herein. Each of the positive G1 birds include the artificial chromosome in substantially all of their somatic cells as demonstrated by FISH. The germline transgenic G1 birds produce substantial quantities of polyclonal antibodies which are deposited in the egg. For example, human polyclonal antibody is present in an amount greater than about 10 μ g/egg or greater than about 0.1 mg/egg.

Example 30: Isolation and characterization of human immunoglobulin loci YACs

Two YACs that contain substantial portions of the human light-chain and heavy-chain immunoglobulin loci are shown in Fig. 27A and 27B. These constructs contain multiple variable, D, J and constant regions, as well as elements required for gene expression, gene rearrangement and constant chain switch. The lambda light-chain construct, IgLambda, is a 410 kb YAC that has been previously used to express human polyclonal antibodies in transgenic mice. See, for example, US patent

application No. 2004/0231012, published November 18, 2004 and Popov et al (1999) J. Exp. Med. 189:1611-1619, the disclosures of which are incorporated in their entireties herein by reference. The heavy-chain construct, IgHeavy-2, is a 300 kb derivative of the YAC shown in Fig. 27A that has been used to express human polyclonal in mice (Nicholson et al (1999) J Immunol 163:6898-6906) to which a functional human gamma-constant gene segment has been added 3' of the C δ region.

YAC containing strains of *Saccharomyces cerevisiae* were grown in a yeast nitrogen base medium with 2% glucose and an appropriate selective amino acid at 30°C for 4 days. Total DNA agarose plugs were prepared from the yeast strains using the protocol of Iadonato, S. P., and A. Gnirke (1996) modified as follows:

Yeast cells were centrifuged, washed with 50 mM EDTA pH 8 and resuspended at 2×10^9 cells/ml in 50 mM EDTA pH 8. The cell suspension was heated to 45-50°C and added to an equal volume of 2% LMP agarose that had been melted and brought to 45-50°C. Cells and agarose were mixed and dispensed into plug molds which were then placed at 4°C. Hardened plugs were placed in spheroplasting solution (1 M sorbitol, 20 mM EDTA, 10 mM Tris-HCl pH 7.5, 14 mM mercaptoethanol, 3% lyticase solution (#170-3593 Bio-Rad)) at 37°C for 4 hours with gentle agitation. Plugs were then washed in LDS solution (1% lithium dodecyl sulfate, 100 mM EDTA pH 8, 10 mM Tris-HCl pH 8) for 15 minutes and were then placed in LDS solution for 16 hours at 37°C with gentle agitation. Plugs were then washed 3 times for 30 minutes in NDS solution (500 mM EDTA, 10 mM Tris base, 1% sarkosyl pH 9) and 5 times for 30 minutes with TE (10 mM Tris-HCl pH 8, 1 mM EDTA pH 8) with gentle agitation.

The intact YACs were separated by contour-clamped homogeneous electric field (CHEF) electrophoresis in 1% low-melting point agarose gels using 0.5X TBE buffer at 14°C and a 30 second constant switch time at 5 V/cm for 36 hr. Gel slices containing the YAC of interest were equilibrated 2 hr with microinjection buffer containing 10 mM Tris-Cl pH 7.5, 0.1mM EDTA pH 8.0, 100 mM NaCl, 30 mM spermine, and 70 mM spermidine. The gel slices were melted at 68°C for 20 min and then digested with GELase (5 U/100 mg) at 42°C for 2 hr. Integrity of Each YAC

sample was then confirmed by CHEF electrophoresis on a 1.5% agarose gel with 0.5X TBE buffer at 14°C using a 30 second constant switch time and 5 V/cm for 24 hr.

Example 31; Transgenesis and immunoglobulin expression

5 Purified heavy-chain and light-chain YAC DNAs prepared as disclosed in Example 31 were co-injected into early embryos to generate transgenic animals as essentially disclosed in Example 21. A volume of 50 nl of 110 pg chromosome DNA per µl of microinjection buffer was injected into each of several hundred embryos. Testing for the production of human light-chain in serum of resultant chickens was
10 performed using a human lambda ELISA quantitation kit (#E80-116) from Bethyl Laboratories (Montgomery, TX). In the procedure, both the capture antibody and detection antibodies were diluted 1:2000. Quantitation of antibody containing associated light-chain and heavy-chains was performed by replacing the detection antibody in the above kit with an alkaline phosphatase-conjugated goat anti-human
15 IgG, Fc gamma-antibody (diluted 1:2000) (#109-056-098, Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) and followed by detection using a TMB substrate. At least one bird was shown to express human immunoglobulins by ELISA (Table 5) in the serum.

20 Table 5

Bird	Lambda Light-Chain	Whole IgGλ
#6946	26 ng/ml	24 ng/ml
Control	0	0

Example 32

25 Identification of Target Sequences and Probe Preparation

A 234bp sequence of mouse major satellite DNA sequence (SEQ ID NO: 14) present in SATACs (see, for example, US patent publication No. 2003/0119104, filed May 30, 2002, the disclosure of which is incorporated in its entirety herein by

reference) was scanned for 6mer repeat regions. Two such sequences were identified, each present five times in the 234bp sequence. These consensus sequences can be targets for labeled polyamide probes. The identified sequences are shown in Table 6.

5 SEQ ID NO: 14, Mouse Major Satellite DNA sequence

gacctggaatatcgcgagtaaactgaaaatcacggaaaatgagaaatacacactttaggacgtgaaatatggcgaggaaaa
ctgaaaaagggtggaaaatttagaatgtccactgtaggacgtggaatatggcaagaaaactgaaaatcatggaaaatgagaa
acatccacttgacgacttgaaaaatgacgaaatcactaaaaaacgtgaaaaatgagaaatgcacactgaaggac

10

Table 6

Pattern	Searched	Consensus	Copy No
TGAAAA		TGAAAA	5
GAAAAT		GAAAAT	5

15

The centromeric region of SATACs contain copies of the mouse minor satellite sequence which can also be targeted. For example, polyamide probes can be designed to target pentameric repeat sequences and octameric repeat sequences located within the 120 bp sequence of the mouse minor satellite (SEQ ID NO: 15). The consensus or repeat sequences identified are shown in Table 7.

SEQ ID NO: 15, Mouse minor satellite DNA Sequence

20

gagtgagttacactgaaaaacacatacgttgaaaccggcattgtagaacagtgtatatcaatgagttacaatgagaaaaatg
gaaaatgataaaaaccacagtgtagaacatattagatgtgtgagttacactgaaaaacacattccttggaacgggattttag
aactgtgtatatcaatgagttacaatgagaacatggaaaatgataaaaaccacactgtagaacatttagatgagtgagttaca
ctgaaaaacacatatgttgaaa

25

Table 7

Pattern Searched	Consensus	Copy No
GAAAA	GAAAA	6
AAAAA	AAAAA	6
AATGA	AATGA	6
TGAGTTAC	TGAGTTAC	5
GAGTTACA	GAGTTACA	5

The heterochromatic region of SATACs contain copies of the mouse rDNA (ribosomal RNA encoding DNA) sequence can also be targeted. For example, polyamide probes can be designed to target 5-mer repeat sequences, six-mer repeat sequences and seven-mer repeat sequences located within the 120 bp sequence of mouse rDNA shown in SEQ ID NO. 16. The consensus or repeat sequences are shown in Table 8.

10

Table 8

Pattern Searched	Consensus	Copy	No
TGTGC	TGTGC	6	
TTCCC	TTCCCJL	6	
CGTGC	CGTGC L	8	
CCGCC	CCGCC L	21	
CGCCG	CGCCG LL	25	
CCCGCG	CCCGCG LL	15	
CCCGTC	CCCGTC LL	5	
CCGGCG	CCGGCG I	7	
CCCGGG	CCCGGGJL	5	
TCTCTCG	TCTCTCGJI	6	

Polyamide probes can be constructed to recognize the repeat sequences shown in Tables 6, 7 and 8 and/or other sequences contained in artificial chromosomes that will provide for a facilitated isolation of the artificial chromosomes, for example, by flow cytometry. Methods of making polyamide probes are well known in the art and are disclosed, for example, in the certain references cited herein and in the specification.

Example 33

Preparation and flow sorting of chromosomes

Chromosome suspensions are prepared from CHO cells (e.g., chromosome suspensions containing artificial chromosomes such as those disclosed in US patent publication No. 2003/0119104, filed May 30, 2002) using a modification of the polyamine-based method described by Sillar and Young (1981) A new method for the preparation of metaphase chromosomes for flow analysis, J. Histochem. Cytochem., 29, 74-78 and Lalande et al (1984) Development and use of metaphase chromosome flow-sorting methodology to obtain recombinant phage libraries enriched for parts of the human X chromosome, Cytometry, 5, 101-107, the disclosures of which are incorporated herein in their entirety by reference. Briefly, cells cultured in RPMI medium containing 20% fetal bovine serum are arrested at mitosis by incubation in 0.1 $\mu\text{g/ml}$ colcemid for about 16 h. The cells are collected by centrifugation, resuspended in 40 mM KCl for 10 min and then centrifuged again. The pellet is resuspended in cold buffer containing 80 mM KCl, 20 mM NaCl, 15 mM Tris-HCl pH 7.2, 2 mM EDTA, 0.5 mM EGTA, 7 mM β -mercaptoethanol, 0.2 mM spermine, 0.5 mM spermidine and 0.12% digitonin, and incubated on ice for 10 min. The suspension is vortexed vigorously for 2 min and then stored for up to 90 days at 4°C before use.

Prior to flow analysis, the chromosomes are stained for about 2 h with 1 μM of fluorescently labeled polyamide probe and 2 $\mu\text{g/ml}$ HO (Hoechst 33258). The polyamide probe targets one or more of the nucleotide sequences specified in Table 6, Table 7 and/or Table 8 and is produced essentially as disclosed in Dervan (2001) Molecular recognition of DNA by small molecules. Bioorg Med Chem 9: 2215-35.

Sodium citrate and sodium sulfite are added to chromosomes 15 to 30 min before flow analysis at final concentrations of 10 and 25 mM, respectively, to improve chromosome resolution.

Chromosomes are separated using an Influx flow sorter (Cytospeia, Inc., Seattle, WA). One laser is tuned to emit ultraviolet light (351-364 nm, 250 mW) to excite HO, and HO fluorescence is measured after passing through a 425-nm long-pass filter and a 458-nm rejection-band filter. A second laser is tuned to 458 nm (250 mW) to excite the fluorescently labeled polyamide probe. The fluorescently labeled polyamide probe fluorescence is measured after passing through a 500-nm long-pass filter and a 458-nm rejection-band filter. Alternatively, fluorescein fluorescence is measured following excitation at 488 nm (250 mW) after passing through a 530/40 band-pass filter. The fluorescence pulses from the individual chromosomes are integrated by a data acquisition system, and are collected in listmode at a rate of about 1000 chromosomes per second.

15

Example 34

Purification of SATACs Contained in Micronuclei

Micronucleation of chromosomes in ChY1 cells containing the artificial chromosome (e.g., SATACs) is induced by incubation for 72 h in the presence of 1 µg/ml of colchicine in growth medium.

Micronuclei are isolated essentially as described in Labidi (Labidi, B, et al. Procedure for isolating micronuclei from rat kangaroo cultured cells containing individualized chromosomes. Eur J Cell Biol 38: 165-70, 1985), the disclosure of which is incorporated in its entirety herein by reference. Briefly, micronucleated cells are harvested by trypsin-EDTA treatment, rinsed twice in PBS, and resuspended in 2 vol TKM buffer (10 mM Tris-HCl, pH 7.4, 10 mM KCl, and 3 mM MgCl₂) containing 0.05% collagenase 1 A (Sigma). Cell lysis is accelerated by gentle shearing of the suspension through a 26-gauge needle and 1 mM phenyl methyl sulfonyl fluoride (PMSF) is added. Isolated micronuclei are collected by low-speed centrifugation (1500g) and washed twice with 4 vol of Tris-polyamine buffer (TPB; 15

niM Tris-HCl, pH 7.4, 0.2 mM spermine, 0.5 niM spermidine, 2 niM EDTA, 0.5 mM EGTA, 80 mM KCl, 20 mM NaCl, and 14 mM B-mercaptoethanol).

The micronuclei containing chromosomes are stained and purified by flow cytometry in essentially the same manner as described for the staining and flow
5 cytometry purification of artificial chromosomes, as disclosed in Example 33.

During the flow cytometry, micronuclei collection is limited to the window in the fluorescence histogram where micronuclei containing a single SATAC are located, which can be defined by conventional methodologies.

10 **Example 35**

Production of Transchromosomic Chickens Using Satellite DNA-Based Artificial Chromosomes

The flow-sorted artificial chromosomes of Example 33 or micronuclei containing the artificial chromosomes of Example 34 are pelleted by centrifugation of
15 a 750 μ l sample containing approximately 10^8 chromosomes (artificial chromosomes or micronuclei containing artificial chromosomes) at 2500 x g for 30 min at 4°C. The supernatant, except the bottom 30 microliters (μ l) containing the chromosomes, is removed resulting in a concentration of about 7000 to 11,500 chromosomes per μ l of injection buffer (Monteith, et al. Methods Mol Biol 240: 227-242, 2004).
20 Approximately 25 to 100 nanoliters (nl) of injection buffer is injected per embryo.

Early stage embryos (e.g., stage I embryos) are collected from 24 to 36 week-old hens from commercial White Leghorn variety of *G. gallus*. Embryo donor hens are inseminated weekly using pooled semen from roosters of the same breed to produce eggs for injection.

25 On the day of egg collection, fertile hens are euthanized 2h post oviposition by cervical dislocation. Typically, oviposition is followed by ovulation of the next egg after about 24 minutes (Morris, Poultry Science 52: 423-445, 1973). The recently ovulated and fertilized eggs are collected from the upper magnum region of the oviduct under sterile conditions and placed in a glass well and covered with Ringers'

Medium (Tanaka, et al. J Reprod Fertil 100: 447-449, 1994) and maintained at 41⁰C until microinjection.

Cytoplasmic injection of artificial chromosomes is achieved using the microinjection apparatus disclosed in US Patent Application No. 11/159,973, filed
5 June 23, 2005, the disclosure of which is incorporated in its entirety herein by reference. Chromosomes are injected into the Stage I embryos at a single site. Each embryo is cytoplasmically injected with approximately 400 to 1000 chromosomes. The chromosomes are injected in a suspension of 25 to 100 nanoliters (nl) of injection buffer.

10 Following microinjection, the embryos are transferred to the oviduct of recipient hens using the ovum transfer (OT) procedure of Olsen (Olsen, M and Neher, B. J Exp Zool 109: 355-66, 1948), with the exception that the hens are anesthetized by isofluorane gas. Typically, about 26h after OT₅ the recipient hens lay a hard shell egg containing the manipulated ovum. Eggs are incubated for 21 days in a regular
15 incubator until hatching of the birds.

Transchromosomic founders are identified based on PCR analysis and FISH analysis. Analysis of G1 embryos from a test bird show the artificial chromosome to be transmitted through the germline.

While this invention has been described with respect to various specific
20 examples and embodiments, it is to be understood that the invention is not limited thereto and that it can be variously practiced with the scope of the following claims.

What is claimed is:

1. A method comprising:
isolating an artificial chromosome;
5 introducing the artificial chromosome into an avian embryo;
maintaining the embryo under conditions suitable for the embryo to
develop and hatch as a chick; and
maintaining the chick under conditions suitable to obtain a mature
avian wherein the artificial chromosome is present in the genome of the mature avian.
10
2. The method of claim 1 wherein isolating the artificial chromosome is
facilitated by flow cytometry.
3. The method of claim 2 wherein the flow cytometry is facilitated by a
15 probe which is associated with the artificial chromosome.
4. The method of claim 3 wherein the probe is a polyamide probe.
5. The method of claim 2 wherein the artificial chromosome is present in
20 a micronuclei.
6. The method of claim 1 wherein the artificial chromosome is introduced
into the avian embryo by injection.
7. The method of claim 1 comprising transferring the embryo to a recipient
25 female avian.
8. The method of claim 1 wherein the embryo is an early stage embryo.
9. The method of claim 1 wherein the embryo is a stage I embryo.
30

10. The method of claim 1 wherein the avian is a chicken.
11. The method of claim 1 wherein the artificial chromosome comprises a
5 heterologous recombination site.
12. The method of claim 11 wherein the artificial chromosome comprises
more than one heterologous recombination site.
- 10 13. The method of claim 1 wherein the artificial chromosome comprises a
heterologous coding sequence.
14. The method of claim 13 wherein the heterologous coding sequence
comprises a pharmaceutical protein coding sequence.
- 15 15. The method of claim 13 wherein the heterologous coding sequence
encodes an immunoglobulin polypeptide.
16. The method of claim 13 wherein the heterologous coding sequence
20 encodes a cytokine.
17. The method of claim 1 wherein the artificial chromosome comprises a
promoter.
- 25 18. The method of claim 1 wherein the artificial chromosome comprises a
promoter which functions in tubular gland cells.
19. The method of claim 1 wherein the artificial chromosome comprises an
IRES.
- 30

20. The method of claim 1 comprising obtaining an offspring from the mature avian wherein the offspring contains an artificial chromosome in its genome.

21. The method of claim 1 wherein the artificial chromosome is a stabilized
5 artificial chromosome isolated from an avian cell.

22. An avian produced by the method of claim 1.

23. A method comprising:
10 isolating an artificial chromosome;
introducing the artificial chromosome into an avian embryo by
injection;
maintaining the embryo under conditions suitable for the embryo to
develop and hatch as a chick; and
15 maintaining the chick under conditions suitable to obtain a mature
avian wherein the artificial chromosome is present in the genome of the mature avian.

24. The method of claim 23 comprising transferring the embryo to a recipient
female avian.
20

25






Fig. 1

*Fig. 2*

**Fig. 3**




Fig. 4







Fig. 5

**Fig. 6**

***Fig. 7***

**Fig. 8**

pCMV-C31int (SEQ ID NO: 1)

CATTGCGCCATTTCAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTATT
ACGCCAGCCAATACGCAAACCGCTCTCCCCGCGCGTTGGCCGATTCATTAATGCAGGATCG
ATCCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGAAAA
AAATGCTTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAA
TAAACAAGTTAACAACAACAATTGCATTCATTTTATGTTTCAGGTTTCAGGGGGAGGTGTGGG
AGGTTTTTTAAAGCAAGTAAACCTCTACAAATGTGGTATGGCTGATTATGATCATGAACAG
ACTGTGAGGACTGAGGGGCCTGAAATGAGCCTTGGGACTGTGAATCTAAAATACACAAACAA
TTAGAATCACTAGCTCCTGTGTATAATATTTTCATAAATCATACTCAGTAAGCAAAACTCTC
AAGCAGCAAGCATATGCAGCTAGTTTAAACACATTATACACTTAAAAATTTTATATTTACCTT
AGAGCTTTAAATCTCTGTAGGTAGTTTGTCCAATTATGTCACACCACAGAAGTAAGGTTTCT
TCACAAAGATCCCAAGCTAGCTTATAATACGACTCACTATAGGGAGAGAGCTATGACGTCGC
ATGCACGCGTAAGCTTGGGCCCCCTCGAGGGATCCGGGTGTCTCGCTACGCCGCTACGTCTTC
CGTGCCGCTCCTGGGCGTCGTCTTCGTGCTGCTCGGTGCGGCGGCTTCGCCACGTGATCGAAG
CGCGCTTCTCGATGGGCGTTCCCTGCCCCCTGCCCGTAGTCGACTTCGTGACAACGATCTTG
TCTACGAAGAGCCGACGAACACGCGCTTGTCTACTGACGCGCGCCCCACACGACTT
AGGGCCGGTCGGGTGAGCGTCGGCGTCTTCGGGGAACCATTTGGTCAAGGGGAAGCTTCGGGG
CTTCGGGCGGCTTCAAGTTCGGCAAGCCGCTCTTCGCCCTTGTGCGGGAGCGTCAGCGCT
GCCTGTTGCTTCCGGAAGTGTCTTCCCTGCCAACGGGTCCGTGCTACGCGCCTGCCGCGCGGTC
TTCGTACAGCTCTTCAAGGGCGTTCAGGGCGTCGGCGCGCTCCGCAACAAGGTTCCGCCGT
CGCCGCTCTTCTCAGGCGCCTCAGTGAGCTTGCCGAAGCGTCGGGCGGCTTCCACAGAAGC
GCCAACGCTCTTCTGTCGCCCTTCGGCGTGCTTGCCTGATCTTGTGAAGATGCGTTCCGCGCGG
CTTGTGCGAGTCCCGCATGCTGACGTTGCACGTGCGCTTCTGCTGCTGCCAGGTGCCGACGGGT
CGACCACCTTCCGGCGACGGCAGCGGTAAGAGTCTTGATCGATTCTTCCCCGCGCTTCGAA
GTCATGACGGCGCCACACTCGCAGTACAGCTTGTCCATGGCGGACAGAATGGCTTGCCCCCG
GGAAAGCCCCCTTGCCGCGCCCCCTGCCGTCCAACCACGCCTGAAGCTCATACCACTCAGCGG
GCTCGATGATCGGTCCGCAATCAAGCTCGACCGGCGGAGCGTGATCGGGTCGCGCTGAATG
CGGTAACCCCTCAATCTTCGTGGTCGGCGTGCCGTCCGGCTTCTTCTGTAGATCACCTCAGC
GGCGAAGCCCGCAATACGCGGGTCCCGAAGGATTCCGCATAACGGTTGCCGGGTCCGACGCG
TTGAAGCGGTCTTCTTCCCAATCGTCTCGCCCCGGGTGCGCACGGCGTCAGCGTCCATGCGC
TTACAAAGCCCCGTGATGCTGCCCGGGTGAATGGCGGCTTGACTGCCCGGCTTGAAGGGAAG
GTGTTTGTGCGTCTTGATCTCACGCCACCACCACGGATTACGTGCGGCTCGAAGTCAAGG
GTCCGGTAAGGGGAGTGGTCGAGTGCAGCAAGCTTGTGATGACGACATTGACCATTCGGCCG
TTGCGCGTGATCTCCTTCGTCTCCGAAACAAGCTCGAAGCCGTAAGGCGCCTTCCCGCCGAC
GTACCCGCCCAATTTCGCGCTGAAGGTTCTTCGTGTCGAGAATCTTCGCCGACTTCAGCGAAG
ATTCTTTGTGCGACGCGTCGAGCCGCAATAATCAGGTGAATCAGGTCCATGACGTTTCCCTGC
CGGAAGACGCCTTCCCTGAGTGGAACAATCGTCACGCCCAGGGCGAGCAATTCGAGACAAT
CGGAATCGCGTCCATGACCTTCAGGCGCGAGAAGCGCGACACGTCATAGACAATGATCATGT
TGAGCCGCCCGGCGCGGCATTTCGTTTCAAGGATGCGTTTCAAGTCCGGGCGCTCCGCCGTCCCG
AACGCCGACGTGCCCGGCGCTTCGCTGAAATGCCCGACGAACCTGAACCGGCCCCCGTCGCG
CTCGACTTCGCGCTGAAGGTGCGCCGCTTGTCTTCGTTGGCGCTACGCTGTGTGCTGGGC
TTGCTGCGCTCGAATTCTCGCGCTCGCGCGACTGACGGTCGTAAGCACCCGCGTACGTGTCC
ACCCCGTACAAACCCCTTGTGTCATGTCGGCGACCCCTACGACTAGTGAGCTCGTCGACCCG
GGAATTCCCGACCGGTACCTGCAGGCGTACCTTCTATAGTGTACCTAAATAGCTTTTTTGCA
AAAGCCTAGGCTAGAGTCCGGAGGCTGGATCGGTCCCGGTGTCTTCTATGGAGGTCAAACA
GCGTGGATGGCGTCTCCAGGCGATCTGACGGTTCACTAAACGAGCTCTGCTTATATAGACCT
CCCACCGTACACGCTTACCGCCCATTTGCGTCAATGGGGCGGAGTTGTTACGACATTTTGA
AAGTCCCGTTGATTTTGGTGCCAAAACAACTCCATTGACGTCAATGGGGTGGAGACTTGG
AAATCCCCGTGAGTCAAACCGCTATCCACGCCCATTTGATGTACTGCCAAAACCGCATACCA
TGTTAATAGCATGACTAATACGTAGATGATGACTGCCAAGTAGGAAAGTCCCATAGGTATG
TACTGGGCATAATGCCAGGCGGGCCATTTACCGTCATTGACGTCAATAGGGGGCGTACTTGG
CATATGATACACTTGATGTACTGCCAAGTGGGCAGTTTACCGTAAATACTCCACCCATTGAC
GTCAATGGAAAGTCCCTATTGGCGTTACTATGGGAACATACGTCAATTATGACGTCAATGGG
CGGGGTCGTTGGGCGGTGAGCCAGGCGGGCCATTTACCGTAAGTTATGTAACGACCTGCAC

GATGCTGTTTCCTGTGTGAAATTGTTATCCGCTCACAATTCACACATTATACGAGCCGGAA
GCTATAAAGTGTAAAGCCTGGGGTGCCTAATGAGTGAAAGGGCCTCGTATACGCCTATTTTT
ATAGGTTAATGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATG
TGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGA
CAATAACCCTGATAAATGCTTCAATAATATTGAAAAACGCGCGAATTGCAAGCTCTGCATTA
ATGAATCGGCCAACGCGCGGGGAGAGCGGTTTGGCGTATTGGGCGCTCTTCCGCTTCCTCGC
TCACTGACTCGCTGCGCTCGGTCTGCTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCG
GTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCA
GCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTGGCTGGCGTTTTTCCATAGGCTCCGCCCC
CTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAA
AGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCT
TACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGCGCTTTCTCAATGCTCAGCT
GTAGGTATCTCAGTTCGGTGTAGGTCTGCTCCGCTCAAGCTGGGCTGTGTGCACGAACCCCC
GTTTACGCCCCGACCGCTGCGCCTTATCCGGTAACATCTGCTCTTGAATCCACCCGGTTAACA
CGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCG
GTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGT
ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAA
ACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAA
AAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAAAA
TCACGTTAAGGGATTTTGGTCATGCCATAACTTCGTATAGCATACATTATACGAAGTTATGG
CATGAGATTATCAAAAAGGATCTTACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAAT
CAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCA
CCTATCTCAGCGATCTGTCTATTTCTGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGAT
AACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCAC
GCTCACC GGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGT
GGTCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAG
TAGTTGCGCAGTTAATAGTTTGC GCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTAC
GCTCGTCTGTTTGGTATGGCTTCATTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGA
TCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTCAGAAAGTAA
GTTGGCCGCAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGC
CATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGT
ATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAG
AACTTTAAAAGTGCTCATCATTTGGAACAGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTAC
CGCTGTTGAGATCCAGTTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTT
ACTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGC AAAATGCCGCAAAAAAGGGAAT
AAGGGCGACACGGAAATGTTGAATACTCATACTCTTCTTTTCAATATTATTGAAGCATTT
ATCAGGGTTATTGTCTCATGCCAGGGGTGGGCACACATATTTGATACCAGCGATCCCTACAC
AGCACATAATTCAATGCGACTTCCCTCTATCGCACATCTTAGACCTTTATTCTCCCTCCAGC
ACACATCGAAGCTGCCGAGCAAGCCGTTCTCACCAGTCCAAGACCTGGCATGAGCGGATACA
TATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCGAAAAAGTG
CCACCTGAAAATTGTAAACGTTAATATTTTGTAAATTCGCGTTAAATTTTTTTTAAATCAG
CTCATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAAGAATAGACCG
AGATAGGGTTGAGTGTGTTCCAGTTTGGAAACAAGAGTCCACTATTAAAGAACGTGGACTCC
AACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGATGGCCCACTACGTGAACCATCACCCCTA
ATCAAGTTTTTTGGGGTCGAGGTGCCGTAAAGCACTAAATCGGAACCCTAAAGGGAGCCCCC
GATTTAGAGCTTGACGGGGAAAGCCGGCGAACGTGGCGAGAAAGGAAGGGAAGAAAGCGAAA
GGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTACGCTGCGCGTAACCACCACACCCGC
CGCGCTTAATGCGCCGCTACAGGGCGCGCT

Fig. 9

pCMV-luc-attB (SEQ ID NO: 2)

CTCTATCGATAGGTACCGAGCTCTTACGCGTGCTAGCCCTCGAGCAGGATCTATACATTGAA
TCAATATTGGCAATTAGCCATATTAGTCATTGGTTATATAGCATAAATCAATATTGGCTATT
GGCCATTGCATACGTTGTATCTATATCATAATATGTACATTTATATTGGCTCATGTCCAATA
TGACCGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATT
AGTTCATAGCCCATATATGGAGTTCCGCGTTACATAAATACGGTAAATGGCCCGCCTGGCT
GACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCA
ATAGGGACTTTCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGT
ACATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCCCG
CCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTTCTACTTGGCAGTACATCTACGTA
TTAGTCATCGCTATTACCATGGTGATGCGGTTTGGCAGTACATCAATGGGCGTGGATAGCG
GTTTGACTCAGGGGATTTCCAAGTCTCCACCCCAATTGACGTCAATGGGAGTTGTTTTGGC
ACCAAAATCAGGGGACTTTCCAATAATGTCGTAACAACTCCGCCCCATTGCGCAAAATGGGC
GGTAGGCGTGACGGTGGGAGGTCATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGC
CTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCC
CCTCGAAGCTCGACTCTAGGGGCTCGAGATCTGCGATCTAAGTAAGCTTGGCATTCCGGTAC
TGTTGGTAAAGCCACCATGGAAGACGCCAAAAACATAAAGAAAGGCCCGGCCATTCTATC
CGCTGGAAGATGGAACCGCTGGAGAGCAACTGCATAAGGCTATGAAGAGATACGCCCTGGTT
CCTGGAACAATTGCTTTACAGATGCATATCGAGGTGGACATCACTTACGCTGAGTACTT
CGAAATGTCCGTTCCGTTGGCAGAAGCTATGAAACGATATGGGCTGAATACAAATCACAGAA
TCGTCGTATGCAGTGAAACTCTCTCAATTCTTTATGCCGGTGTTGGGCGCGTTATTTATC
GGAGTTGCAGTTGCGCCCGCGAACGACATTTATAATGAACGTGAATTGCTCAACAGTATGGG
CATTTGCGAGCCTACCGTGGTGTTTCGTTTCCAAAAAGGGGTTGCAAAAAATTTTGAACGTGC
AAAAAAGCTCCCAATCATCCAAAAAATTATTATCATGGATTCTAAAACGGATTACCAGGGA
TTTCAGTCGATGTACACGTTTCGTCACATCTCATCTACCTCCCGGTTTTAATGAATACGATTT
TGTGCCAGAGTCCTTCGATAGGGACAAGACAATTGCACACTGATCATGAACCTCCTCTGGATCTA
CTGGTCTGCCTAAAGGTGTCGCTCTGCCTCATAGAAGTGCCTGCGTGAGATTCTCGCATGCC
AGAGATCCTATTTTTTGGCAATCAAATCATTCGGGATACTGCGATTTTAAGTGTGTTCCATT
CCATCACGGTTTTTGAATGTTTACTACACTCGGATATTTGATATGTGGATTTTCGAGTCTGCT
TAATGTATAGATTTGAAGAAGAGCTGTTTCTGAGGAGCCTTCAGGATTACAAGATTCAAAGT
GCGCTGCTGGTGCCAACCCTATTCTCCTTCTTCGCCAAAAGCACTCTGATTGACAAATACGA
TTTATCTAATTTACACGAAATTGCTTCTGGTGGCGCTCCCTCTCTAAGGAAGTCGGGGAAG
CGGTTGCCAAGAGGTTCCATCTGCCAGGTATCAGGCAAGGATATGGGCTCACTGAGACTACA
TCAGCTATTCTGATTACACCCGAGGGGGATGATAAACCAGGGCGCGGTGCGGTAAAGTTGTTCC
ATTTTTTGAAGCGAAGGTTGTGGATCTGGATACCGGGAAAACGCTGGGCGTTAATCAAAGAG
GCGAAGTGTGTGTGAGAGGTCTATGATTATGTCCGGTTATGTAAACAATCCGGAAGCGACC
AACGCCTTGATTGACAAGGATGGATGGCTACATTCTGGAGACATAGCTTACTGGGACGAAGA
CGAACACTTCTTCATCGTTGACCGCCTGAAGTCTCTGATTAAGTACAAAGGCTATCAGGTGG
CTCCCGCTGAATTGGAATCCATCTTGCTCCAACACCCCAACATCTTCGACGCAGGTGTGCGA
GGTCTTCCCGACGATGACGCCGGTGAAGTCTCCCGCCCGCGTTGTTGTTTTGGAGCACGGAAA
GACGATGACGGAAAAAGAGATCGTGGATTACGTGCGCAGTCAAGTAACAACCGCGAAAAAGT
TGCGCGGAGGAGTTGTGTTTGTGGACGAAGTACCGAAAGGTCTTACCGGAAAACCTCGACGCA
AGAAAAATCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGAAAGATCGCCGTGTAATTCTA
GAGTCGGGGCGGCGCGCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGACAAA
CCACAAC TAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTA
TTTGTAAACCATATAAGCTGCAATAAACAGTTAACAACAACAAATTGCATTCAATTTATGTT
TCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAAACCTTACAAATGTGGTA
AAATCGATAAGGATCAATTCCGGCTTCAGGTACCGTGCACGATGTAGGTACGGTCTCGAAGC
CGCGGTGCGGGTGCCAGGGCGTGCCCTTGGGCTCCCCGGGCGCGTACTCCACCTCACCCATC
TGGTCCATCATGATGAACGGGTGAGGTGGCGGTAGTTGATCCCGGCGAACGCGCGGCGCAC
CGGGAAGCCCTCGCCCTCGAAACCGCTGGGCGCGGTGGTCACGGTGAGCACGGGACGTGCGA
CGGCGTGGGCGGTGCGGATACGCGGGGCGAGCGTCAGCGGGTTCTCGACGGTCACGGCGGGC
ATGTCGACAGCCGAATTGATCCGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAGCTC
CTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTATCATGC

AACTCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTC
GGTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAG
AATCAGGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGT
AAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAA
TCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTTCCCC
CTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCCTGCCGCTTACCGGATACCTGTCCGCC
TTTCTCCCTTCGGGAAGCGTGGCGTTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCCGT
GTAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTACGCCCCGACCGCTGCG
CCTTATCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCA
GCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAA
GTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGC
CAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCACCGCTGGTAGC
GGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCC
TTTGATCTTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTG
TCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAATGAAGTTTTTAA
TCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGC
ACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGA
TAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCA
CGCTCACCGGCTCCAGATTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAG
TGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGAAGCTAGAGTAA
GTAGTTCGCCAGTTAATAGTTTGCAGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTCA
CGCTCGTCGTTTTGGTATGGCTTCATTACGCTCCGGTTCCCAACGATCAAGGCGAGTTACATG
ATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAGTA
AGTTGGCCGCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATG
CCATCCGTAAAGATGCTTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTG
TATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCA
GAACTTTAAAGTGCTCATCTATTGGAAAAACGTTCTTTCGGGGCGAAAACTCTCAAGGATCTTA
CCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTT
TACTTTTACCAGCGTTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAA
TAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAGCATT
TATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAAT
AGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGCGCCCTGTAGCGGCGCATTA
GCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCC
GCTCCTTTTCGCTTTCTTCCCTTCCCTTCTCGCCACGTTTCGCCGGCTTTCCCGTCAAGCTCT
AAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAAAAAC
TTGATTAGGGTGATGGTTTACGTTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTG
ACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAACCC
TATCTCGGTCTATTCTTTTGAATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTAAAA
ATGAGCTGATTTAACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTACAATTTCC
CATTCGCCATTTCAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTATT
ACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTTGAGCGG
CCGCAATAAAATATCTTTATTTTATTACATCTGTGTGTTGGTTTTTTGTGTGAATCGATAG
TACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAACTAGCAAAATAGGCTGTC
CCAGTGCAAGTGCAGGTGCCAGAACATTT

Fig. 10

pCMV-luc-attP (SEQ ID NO: 3)

CTCTATCGATAGGTACCGAGCTCTTACGCGTGCTAGCCCTCGAGCAGGATCTATACATTGAA
TCAATATTGGCAATTAGCCATATTAGTCATTGGTTATATAGCATAAATCAATATTGGCTATT
GGCCATTGCATACGTTGTATCTATATCATAATATGTACATTTATATTGGCTCATGTCCAATA
TGACCGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATT
AGTTCATAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCCGCTGGCT
GACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCA
ATAGGGACGTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGT
ACATCAAGTGTATCATATGCCAAGTCGCCCCCTATTGACGTCAATGACGGTAAATGGCCCG
CCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTCCTACTTGGCAGTACATCTACGTA
TTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGCGTGGATAGCG
GTTTGACTCACGGGGATTTCGAAGTCTCCACCCCATGACGTCAATGGGAGTTTGTTTTGGC
ACCAAAATCAACGGGACTTTCAAAAATGTCGTAACAACTCCGCCCCATTGACGCAAAATGGGC
GGTAGGCGGTGACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGC
CTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCC
CCTCGAAGCTCGACTCTAGGGGCTCGAGATCTGCGATCTAAGTAAGCTTGGCATTCCGGTAC
TGTTGGTAAAGCCACCATGGAAGACGCCAAAAACATAAAGAAAGGCCCGCGCCATTCTATC
CGCTGGAAGATGGAACCGCTGGAGAGCAACTGCATAAGGCTATGAAGAGATACGCCCTGGTT
CCTGGAACAATTGCTTTTACAGATGCACATATCGAGGTGGACATCACTTACGCTGAGTACTT
CGAAATGTCCGTTCCGTTGGCAGAAGCTATGAAACGATATGGGCTGAATACAAATCACAGAA
TCGTGCTATGCAGTGAAACTCTCTTCAATTCTTTATGCCGGTGTGGGCGCTTATTTATC
GGAGTTGCAGTTGCGCCCGCGAACGACATTTATAATGAACGTGAATTGCTCAACAGTATGGG
CATTTTCGACGCTACCGTGGTGTCTGTTTTCAAAAAGGGGTTGCAAAAAATTTTGAACGTGC
AAAAAAAGCTCCCAATCATCCAAAAAATTATTATCATGGATTCTAAAACGGATTACCAGGGA
TTTCAGTCGATGTACACGTTTCGTCACATCTCATCTACCTCCCGGTTTTAATGAATACGATTT
TGTGCCAGAGTCCTTCGATAGGGACAAGACAATTGCACTGATCATGAACTCCTCTGGATCTA
CTGGTCTGCCTAAAGGTGTCGCTCTGCCTCATAGAAGTGCCTGCGTGAGATTCTCGCATGCC
AGAGATCCTATTTTTGGCAATCAAATCAATTCGCGGATCTGCGATTTTAAGTGTGTTCCATT
CCATCACGGTTTTTGGAAATGTTTACTACACTCGGATATTTGATATGTGGATTTTCGAGTCGTCT
TAATGTATAGATTTGAAGAAGAGCTGTTTCTGAGGAGCCTTCAGGATTACAAGATTCAAAGT
GCGCTGCTGGTGCCAACCCTATTCTCCTTCTTCGCCAAAAGCACTCTGATTGACAAATACGA
TTTATCTAATTTACACGAAATTGCTTCTGGTGGCGCTCCCTCTCTAAGGAAGTCGGGGAAG
CGGTTGCCAAGAGGTTCCATCTGCCAGGTATCAGGCAAGGATATGGGCTCACTGAGACTACA
TCAGCTATTCTGATTACACCCGAGGGGATGATAAACCGGGCGCGGTGCGTTAAAGTTGTTCC
ATTTTTTGAAGCGAAGGTTGTGGATCTGGATACCGGGAAAACGCTGGGCGTTAATCAAAGAG
GCGAACTGTGTGTGAGAGGTCCATGATTATGTCCGGTTATGTAAACAATCCGGAAGCGACC
AACGCCTTGATTGACAAGGATGGATGGCTACATTCTGGAGACATAGCTTACTGGGACGAAGA
CGAACACTTCTTCATCGTTGACCGCCTGAAGTCTCTGATTAAAGTACAAAGGCTATCAGGTGG
CTCCCGCTGAATTTGGAATCCATCTTGCTCCAACACCCCAACATCTTCGACGCAGGTGTGCA
GGTCTTCCCGACGATGACGCCGGTGAACCTCCCGCCCGCGTTGTTGTTTTGGAGCACGGAAA
GACGATGACGGAAAAAGAGATCGTGGATTACGTGCGCAGTCAAGTAACAACCGCGAAAAAGT
TGCGCGGAGGAGTTGTGTTTTGTGGACGAAGTACCGAAAGGTCTTACCGGAAAACCTCGACGCA
AGAAAAATCAGAGAGATCCTCATAAAGGCCAAGAAGGGCGGAAAGATCGCCGTGTAATTCTA
GAGTCGGGGCGGCCCGCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGACAAA
CCACAAGTAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTA
TTTGTAACCATTTATAAGCTGCAATAAACAAGTTAAACAACAATTCATTCTATTTATGTT
TCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAACCTCTACAAATGTGGTA
AAATCGATAAGGATCAATTCGGCTTCGACTAGTACTGACGGACACACCGAAGCCCGCGGC
AACCCTCAGCGGATGCCCCGGGCTTCACGTTTTCCAGGTGAGAAGCGGTTTTTCGGGAGTA
GTGCCCCAACTGGGGTAACCTTTGAGTTCTCTCAGTTGGGGGCGTAGGGTCGCCGACATGAC
ACAAGGGGTTGTGACCGGGGTGGACACGTACGCGGGTGCTTACGACCGTCAGTCGCGCGAGC
GCGACTAGTACAAGCCGAATTGATCCGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCA
GCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATC
ATGCAACTCGTAGGACAGGTGCCGGCAGCGCTCTCCGCTTCTCGCTCACTGACTCGCTGC

GCTCGGTTCGTTCCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCC
ACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAA
CCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACA
AAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTT
CCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCCTGCCGCTTACCGGATACCTGTC
CGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTT
CGGTGTAGGTTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCCGACCGC
TGCGCCTTATCCGGTAACCTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACT
GGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCT
TGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTG
AAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCACCGCTGG
TAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAG
ATCCTTTGATCTTTTTCTACGGGGTCTGACGCTCAGTGGAAACGAAAACTCAGTTAAGGGATT
TTGGTTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTT
TAAATCAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTG
AGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTG
TAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGA
CCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCA
GAAGTGGTCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGA
GTAAGTAGTTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGT
GTCACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTTCCTCAACGATCAAGGCGAGTTA
CATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGA
AGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGT
CATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTTCTGAGAAT
AGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACAT
AGCAGAACTTTAAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGAT
CTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACCTGATCTTCAGCAT
CTTTTACTTTACCCAGCGTTTCTGGGTGAGCAAAAAACAGGAAGGCAAAATGCCGCAAAAAAG
GGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAG
CATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAAC
AAATAGGGGTTCGCGCGACATTTCCCCGAAAAGTGCCACCTGACGCGCCCTGTAGCGGCGCA
TTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGC
GCCGCTCCTTTTCGCTTTCTTCCCTTCTTTCTCGCCACGTTTCGCCGGCTTTCCCGTCAAG
CTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAA
AACTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCCC
TTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAACTGGAACAACACTCA
ACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTA
AAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTTAACAAAAATTAACGTTTACAAT
TTCCCATTCGCCATTCAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGC
TATTACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTTGGGA
GCGGCCGCAATAAAATATCTTTATTTTCATTACATCTGTGTGTTGGTTTTTTTGTGTGAATCG
ATAGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAACTAGCAAAATAGGC
TGTCCCCAGTGCAAGTGCAGGTGCCAGAACATTT

Fig. 11

pCMV-pur-attB (SEQ ID NO: 4)

CTAGAGTCGGGGCGGCCGCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGAC
AAACCACAACCTAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCT
TTATTTGTAACCATTATAAGCTGCAATAAACAAGTTAACAACAACAATTGCATTCAATTTAT
GTTTCAGGTTTCAGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAACCTCTACAAATGTG
GTAAAAATCGATAAGGATCAATTTCGGCTTCAGGTACCGTCGACGATGTAGGTCACGGTCTCGA
AGCCGCGGTGCGGGTGCCAGGGCGTGCCCTTGGGCTCCCCGGGCGCGTACTCCACCTCACCC
ATCTGGTCCATCATGATGAACGGGTTCGAGGTGGCGGTAGTTGATCCCGGCGAACGCGCGGGCG
CACCGGGAAGCCCTCGCCCTCGAAACCGCTGGGCGCGGTGGTCACGGTGAGCACGGGACGTG
CGACGGCGTTCGGCGGGTGCCGATACGCGGGGCGAGCGTCAGCGGGTCTCTCGACGGTCACGGCG
GGCATGTGCGACAGCCGAATTGATCCGTTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAG
CTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCGCACTTATGACTGTCTTCTTTATCA
TGCAACTCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCTCGCTCACTGACTCGCTGCG
CTCGGTTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCA
CAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAAC
CGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAA
AAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTC
CCCCTGGAAGCTCCCTCGTGCCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCC
GCCTTCTCCCTTCGGGAAGCGTGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTC
GGTGTAGGTTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTTCAGCCCGACCGCT
GCGCCTTATCCGGTAACCTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTG
GCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTT
GAAGTGGTGGCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGA
AGCCAGTTACCTTCGGA AAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCACCGCTGGT
AGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGA
TCCTTTGATCTTTTCTACGGGGTCTGACGCTTCAGTGGAACGAAACTCACGTTAAGGGATTT
TGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTA AAAATGAAGTTTT
AAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGA
GGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCTGT
AGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAC
CCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAG
AAGTGGTCCGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAG
TAAGTAGTTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTG
TCACGCTCGTCTTTTGGTATGGCTTCATTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTAC
ATGATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCTCCTCGATCGTTGTGAGAA
GTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTC
ATGCCATCCGTAAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATA
GTGTATGCGGCGACCGAGTTGCTCTTGCCGGCGTCAATAACGGGATAATACCGCGCCACATA
GCAGAACTTTAAAGTGCTCATCATTTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATC
TTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATC
TTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGG
GAATAAGGGCGACACGAAATGTTGAATACTCATACTCTTCTTTTCAATATTATTGAAGC
ATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAACA
AATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGCGCCCTGTAGCGGCGCAT
TAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCG
CCGCTCCTTTCGCTTTCTTCCCTTCTTCTCGCCACGTTGCGCGGCTTTCCCCGTCAAGC
TCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAA
AACTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCT
TTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAA
CCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTAA
AAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTAACA AAAATATTACGTTTACAATT
TCCCATTCGCCATTACGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCT
ATTACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTTGGAG
CGGCCGCAATAAAAATATCTTTATTTTATTACATCTGTGTGTTGGTTTTTTGTGTGAATCGA

TAGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAAACTAGCAAAATAGGCT
GTCCCCAGTGCAAGTGCCAGGTGCCAGAACATTTCTCTATCGATAGGTACCGAGCTCTTACGC
GTGCTAGCCCTCGAGCAGGATCTATACATTGAATCAATATTGGCAATTAGCCATATTAGTCA
TTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATATCAT
AATATGTACATTTATATTTGGCTCATGTCCAATATGACCGCCATGTTGACATTGATTATTGAC
TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCGCG
TTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCATTGACG
TCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGT
GGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTCCGC
CCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCAGTACATGACCTTA
CGGGACTTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCG
GTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCC
ACCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAATCAACGGGACTTTCCAAAATGT
CGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATAT
AAGCAGAGCTCGTTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACC
TCCATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGAAGCTCGACTCTAGGGGCTCGAGA
TCTGCGATCTAAGTAAGCTTGCATGCCTGCAGGTGCGCCGCCACGACCGGTGCCGCCACCAT
CCCCTGACCCACGCCCCCTGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGCCC
ACGGTGCGCCCTCGCCACCCGCGACGACGTCCCCCGGGCCGTACGCACCCTCGCCGCCGCGTT
CGCCGACTACCCCGCCACGCGCCACACCGTCGACCCGGACCGCCACATCGAGCGGGTCACCG
AGCTGCAAGAACTCTTCCTCACGCGCGTCGGGCTCGACATCGGCAAGGTGTGGGTTCGCGGAC
GACGGCGCCGCGGTGGCGGTCTGGACCACGCGGAGAGCGTCGAAGCGGGGGCGGTGTTCGC
CGAGATCGGCCCCGCGCATGGCCGAGTTGAGCGGTTCCTGGCTGGCCGCGCAGCAACAGATGG
AAGGCCTCCTGGCGCCGCACCGGCCCAAGGAGCCCGCGTGGTTTCTGGCCACCGTCGGCGTC
TCGCCCCGACCACAGGGCAAGGGTCTGGGACGCGCCGTGCTGCTCCCCGGAGTGAGGGCGGC
CGAGCGCGCCGGGGTGGCCGCTTCTTGGAGACCTCCGCGCCCCGCAACCTCCCCTTCTACG
AGCGGCTCGGCTTCACCGTCACCGCCGACGTGAGGTGCCCCGAAGGACCGCGCACCTGGTGC
ATGACCCGCAAGCCCGGTGCCTGACGCCCCGCCCCACGACCCGCGAGCGCCGACCGAAAAGGAG
CGCACGACCCCATGGCTCCGACCGAAGCCGACCCGGGCGGCCCGCGGACCCCGCACCCGCC
CCCGAGGCCACCGACT

Fig. 12

pCMV-pur-attP (SEQ ID NO: 5)

CTAGAGTCGGGGCGGCCGCGCTTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGAC
AAACCACAAC TAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCT
TTATTTGTAACCATTATAAGCTGCAATAAACAAGTTAACAACAACAATTGCATTCATTTTAT
GTTTCAGGTT CAGGGGGAGGTGTGGGAGGTTTTTTTAAAGCAAGTAAAACCTCTACAAATGTG
GTAAAATCGATAAGGATCAATTCGGCTTCGACTAGTACTGACGGACACACCGAAGCCCCGGC
GGCAACCCCTCAGCGGATGCCCCGGGGCTTCACGTTTCCCAGGTCAGAAGCGGTTTTCGGGA
GTACGTGCCCCAAGTGGGGTAACCTTTGAGTTTCTCTCAGTTGGGGGCGTAGGGTGCCCGACAT
GACACAAGGGGTTGTGACCGGGGTGGACACGTACGCGGGTGCTTACGACCGTCAGTCGCGCG
AGCGCGACTAGTACAAGCCGAATTGATCCGTGACCGATGCCCTTGAGAGCCTTCAACCCAG
TCAGCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTT
ATCATGCAACTCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGC
TGCGCTCGGTGCTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTA
TCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAG
GAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATC
ACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAAACCCGACAGGACTATAAAGATACCAGGCG
TTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCTGTTCCGACCCTGCCGCTTACCGGATACCT
GTCCGCCTTTCTCCCTTCGGGAAGCGTGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCA
GTTCCGGTGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTT CAGCCCGAC
CGCTGCGCCTTATCCGGTAACCTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCC
ACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGT
TCTTGAAGTGGTGGCCTAACTACGGCTACACTAGGAAGGACAGTATTTGGTATCTGCGCTCTG
CTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGC
TGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAG
AAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGAACGAAAACCTCACGTTAAGGG
ATTTTGGTCA TGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTA AAAATGAAG
TTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCA
GTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGT
GTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATAACCGCG
AGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGC
GCAGAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCT
AGAGTAAGTAGTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGT
GGTGTACGCTCGTCTGTTTGGTATGGCTTCATT CAGCTCCGGTTCCCAACGATCAAGGCGAG
TTACATGATCCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTC
AGAAGTAAGTTGGCCGAGTGTTATCACTACATGGTTATGGCAGCACTGCATAATTCTCTTAC
TGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAG
AATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCA
CATAGCAGAACTTTAAAAGTGCTCATCATTTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAG
GATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAG
CATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAA
AAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCTTTTCAATATTATTG
AAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATA
AACAAATAGGGGTTCCGCGCACATTTCCCGAAAAGTGCCACCTGACGCGCCCTGTAGCGGC
GCATTAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCT
AGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTTCGCCGGCTTTCCCCGTC
AAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCC
AAAAA ACTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGATTTTCG
CCCTTTGACGTTGGAGTCCACGTTCTTAAATAGTGGAAGTCTTGTTCCAAACCTGGAAACAAC
TCAACCCCTATCTCGGTCTATTTCTTTGATTTATAAGGGATTTTGCCGATTTGCGCCTATTGG
TTAAAAAATGAGCTGATTTAAACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTAC
AATTTCCCATTCGCCATT CAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTT
CGCTATTACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTT
GGAGCGGCCGCAATAAAATATCTTTATTTTCATTACATCTGTGTGTGGTTTTTTGTGTGAA
TCGATAGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAACTAGCAAAATA

GGCTGTCCCCAGTGCAAGTGCAGGTGCCAGAACATTTCTCTATCGATAGGTACCGAGCTCTT
ACGCGTGCTAGCCCTCGAGCAGGATCTATACATTGAATCAATATTGGCAATTAGCCATATTA
GTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATA
TCATAATATGTACATTTATATTGGCTCATGTCCAATATGACCGCCATGTTGACATTGATTAT
TGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTC
CGCGTTACATAACTTACGGTAAATGGCCCCGCTGGCTGACCGCCCAACGACCCCCGCCATT
GACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAAT
GGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGT
CCGCCCCCTATTGACGTCAATGACGGTAAATGGCCCCGCTGGCATTATGCCCAGTACATGAC
CTTACGGGACTTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGA
TGC GGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTTGACTCACGGGGATTTCCAAGT
CTCCACCCCATTTGACGTCAATGGGAGTTTTGTTTTTGGCACCAAAATCAACGGGACTTTCCAAA
ATGTCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCT
ATATAAGCAGAGCTCGTTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTT
GACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGAAGCTCGACTCTAGGGGCTC
GAGATCTGCGATCTAAGTAAGCTTGCATGCCTGCAGGTGCGCCGCCACGACCGGTGCCGCCA
CCATCCCCTGACCCACGCCCCTGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAA
GCCCACGGTGCGCCTCGCCACCCGCGACGACGTCCCCCGGGCCGTACGCACCCCTCGCCGCCG
CGTTTCGCCGACTACCCCGCCACGCGCCACACCGTCGACCCGGACCGCCACATCGAGCGGGTC
ACCGAGCTGCAAGAACTCTTCCTCACGCGCGTCGGGCTCGACATCGGCAAGGTGTGGGTTCGC
GGACGACGGCGCCGCGGTGGCGGTCTGGACCACGCCGGAGAGCGTCGAAGCGGGGGCGGTGT
TCGCCGAGATCGGCCCGCGCATGGCCGAGTTGAGCGGTTCCCGGCTGGCCGCGCAGCAACAG
ATGGAAGGCCTCCTGGCGCCGCACCGGCCCAAGGAGCCCGCGTGGTTTCTTGGCCACCGTCGG
CGTCTCGCCCGACCAACAGGGCAAGGCTCTGGGCAGCGCCGTCGTGCTCCCCGGAGTGGAGG
CGGCCGAGCGCGCCGGGGTGCCCGCCTTCTTGGAGACCTCCGCGCCCCGCAACCTCCCCTTC
TACGAGCGGCTCGGCTTCACCGTCACCGCCGACGTGAGGTGCCCGAAGGACCGCGCACCTG
GTGCATGACCCGCAAGCCCGGTGCCTGACGCCCGCCCCACGACCCGAGCGCCCGACCGAAA
GGAGCGCACGACCCCATGGCTCCGACCGAAGCCGACCCGGGCGGCCCGCCGACCCCGCACCC
CGCCCCCGAGGCCACCGACT

Fig. 13

pCMV-EGFP-attB (SEQ ID NO: 6)

CTAGAGTCGGGGCGGCCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGAC
AAACCACAAC TAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCT
TTATTTGTAAACCATTATAAGCTGCAATAAACAAAGTTAACAACAACAATTGCATTCATTTTAT
GTTTCAGGTT CAGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAACCTCTACAAATGTG
GTAAATCGATAAGGATCAATTCGGCTTCAGGTACCGTCGACGATGTAGGTACGGTCTCGA
AGCCGCGGTGCGGGTGCCAGGGCGTGCCCTTGGGCTCCCCGGGCGCGTACTCCACCTCACCC
ATCTGGTCCATCATGATGAACGGGTGCGAGGTGGCGGTAGTTGATCCCGGCGAACGCGCGGCG
CACCGGGAAGCCCTCGCCCTCGAAACCGCTGGGCGCGGTGGTCACGGTGAGCACGGGACGTG
CGACGCGTTCGGCGGGTGCGGATACGCGGAGCCAGCGTCAGCGGGTTCTCGACGGTCACGGCG
GGCATGTGACAGCCGAATTGATCCGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAG
CTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCA
TGCAACTCGTAGGACAGGTGCCGGCAGCGCTCTTCGGCTTCTCGCTCACTGACTCGCTGCG
CTCGGTCTGTTCCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCA
CAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAAC
CGTAAAAAGCCCGTGTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAA
AAATCGACGCTCAAGTCAGAGGTGGCGAAACCCAGCAGGACTATAAAGATACAGGCGTTTC
CCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCC
GCCTTCTCCTTCCGGGAAGCGTGGCGCTTCTCAATGCTCACGCTGTAGGTATCTCAGTTC
GGTGTAGGTCTGTTCCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTACGCCGACCGCT
GCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTG
GCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTT
GAAGTGGTGGCTAACCTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGA
AGCCAGTTACCTTCGGAAAAAGAGTTGGTATGCTCTTGATCCGGCAAAACAAACCCGCTGGT
AGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGA
TCCTTTGATCTTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAACTCACGTTAAGGGATTT
TGGTCATGAGATTATCAAAAAGGATCTTACCTAGATCCTTTTAAATTAAAAATGAAGTTTT
AAATCAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGA
GGCACCTATCTCAGCGATCTGTCTATTTCTGTTTCAATCCATAGTTGCCTGACTCCCCGTCTGT
AGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAC
CCACGCTCACCGGCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGCAG
AAGTGGTCTTCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAG
TAAGTAGTTCCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTG
TCACGCTCGTCTGTTTGGTATGGCTTCATTACGCTCCGGTTCCCAACGATCAAGGCGAGTTAC
ATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCTCGATCGTTGTGAGAA
GTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTC
ATGCCATCCGTAAGATGCTTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTTCTGAGAATA
GTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATA
GCAGAACTTTAAAAGTGCTCATCATTGGAACCGTTCTTCGGGGCGAAACTCTCAAGGATC
TTACCGCTGTTGAGATCCAGTTGATGTAACCCACTCGTGACCCAACTGATCTTCAGCATC
TTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAAACAGGAAGGCAAAATGCCGCAAAAAAGG
GAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCTTTTCAATATTATTGAAGC
ATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAACA
AATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGCGCCCTGTAGCGGCGCAT
TAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTGCCAGCGCCCTAGCG
CCCGCTCCTTTCGCTTCTTCCCTTCTTCTCGCCACGTTCCGCGGCTTTCCTCCGTCAGC
TCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAA
AACTTGATTAGGGTGATGGTTACGTTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCT
TTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCAAACTGGAACAACACTCAA
CCATATCGGTCTATTCTTTTGAATTTAAGGGATTTTGCCGATTTCCGGCTATTGGTTAA
AAATGAGTGATGATTTAACAATAATTTAACGCAATTTTAAACAAATATTAACCTTTACAATT
TCCCATTCGCCATTACAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCT
ATTACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTTGGAG
CGGCCGCAATAAAATATCTTTATTTTATTACATCTGTGTGTTGGTTTTTTGTGTGAATCGA

TAGTACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAAACTAGCAAAATAGGCT
GTCCCCAGTGCAAGTGCAGGTGCCAGAACATTTCTCTATCGATAGGTACCGAGCTCTTACGC
GTGCTAGCCCTCGAGCAGGATCTATACATTGAATCAATATTGGCAATTAGCCATATTAGTCA
TTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATATCAT
AATATGTACATTTATATTGGCTCATGTCCAATATGACCGCCATGTTGACATTGATTATTGAC
TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCG
TTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCATTGACG
TCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCATTGACGTCAATGGGT
GGAGTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTCCGC
CCCCATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGACCTTA
CGGGACTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCG
GTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCC
ACCCCATTTGACGTCAATGGGAGTTTGTGTTTTGGCACAAAATCAACGGGACTTTCCAAAATGT
CGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATAT
AAGCAGAGCTCGTTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACC
TCCATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGAAGCTCGACTCTAGGGGCTCGAGA
TCCCCGGGTACCGGTGCGCCACCATGGTGAGCAAGGGCGAGGAGCTGTTACCGGGGTGGTG
CCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGC
GAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCC
CGTGCCCTGGCCCAACCTCGTGACCACCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACC
CCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAG
CGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGG
CGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCC
TGGGGCACAAGCTGGAGTACAACACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAG
AAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCT
CGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACC
ACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTC
CTGCTGGAGTTCTGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAAAG
CGGCCGCTCGAGCATGCAT

Fig. 14

p-12.0-lys-LSPIFNMM-CMV-pur-attB (SEQ ID NO: 7)

GGGCTGCAGGAATTTCGATTGCCGCCTTCTTTGATATTCACCTCTGTTGTATTTTCATCTCTTCT
TGCCGATGAAAGGATATAACAGTCTGTATAACAGTCTGTGAGGAAATACTTGGTATTTCTTC
TGATCAGTGTTTTTATAAGTAATGTTGAATATTGGATAAGGCTGTGTGTCCTTTGTCTTGGG
AGACAAAGCCACAGCAGGTGGTGGTTGGGGTGGTGGCAGCTCAGTGACAGGAGAGGTTTTT
TTGCCCTGTTTTTTTTTTTTTTTTTTTTTAAAGTAAGGTGTTCTTTTTCTTAGTAAATTTT
CTACTGGACTGTATGTTTTGACAGGTCAGAAACATTTCTTCAAAGAAGAACCTTTTGGAAA
CTGTACAGCCCTTTTCTTTCATTCCTTTTGTCTTCTGTGCCAATGCCTTTGGTTCTGATT
GCATTATGAAAACGTTGATCGGAACCTTGAGGTTTTTATTTATAGTGTGGCTTGAAAGCTTG
GATAGCTGTTGTTACACGAGATACCTTATTAAGTTTAGGCCAGCTTGATGCTTTATTTTTTC
CCTTTGAAGTAGTGAGCGTTCTCTGGTTTTTTTCTTTGAAACTGGTGAGGCTTAGATTTTT
CTAATGGGATTTTTTACCTGATGATCTAGTTGCATACCCAAATGCTTGTAATGTTTTCTTA
GTTAACATGTTGATAACTTCGGATTTACATGTTGTATATACTTGTCTATCTGTGTTCTAGTA
AAAATATATGGCATTATAGAAATACGTAATTCCTGATTTCTTTTTTTTTTATCTATGCT
CTGTGTGTACAGGTCAAACAGACTTCACCTCCTATTTTTATTTATAGAATTTTATATGCAGTC
TGTCGTTGGTTCTTGTGTTGTAAGGATACAGCCTTAAATTTCTAGAGCGATGCTCAGTAAG
GCGGTTGTGCATGGGTTCAAATGTAAAACGGGCACGTTTGGCTGCTGCCTTCCCAGATC
CAGGACACTAAACTGCTTCTGCACTGAGGTATAAATCGCTTCAGATCCCAGGGAAGTGCAGA
TCCACGTGCATATTCTTAAAGAAGAATGAATACTTTCTAAAATATTTTGGCATAGGAAGCAA
GCTGCATGGATTTGTTTGGGACTTAAATTTTGGTAACGGAGTGCATAGGTTTTAAACAC
AGTTGCAGCATGCTAACGAGTCACAGCGTTTATGTCAGAAAGTGATGCCTGGATGCCTGTTGCA
GCTGTTTACGGCACTGCCTTGCAGTGAGCATTGCAGATAGGGGTGGGGTGCTTTGTGTCTG
TTCCACACGCTGCCACACAGCCACCTCCCGGAACACATCTCACCTGCTGGGTACTTTTTCAA
ACCATCTTAGCAGTAGTAGATGAGTTACTATGAAACAGAGAAGTTCCCTCAGTTGGATATTCT
CATGGGATGTCTTTTTTCCCATGTTGGGCAAAGTATGATAAAGCATCTCTATTTGTAAATTA
TGCATTGTTAGTTTCTGAATCCTTTCTATAGCACCACTTATTGCAGCAGGTGTAGGCTCTG
GTGTGGCCTGTGTCTGTGCTTCAATCTTTTAAAGCTTCTTTGGAATACTGACTTGATTG
AAGTCTCTTGAAGATAGTAAACAGTACTTACCTTTGATCCCAATGAAATCGAGCATTTCAGT
TGTAAGAAGAAATCCGCCTATTCATACCATGTAATGTAATTTTACACCCCCAGTGCTGACACT
TTGGAATATATTCAAGTAATAGACTTTGGCCTCACCTCTTGTGTACTGTATTTGTAAATAG
AAAATATTTTAAACTGTGCATATGATTATTACATTATGAAAGAGACATTCTGCTGATCTTCA
AATGTAAGAAAATGAGGAGTGCGTGTGCTTTTATAAATACAAGTGATTGCAAAATAGTGCA
GTGTCCTTAAAAAAGTAATATAAAAAGGACCAGGTGTTTTACAAGTGAAT
ACATTCCTATTGTTAAACAGTTACATTTTATGAAGATTACCAGCGCTGCTGACTTTCTAA
ACATAAGGCTGTATTGTCTTCTGTACCATTTGCATTTCTCATTCCCAATTTGCACAAGGAT
GTCTGGGTAAACTATTCAAGAAATGGCTTTGAAATACAGCATGGGAGCTTGTCTGAGTTGGA
ATGCAGAGTTGCACTGCAAAATGTCAGGAAATGGATGTCTCTCAGAATGCCCACTCCAAAG
GATTTTATATGTGTATATAGTAAGCAGTTTCTGATTCCAGCAGGCCAAAGAGTCTGCTGAA
TGTTGTGTTGCCGGAGACCTGTATTTCTCAACAAGGTAAGATGGTATCCTAGCAACTGCGGA
TTTTAATACATTTTTCAGCAGAAGTACTTAGTTAATCTCTACCTTTAGGGATCGTTTCATCAT
TTTTAGATGTTATACTTGAAATACTGCATAACTTTTAGCTTTCATGGGTTCTTTTTTTTCAG
CCTTTAGGAGACTGTTAAGCAATTTGCTGTCCAACCTTTTGTGTTGGTCTTAAACTGCAATAG
TAGTTTACCTTGTATTGAAGAAATAAAGACCATTTTATATTAATAAATACTTTTGTCTGTC
TTCATTTTGACTTGTCTGATATCCTTGCAGTGCCCATTTATGTCAGTTCTGTGAGATATTGAG
ACATCAAACTTAACGTGAGCTCAGTGGAGTTACAGCTGCGGTTTTTGATGCTGTTATTATTT
CTGAACTAGAAATGATGTTGTCTTCATCTGCTCATCAAACTTCATGCAGAGTGTAAGGC
TAGTGAGAAATGCATACATTTATTGATACTTTTTTAAAGTCAACTTTTTATCAGATTTTTTT
TTCATTTTGGAAATATATTGTTTTCTAGACTGCATAGCTTCTGAATCTGAAATGCAGTCTGAT
TGGCATGAAGAAGCACAGCACTCTTCATCTTACTTAACTTCATTTTGGAAATGAAGGAAGTT
AAGCAAGGGCACAGGTCCATGAAATAGAGACAGTGCGCTCAGGAGAAAGTGAACCTGGATTT
CTTTGGCTAGTGTTCTAAATCTGTAGTGAGGAAAGTAACACCCGATTCTTTGAAAGGGCTCC
AGCTTTAATGCTTCCAAATTGAAGGTGGCAGGCAACTTGGCCACTGGTTATTTACTGCATTA
TGTCTCAGTTTCGCAGCTAACCTGGCTTCTCCACTATTGAGCATGGACTATAGCCTGGCTTC
AGAGGCCAGGTGAAGGTTGGGATGGGTGGAAGGAGTGCTGGGCTGTGGCTGGGGGGACTGTG

GGGACTCCAAGCTGAGCTTGGGGTGGGCAGCACAGGGAAGTGTGGGTAACATTTTTTAAG
TACTGTGTTGCAAACGTCTCATCTGCAAATACGTAGGGTGTGTACTCTCGAAGATTAACAGT
GTGGGTTTCAAGTAATATATGGATGAATTCACAGTGGAAGCATTCAGGGTAGATCATCTAACG
ACACCAGATCATCAAGCTATGATTGGAAGCGGTATCAGAAGAGCGAGGAAGGTAAGCAGTCT
TCATATGTTTTCCCTCCACGTAAAGCAGTCTGGGAAAGTAGCACCCCTTGAGCAGAGACAAG
GAAATAATTCAAGGAGCATGTGCTAGGAGAACTTTCTTGCTGAATTCTACTTGCAAGAGCTTT
GATGCCTGGCTTCTGGTGCCTTCTGCAGCACCTGCAAGGCCAGAGCCTGTGGTGAGCTGGA
GGGAAAGATTCTGCTCAAGTCCAAGCTTCAGCAGGTCATTGTCTTTGCTTCTTCCCCAGCA
CTGTGCAGCAGAGTGGAAGTGTGCGAAGCCTCCTGTCCACTACCTGTTGCTGCAGGCAGA
CTGCTCTCAGAAAAAGAGAGCTAACTCTATGCCATAGTCTGAAGGTAAAATGGGTTTTAAAA
AAGAAAACACAAAGGCCAAACCGGCTGCCCATGAGAAGAAAGCAGTGGTAAACATGGTAGA
AAAGGTGCAGAAGCCCCAGGCAGTGTGACAGGCCCTCCTGCCACCTAGAGGCGGGAAACAA
GCTTCCCTGCCTAGGGCTCTGCCGCGAAGTGCGTGTCTTTGGTGGGTTTTGTTTGGCGT
TTGGTTTTGAGATTTAGACACAAGGGAAGCCTGAAAGGAGGTGTTGGGCACTATTTTGGTTT
GTAAAGCCTGACTTCAAATATATATTTTGTGAGGTGAGTGTAGCGAATTGGCCAATTTAAAA
TAAAGTTGCAAGAGATTGAAGGCTGAGTAGTTGAGAGGGTAACACGTTTAATGAGATCTTCT
GAACTACTGCTTCTAAACACTTGTGTGAGTGGTGAGACCTTGGATAGGTGAGTGCTCTTGT
TACATGTCTGATGCACTTGCTTGTCTTTTCCATCCACATCCATGCATTCCACATCCACGCA
TTTGTCACTTATCCCATATCTGTCTATCTGACATACCTGTCTCTTCGTCACTTGGTCAGAA
GAAACAGATGTGATAATCCCCAGCCGCCCAAGTTTGAGAAGATGGCAGTTGCTTCTTTCCC
TTTTCTCTGCTAAGTAAGGATTTTCTCTGGCTTTGACACCTCACGAAATAGTCTTCTCTGCC
TTACATTCTGGGCATTATTTCAAATATCTTTGGAGTGCGCTGCTCTCAAGTTTGTGTCTTCC
TACTCTTAGAGTGAATGCTCTTAGAGTGAAAGAGAAGGAAGAGAAGATGTTGGCCGAGTTC
TCTGATGAACACACCTCTGAATAATGGCCAAAGGTGGGTGGGTTTCTCTGAGGAACGGGCAG
CGTTTGCCTCTGAAAGCAAGGAGCTCTGCGGAGTTGCAGTTATTTTGCAACTGATGGTGAA
CTGGTGCTTAAAGCAGATTCCCTAGGTTCCCTGCTACTTCTTTTCTTCTTGGCAGTCAGTT
TATTTCTGACAGACAAACAGCCACCCCCACTGCAGGCTTAGAAAGTATGTGGCTCTGCCTGG
GTGTGTTACAGCTCTGCCCTGGTGAAAGGGGATTAAAACGGGCACCATTTCATCCCAAACAGG
ATCCTCATTTCATGGATCAAGCTGTAAGGAACCTGGGCTCCAACCTCAAACATTAATTGGAG
TACGAATGTAATTAATAACTGCATTCTCGCATTCCTAAGTCATTTAGTCTGGACTCTGCAGCA
TGTAGGTGCGCAGCTCCCACTTTCTCAAAGACCACTGATGGAGGAGTAGTAAAAATGGAGAC
CGATTTCAGAACAAACCAACGGAGTGTTGCCGAAGAACTGATGGAAATAATGCATGAATTGTG
TGGTGGACATTTTTTTTAAATACATAAACTACTTCAAATGAGGTGCGAGAAGGTCAGTGTTT
TATTAGCAGCCATAAAACCAGGTGAGCGAGTACCATTTTTCTCTACAAGAAAAACGATTTCT
AGCTGTGCGTAAGTATAAGTTCTCCATAGCGGCTGAAGCTCCCCCTGGCTGCCATCT
CAGCTGGAGTGAGTGCCATTTCCCTGGGGTTTCTCTCACAGCAGTAATGGGACAATACTTC
ACAAAAATTCTTTCTTTTCTGTCTATGTGGGATCCCTACTGTGCCCTCCTGGTTTTTACGTTA
CCCCCTGACTGTTCCATTTCAGCGGTTTGGAAGAGAAAAAGAATTTGGAAATAAAACATGTC
TACGTTATCACCTCCTCCAGCATTTTGGTTTTTAAATTATGTCAATAACTGGCTTAGATTTGG
AAATGAGAGGGGGTGGGTGTATTACCGAGGAACAAAGGAAGGCTTATATAAACTCAAGTCT
TTTATTTAGAGAACTGGCAAGCTGTCAAAAACAAAAGGCCCTTACCACCAAATTAAGTGAAT
AGCCGCTATAGCCAGCAGGGCCAGCACGAGGGATGGTGCACTGCTGGCACTATGCCACGGCC
TGCTTGTGACTCTGAGAGCAACTGCTTTGGAAATGACAGCACTTGGTGCAATTTCCCTTGT
TCAGAATGCGTAGAGCGTGTGCTTGGCGACAGTTTTTCTAGTTAGGCCACTTCTTTTTTCT
TCTCTCCTCATTCTCCTAAGCATGTCTCCATGCTGGTAATCCAGTCAAGTGAACGTTCAAA
CAATGAATCCATCACTGTAGGATTCTCGTGGTGATCAAACTTTGTGTGAGGTCTATAAAAT
ATGGAAGCTTATTTATTTTTCGTTCTTCCATATCAGTCTTCTCTATGACAATTCACATCCAC
CACAGCAAAATTAAGGTGAAGGAGGCTGGTGGATGAAGAGGCTTCTAGCTTTACGTTCT
TCCTTGCAAGGCCACAGGAAATGCTGAGAGCTGTAGAATACAGCCTGGGGTAAGAAGTTCA
GTCTCCTGCTGGGACAGCTAACCGCATCTTATAACCCCTTCTGAGACTCATCTTAGGACCAA
ATAGGGTCTATCTGGGGTTTTTGTTCCTGCTGTTCTCCTGGAAGGCTATCTCACTATTTCA
CTGCTCCACGGTTACAAACCAAGATACAGCCTGAATTTTTTCTAGGCCACATTACATAAA
TTTGACCTGGTACCAATATTGTTCTCTATATAGTTATTTCTTCCCCACTGTGTTTAAACCC
TTAAGGCATTCAGAACAACTAGAATCATAGAATGGTTTGGATTGGAAGGGGCCCTTAAACATC

ATCCATTTCCAACCCTCTGCCATGGGCTGCTTGCCACCCACTGGCTCAGGCTGCCCAGGGCC
CCATCCAGCCTGGCCTTGAGCACCTCCAGGGATGGGGCACCCACAGCTTCTCTGGGCAGCCT
GTGCCAACACCTCACCCTCTCTGGGTAAAGAATTCTCTTTTAACATCTAATCTAAATCTCT
TCTCTTTTAGTTTAAAGCCATTCTCTTTTCCCGTTGCTATCTGTCCAAGAAATGTGTATT
GGTCTCCCTCCTGCTTATAAGCAGGAAGTACTGGAAGGCTGCAGTGAGGTCTCCCCACAGCC
TTCTCTTCTCCAGGCTGAACAAGCCCAGCTCCTTCAGCCTGTCTTCGTAGGAGATCATCTTA
GTGGCCCTCCTCTGGACCCATTCCAACAGTTCCACGGCTTTCTTGTGGAGCCCCAGGTCTGG
ATGCAGTACTTCAGATGGGGCCTTACAAAGGCAGAGCAGATGGGGACAATCGCTTACCCCTC
CCTGCTGGCTGCCCCTGTTTTGATGCAGCCCAGGGTACTGTTGGCCTTTCAGGCTCCCAGAC
CCCTTGCTGATTTGTGTCAAGCTTTTTCATCCACCAGAACCACGCTTCCTGGTTAATACTTC
TGCCCTCACTTCTGTAAGCTTGTTTTAGGAGACTTCCATTCTTTAGGACAGACTGTGTTACA
CCTACCTGCCCTATTCTTGCATATATACAGATTTTCACTTCATGTTTCTGTAAACAGGACAGAAT
ATGTATTCTCTAACAAAAATACATGCAGAAATTCCTAGTGCCATCTCAGTAGGGTTTTTCATG
GCAGTATTAGCACATAGTCAATTTGCTGCAAGTACCTTCCAAGCTGCGGCCTCCATAAATC
CTGTATTTGGGATCAGTTACCTTTTGGGGTAAGCTTTTGTATCTGCAGAGACCCTGGGGGT
CTGATGTGCTTCAGCTCTGCTCTGTTCTGACTGCACCATTTTCTAGATCACCCAGTTGTTC
TGTACAACCTTCTTGTCTCCATCCTTTCCAGCTTGTATCTTTGACAAATACAGGCCTATT
TTTGTGTTTGTCTCAGCAGCCATTTAATTCCTCAGTGTCTCTTGTCTGTTGATGCCACTG
GAACAGGATTTTCAGCAGTCTTGCAAAGAACATCTAGCTGAAAACCTTCTGCCATTCAATAT
TCTTACCAGTTCTTCTGTTTGAGGTGAGCCATAAATTACTAGAACCTTCGTCACTGGACAAGT
TTATGCATTTTATTACTTCTATTATGTACTTACTTTGACATAACACAGACACGCACATATTT
TGCTGGGATTTCCACAGTGTCTCTGTGCTTTCACATGGTTTTACTGTCTACTTCCGTTAT
AACCTTGGCAATCTGCCCAGCTGCCCATCACAAGAAAAGAGATTCTTTTTTTATTACTTCTC
TTCAGCCAATAAACAAAATGTGAGAAGCCCCAACAAAGAACTTGTGGGGCAGGCTGCCATCAA
GGGAGAGACAGCTGAAGGGTGTGTAGCTCAATAGAATTAAGAAATAATAAAGCTGTGTGAG
ACAGTTTTGCTGATTATACAGGCACGCCCCAAGCCAGAGAGGCTGTCTGCCAAGGCCACC
TTGCAGTCTTGGTTTGTAAAGATAAGTCATAGGTAACCTTTTCTGGTGAATTGCGTGGAGAAT
CATGATGGCAGTTCTTGCTGTTTACTATGGTAAGATGCTAAAATAGGAGACAGCAAAGTAAC
ACTTGCTGCTGTAGGTGCTCTGCTATCCAGACAGCGATGGCACTCGCACACCAAGATGAGGG
ATGCTCCCAGCTGACGGATGCTGGGGCAGTAACAGTGGGTCCCATGCTGCCTGCTCATTAGC
ATCACCTCAGCCCTCACCAGCCCATCAGAAGGATCATCCCAAGCTGAGGAAAAGTTGCTCATC
TTCTTCACATCATCAAACCTTTGGCCTGACTGATGCCTCCCGGATGCTTAAATGTGGTCACT
GACATCTTTATTTTTCTATGATTTCAAGTCAGAACCTCCGGATCAGGAGGGAACACATAGTG
GGAATGTACCCTCAGCTCCAAGGCCAGATCTTCTTCAATGATCATGCATGCTACTTAGGAA
GGTGTGTGTGTGTGAATGTAGAATTGCCTTTGTATTTTTTCTTCTGCTGTGAGGAACATT
TTGAATACCAGAGAAAAAGAAAAGTGCTCTTCTTGGCATGGGAGGAGTTGTACACTTGCAA
AATAAAGGATGCAGTCCCAAATGTTTCAATCTCAGGGTCTGAAGGAGGATCAGAACTGTG
TATACAATTTCAAGGCTTCTCTGAATGCAGCTTTTGAAAGCTGTTTCTTGGCCGAGGCAGTACT
AGTCAGAACCCCTCGGAAACAGGAACAAATGTCTTCAAGGTGCAGCAGGAGGAAACACCTTGC
CCATCATGAAAGTGAATAACCACTGCCGCTGAAGGAATCCAGCTCCTGTTTGAGCAGGTGCT
GCACACTCCCACACTGAAACAACAGTTCATTTTTATAGGACTTCCAGGAAGGATCTTCTTCT
TAAGCTTCTTAATTATGGTACATCTCCAGTTGGCAGATGACTATGACTACTGACAGGAGAAT
GAGGAACTAGCTGGGAATATTTCTGTTTGACCACCATGGAGTCACCCATTTCTTTACTGGTA
TTTGGAATAATAAATTCTGAATTGCAAAGCAGGAGTTAGCGAAGATCTTCATTTCTTCCATG
TTGGTGACAGCACAGTTCTGGCTATGAAAGTCTGCTTACAAGGAAGAGGATAAAAATCATAG
GGATAATAAATCTAAGTTTGAAGACAATGAGGTTTTAGCTGCATTTGACATGAAGAAATTGA
GACCTCTACTGGATAGCTATGGTATTTACGTGTCTTTTTTGCTTAGTTACTTATTGACCCAG
CTGAGGTCAAGTATGAACTCAGGTCTCTCGGGCTACTGGCATGGATTGATTACATACAACCTG
TAATTTTAGCAGTGATTTAGGGTTTATGAGTACTTTTGCAGTAAATCATAGGGTTAGTAATG
TTAATCTCAGGGAAAAAAGCCAAACCTGACAGACATCCAGCTCAGGTGGAAATC
AAGGATCACAGCTCAGTGCGGTCCCAGAGAACACAGGGACTCTTCTCTTAGGACCTTTATGT
ACAGGGCCTCAAGATAACTGATGTTAGTCAGAAGACTTTCCATTCTGGCCACAGTTTCACTG
AGGCAATCCTGGAATTTTCTCTCCGCTGCACAGTTCCAGTCATCCAGTTTGTACAGTTCTG
GCACTTTTTGGGTGAGGCCGTGATCCAAGGAGCAGAAGTTCCAGCTATGGTCAGGGAGTGCC

TGACCGTCCCAACTCACTGCACTCAAACAAAGGGCGAAACCACAAGAGTGGCTTTTGTGAA
TTGCAGTGTGGCCAGAGGGGCTGCACCAGTACTGGATTGACCACGAGGCAACATTAATCCT
CAGCAAGTGCAATTTGCAGCCATTAAATTTGAACTAACTGATACTACAATGCAATCAGTATCA
ACAAGTGGTTTGGCTTGAAGATGGAGTCTAGGGGCTCTACAGGAGTAGCTACTCTCTAATG
GAGTTGCATTTTGAAGCAGGACACTGTGAAAAGCTGGCCTCCTAAAGAGGCTGCTAAACATT
AGGGTCAATTTTCCAGTGCACCTTTCTGAAGTGTCTGCAGTTCCCCATGCAAAGCTGCCCAA
CATAGCACTTCCAATTGAATACAATTATATGCAGGCGTACTGCTTCTTGCCAGCACTGTCCT
TCTCAAATGAACTCAACAAACAATTTCAAAGTCTAGTAGAAAGTAACAAGCTTTGAATGTCA
TTAAAAAGTATATCTGCTTTTCAGTAGTTTCAGCTTATTTTATGCCCACTAGAAACATCTTGTAC
AAGCTGAACACTGGGGCTCCAGATTAGTGGTAAAACCTACTTTATACAATCATAGAATCATA
GAATGGCCTGGGTTTGAAGGGACCCCAAGGATCATGAAGATCCAACACCCCCGCCACAGGCA
GGGCCACCAACCTCCAGATCTGGTACTAGACCAGGCAGCCCAGGGCTCCATCCAACCTGGCC
ATGAACACCTCCAGGGATGGAGCATCCACAACCTCTCTGGGCAGCCTGTGCCAGCACCTCAC
CACCCTCTCTGTGAAGAACTTTCCCTGACATCCAATCTAAGCCTTCCCTCCTTGAGGTTAG
ATCCACTCCCCCTTGTGCTATCACTGTCTACTCTTGTAAAAAGTTGATTCTCTCTTTTGTG
GAAGGTTGCAATGAGGTCTCCTTGCAGCCTTCTTCTCTTCTGCAGGATGAACAAGCCCAGCT
CCCTCAGCCTGTCTTTATAGGAGAGGTGCTCCAGCCCTCTGATCATCTTTGTGGCCCTCCTC
TGGACCCGCTCCAAGAGCTCCACATCTTTCCTGTACTGGGGGCCCCAGGCCTGAATGCAGTA
CTCCAGATGGGGCCTCAAAGAGCAGAGTAAAGAGGGGACAATCACCTTCCCTCACCTGCTGG
CCAGCCCTCTTCTGATGGAGCCCTGGATACAACCTGGCTTTCTGAGCTGCAACTTCTCCTTAT
CAGTTCCACTATTAAACAGGAACAATAACAAGGTGCTGATGGCCAGTGCAGAGTTTTTCA
CACTTCTTCATTTTCGTTAGATCTTAGATGAGGAACGTTGAAGTTGTGCTTCTGCGTGTGCTT
CTTCTCTCTCAAATACTCCTGCCTGATACCTCACCCACCTGCCACTGAATGGCTCCATGGC
CCCCTGCAGCCAGGGCCCTGATGAACCCGGCCTGCTTCAGATGCTGTTTAAATAGCACAGTA
TGACCAAGTTGCACCTATGAATACACAAACAATGTGTTGCATCCTTCAGCACTTGAGAAGAA
GAGCCAAATTTGCATTGTTCAGGAAATGGTTTAGTAATTCTGCCAATTAACACTTGTTTTATCT
ACCATGGCTGTTTTTATGGCTGTTAGTAGTGGTACACTGATGATGAACAATGGCTATGCAGT
AAAATCAAGACTGTAGATATTGCAACAGACTATAAAATTCCTCTGTGGCTTAGCCAAATGTGG
TACTTCCCACATTTGTATAAGAAATTTGGCAAGTTTAGAGCAATGTTTGAAGTGTGGGAAAT
TTCTGTATACTCAAGAGGGCGTTTTTGGACAACGTAGAACAGAGGAATCAAAAGGGGGTGGG
AGGAAGTTAAAGAAGAGGCAGGTGCAAGAGAGCTTGCAGTCCCGCTGTGTGTACGACACTG
GCAACATGAGGTCTTTGCTAATCTTGGTGCTTTGCTTCCTGCCCTGGCTGCCTTAGGGTGC
GATCTGCCTCAGACCCACAGCCTGGGCAGCAGGAGGACCTGATGCTGCTGGCTCAGATGAG
GAGAATCAGCCTGTTTAGCTGCCTGAAGGATAGGCACGATTTTGGCTTTCCTCAAGAGGAGT
TTGGCAACCCAGTTTCAGAAGGCTGAGACCATCCCTGTGCTGCACGAGATGATCCAGCAGATC
TTTAACCTGTTTAGCACCAAGGATAGCAGCGCTGCTTGGGATGAGACCCCTGCTGGATAAGTT
TTACACCGAGCTGTACCAGCAGCTGAACGATCTGGAGGCTTGCCTGATCCAGGGCGTGGGCG
TGACCGAGACCCCTCTGATGAAGGAGGATAGCATCCTGGCTGTGAGGAAGTACTTTTCAAGG
ATCACCTGTACCTGAAGGAGAAGAAGTACAGCCCTGCGCTTGGGAAGTCGTGAGGGCTGA
GATCATGAGGAGCTTTAGCCTGAGCACCAACCTGCAAGAGAGCTTGAGGTCTAAGGAGTAAA
AAGTCTAGAGTCGGGGCGGCCGCGCTTCGAGCAGACATGATAAGATACATTGATGAGTTT
GGACAAACCACAACCTAGAATGCAGTGAAAAAAATGCTTTATTTGTGAAATTTGTGATGCTAT
TGCTTTATTTGTAAACCATTATAAGCTGCAATAAAACAAGTTAACAACAACAATTGCATTATT
TTATGTTTTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAAACCTCTACAAA
TGTGGTAAATCGATAAGGATCCGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAGCT
CCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATG
CAACTCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCCTCGCTCACTGACTCGCTGCGCT
CGGTCGTTCCGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACA
GAATCAGGGGATAACGACAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCG
TAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAA
ATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCC
CCTGGAAGCTCCCTCGTGCCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGC
CTTCTCCTTCGGGAAGCGTGGCGCTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGG
TGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGC

GCCTTATCCGGTAACCTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGC
AGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGA
AGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAG
CCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAG
CGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATC
CTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGAACGAAAACTCACGTTAAGGGATTTTG
GTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTAA
ATCAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGG
CACCTATCTCAGCGATCTGTCTATTTCTGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAG
ATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCC
ACGCTCACCGGCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGCAGAA
GTGGTCTTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTA
AGTAGTTTCGCCAGTTAATAGTTTGCAGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGT
ACGCTCGTCTGTTGGTATGGCTTCATTCAGCTCCGGTTCCCAACGATCAAGCGCATCTACAT
GATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAGT
AAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCTAT
GCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGT
GTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGC
AGAAGTTTAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTT
ACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGACCCCACTGATCTTCAGCATCTT
TTACTTTACCCAGCGTTTCTGGGTGAGCAAAAAACAGGAAGGCAAAATGCCGCAAAAAAGGGA
ATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCTTTTCAATATTATTGAAGCAT
TTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACA
TAGGGGTTCCGCGCACATTTCCCGGAAAAGTGCCACCTGACGCGCCCTGTAGCGGCGCATTA
AGCGCGGCGGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCC
CGCTCCTTTTCGCTTTCTTCCCTTCTCCTTTCTCGCCACGTTTCGCGGCTTTCCCGCTCAAGCTC
TAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAAAA
CTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCCCTTT
GACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAACTGGAACAACACTCAACC
CTATCTCGGTCTATTTCTTTGATTTATAAGGGATTTTGCCGATTTTCGGCTATTGGTTAAAA
AATGAGCTGATTTAACAATAATTTAACGCGAATTTTAAACAAATATTAACGTTTACAATTTT
CCATTCGCCATTACAGGCTGCGCAACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTAT
TACGCCAGCCCAAGCTACCATGATAAGTAAGTAATATTAAGGTACGGGAGGTACTTGGAGCG
GCCGCTCTAGAAGTAGTGGATCCCCCGGCCGCAATAAAATATCTTTATTTTTCATTACATCTG
TGTGTTGGTTTTTTTGTGTGAATCGATAGTACTAACATACGCTCTCCATCAAAACAAAACGAA
ACAAAACAACTAGCAAAATAGGCTGTCCCCAGTGCAAGTGACAGGTGCCAGAACATTTCTCT
ATCGATAGGTACCGAGCTCTTACGCGTGCTAGCCCTCGAGCAGGATCTATACATTGAATCAA
TATTGGCAATTAGCCATATTAGTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCC
ATTGCATACGTTGTATCTATATCATAATATGTACATTTATATTGGCTCATGTCCAATATGAC
CGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTT
CATAGCCCATATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACC
GCCCCAACGACCCCCGCCCATTTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAG
GGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAACTGCCCACTTGGCAGTACAT
CAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTG
GCATTATGCCAGTACATGACCTTACGGGACTTTCTTACTTGGCAGTACATCTACGTATTAG
TCATCGCTATTACCATGGTGATGCGGTTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTT
GACTCACGGGGATTTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCA
AAATCAACGGGACTTTTCCAAAATGTGTAACAACTCCGCCCCATTGACGCAAAATGGGCGGTA
GGCGTGACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGG
AGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCCCCTC
GAAGCTCGACTCTAGGGGCTCGAGATCTGCGATCTAAGTAAGCTTGCATGCCTGCAGGTCCG
CCGCCACGACCGGTGCCGCCACCATCCCCTGACCCACGCCCTGACCCCTCACAGGAGACG
ACCTTCCATGACCGAGTACAAGCCCACGGTGCGCCTCGCCACCCGCGACGACGTCCCCCGG
CCGTACGCACCTCGCCGCCGCGTTTCGCCGACTACCCCGCCACGCGCCACACCGTCGACCCG

GACCGCCACATCGAGCGGGTCACCGAGCTGCAAGAACTCTTCCTCACGCGCGTCTGGGCTCGA
CATCGGCAAGGTGTGGGTCGCGGACGACGGCGCCGCGGTGGCGGTCTGGACCACGCCGGAGA
GCGTCGAAGCGGGGGCGGTGTTCCCGAGATCGGCCCGCGCATGGCCGAGTTGAGCGGTTCC
CGGCTGGCCGCGCAGCAACAGATGGAAGGCCTCCTGGCGCCGACCGGCCCAAGGAGCCCGC
GTGGTTCTTGGCCACCGTCGGCGTCTCGCCCGACCAAGGGCAAGGGTCTGGGCAGCGCCG
TCGTGCTCCCCGGAGTGGAGGCGGCCGAGCGCGCCGGGGTGCCCGCCTTCCTGGAGACCTCC
GCGCCCCGCAACCTCCCCTTCTACGAGCGGCTCGGCTTCACCGTCACCGCCGACGTGAGGT
GCCCCAAGGACCGCGCACCTGGTGCATGACCCGCAAGCCCGGTGCCTGACGCCCGCCCCACG
ACCCGCAGCGCCCGACCGAAAGGAGCGCACGACCCCATGGCTCCGACCGAAGCCGACCCGGG
CGGCCCCGCGACCCCGCACCCGCCCCCGAGGCCACCGACTCTAGAGTCGGGGCGGCCGGC
CGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAG
TGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAG
CTGCAATAAACAAGTTAACAACAACAATTGCATTCATTTTATGTTTCAGGTTTCAGGGGGAGG
TGTGGGAGGTTTTTTTAAAGCAAGTAAAACCTCTACAAATGTGGTAAAAATCGATAAGGATCAA
TTCGGCTTCAGGTACCGTCGACGATGTAGGTACGGTCTCGAAGCCGCGGTGCGGGTGCCAG
GGCGTGCCCTTGGGCTCCCCGGGCGCGTACTCCACCTCACCCATCTGGTCCATCATGATGAA
CGGGTCGAGGTGGCGGTAGTTGATCCCGGCGAACGCGCGGCGCACCGGGAAGCCCTCGCCCT
CGAAACCGCTGGGCGCGGTGGTCACGGTGAGCACGGGACGTGCGACGGCGTTCGGCGGGTGCG
GATACGCGGGGCAGCGTCAGCGGGTCTCGACGGTCACGGCGGGCATGTGACAGCCGAATT
GATCCGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAGCTCCTTCCGGTGGGCGCGGG
GCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAACTCGTAGGACAGGTG
CCGGCAGCGCTCTTCGGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTTCGGTTCGGCTGCGGC
GAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCA
GGAAAGAACATG

Fig. 15

pOM IFN-Ins-CMV-pur-attB (SEQ ID NO: 8)

GGCCGCCACCGCGGTGGAGCTCCAATTCGCCCTATAGTGAGTCGTATTACAATTCAGTGGCC
GTCGTTTTTACAACGTCGTGACTGGGAAAACCTGGCGTTACCCAACTTAATCGCCTTGCAGC
ACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAAC
AGTTGCGCAGCCTGAATGGCGAATGGGACGCGCCCTGTAGCGGCGCATTAAGCGCGGCGGGT
GTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGC
TTTCTTCCCTTCCCTTTCGCGCACGTTCCGCGGCTTTCCCGTCAAGCTCTAAATCGGGGGC
TCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCCAAAAAACTTGATTAGGGT
GATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTC
CACGTTCTTTAATAGTGGACTCTTGTTCCAACTGGAACAACACTCAACCCTATCTCGGTCT
ATTCTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATT
TAACAAAAATTTAACGCGAATTTTAACAAAAATATTAACGCTTACAATTTAGGTGGCACTTTT
CGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCC
GCTCATGAGACAATAACCCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGATGAGTA
TTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTTCGCGCATTTTTCCTTCTGTTTTTGCT
CACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTA
CATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTTC
CAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGACGCCGGG
CAAGAGCAACTCGGTGCGCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGT
CACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCA
TGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACC
GCTTTTTTGCACAACATGGGGGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAA
TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGC
GCAAATATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATG
GAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGC
TGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATG
GTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGA
AATAGACAGATCGCTGAGATAGGTGCCCTCACTGATTAAGCATTGGTAACCTGTGAGACCAAGT
TTACTCATATATACTTTAGATTGATTTAAAACCTTCATTTTTAATTTAAAAGGATCTAGGTGA
AGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCG
TCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTG
CTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTAC
CAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCTTCTA
GTGTAGCCGTAGTTAGGCCACCACTTCAAGAAGCTCTGTAGCACCGCCTACATACCTCGCTCT
GCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAAGTCGTGCTTACCAGGTTGGACT
CAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTCGTGCACACAG
CCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAG
CGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAG
GAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTGCGGTTT
CGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAA
AAAGCCAGCAACGCGGCCTTTTTACGTTTCTGCGCCTTTTGCTGGCCTTTTGCTCACATGT
TCTTTCCTGCGTTATCCCCTGATTCTGTGGATAAACCCTATTACCGCCTTTGAGTGAGCTGAT
ACCGCTCGCCGACGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCG
CCCAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTCATTAATGCAGCTGGCACGACA
GGTTTCCCGACTGGAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCAT
TAGGCACCCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGG
ATAACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCGAAATTAACCTC
ACTAAAGGGAACAAAAGCTGGGTACCGGGCCCCCCCCCTCGACTAGAGGGACAGCCCCCCCCCA
AAGCCCCCAGGGATGTAATTACGTCCCTCCCCCGCTAGGGGGCAGCAGCGAGCCGCCCGGGG
CTCCGCTCCGGTCCGGCGCTCCCCCGCATCCCCGAGCCGGCAGCGTGCGGGGACAGCCCGG
GCACGGGGAAGGTGGCACGGGATCGCTTTCTCTGAACGCTTCTCGCTGCTCTTTGAGCCTG
CAGACACCTGGGGGGATACGGGGAAAAAGCTTTAGGCTGAAAGAGAGATTTAGAATGACAGA
ATCATAGAACGGCCTGGGTGCAAAGGAGCACAGTGCTCATCCAGATCCAACCCCTGCTAT
GTGCAGGGTCATCAACCAGCAGCCCAGGCTGCCAGAGCCACATCCAGCCTGGCCTGAATG

CCTGCAGGGATGGGGCATCCACAGCCTCCTTGGGCAACCTGTTTCAGTGCCTCACCACCCTCT
GGGGGAAAACTGCCTCCTCATATCCAACCCAAACCTCCCCTGTCTCAGTGTAAGCCATTC
CCCCTTGTCTATCAAGGGGGAGTTTGCTGTGACATTGTTGGTCTGGGGTGACACATGTTTG
CCAATTCAGTGCATCACGGAGAGGCAGATCTTGGGGATAAGGAAGTGCAGGACAGCATGGAC
GTGGGACATGCAGGTGTTGAGGGCTCTGGGACACTCTCCAAGTCACAGCGTTTCAGAACAGCC
TTAAGGATAAGAAGATAGGATAGAAGGACAAAGAGCAAGTTAAAACCCAGCATGGAGAGGAG
CACAAAAAGGCCACAGACACTGCTGGTCCCTGTGTCTGAGCCTGCATGTTTGATGGTGTCTG
GATGCAAGCAGAAGGGGTGGAAGAGCTTGCCCTGGAGAGATACAGCTGGGTTCAGTAGGACTGG
GACAGGCAGCTGGAGAATTGCCATGTAGATGTTTCATACAATCGTCAAATCATGAAGGCTGGA
AAAGCCCTCCAAGATCCCCAAGACCAACCCCAACCCACCCACCGTGCCCACTGGCCATGTCC
CTCAGTGCCACATCCCCACAGTTCTTCATCACCTCCAGGGACGGTGACCCCCCACCTCCGT
GGGCAGCTGTGCCACTGCAGCACCGCTCTTTGGAGAAGGTAAATCTTGCTAAATCCAGCCCCG
ACCTTCCCCTGGCACAACGTAAGGCCATTATCTCTCATCCAACCTCCAGGACGGAGTCAGTGA
GGATTGGGGCTCTAGTCGAGGTTCGACGGTATCGATAAGCTTGATTAGGCAGAGCAATAGGACT
CTCAACCTCGTGAGTATGGCAGCATGTTAACTCTGCCTGGAGTCCAGCGTGGGAAACAATC
TGCCTTGCACATGAGTCTTCGTGGGCCAATATTCCCCAACGGTTTTCTTTCAGCTTGTCTTG
TCTCCTAAGCTCTCAAAACACCTTTTTTGGTGAATAAACTCACTTGGCAACGTTTATCTGTCT
TACCTTAGTGTACGTTTCATCCCTATTCCCCTTTCTCCTCCTCCGTGTGGTACACAGTGGT
GCACATGGTTCTTCTGTGTGATGTTCTGTCTGCAGCCCAATGTGGGTAAAGTTCTTCTCTGC
CACGTGTCTGTGTTGTTTTCACTTCAAAAAGGGCCCTGGGCTCCCCTTGGAGCTGTGTCTTA
TTTTCTTAATCATCACAGTCACGCTGGCAGGATTAGTCCCTCCTAAACCTTAGAATGACCTG
AACGTGTGCTCCCTCTTGTAGTCAGTGCAGGGAGACGTTTGCCTCAAGATCAGGGTCCATC
TCACCCACAGGGCCATTCCCAAGATGAGGTGGATGTTTACTCTCACAAAAAGTTTTCTTAT
GTTTGGCTAGAAAGGAGAACTCACTGCCTACCTGTGAATTCCCCTAGTCCTGGTTCTGCTGC
CACTGCTGCCTGTGCAGCCTGTCCCATGGAGGGGGCAGCAACTGCTGTACAAAAGGTGATCC
CACCCTGTCTCCACTGAAATGACCTCAGTGCCACGTGTTGTATAGGGTATAAAGTACGGGAG
GGGGATGCCCGGCTCCCTTCAGGGTTGCAGAGCAGAAGTGTCTGTGTATAGAGTGTGTCTTA
ATCTATTAATGTAACAGAACAACCTTCAGTCCTAGTGTTTTGTGGGCTGGAATTGCCCATGTG
GTAGGGACAGGCCTGCTAAATCACTGCAATCGCCTATGTTCTGAAGGTATTTGGGAAAGAAA
GGGATTTGGGGGATTGCCTGTGATTGGCTTTAATTGAATGGCAAATCACAGGAAAGCAGTTC
TGCTCAACAGTTGGTTGTTTCAGCCAATTCTTGCAGCCAAAGAGCCGGGTGCCCAGCGATAT
AATAGTTGTCACTTGTGTCTGTATGGATGACAGGGAGGTAGGGTGACCTGAGGACCACCTC
CAGCTTCTGTAGCTAGGTACAGTCACCACTCCAGCTCCACACGAGTCCCATCGTGGTTT
ACCAAAGAAACACAATTATTTGGACCAGTTTGGAAGTCAACCGCTGAATTGTGAGGCTAGA
TTAATAGAGCTGAAGAGCAAATGTTCCCAACTTGGAGATACTAGTTGGTATTAGTATCAGAG
GAACAGGGCCATAGCACCTCCATGCTATTAGATTCCGGCTGGCATGTACTTTTCAAGATGAT
TTGTAACTAACAATGGCTTATTGTGCTTGTCTTAAGTCTGTGTCTAATGTAAATGTTCCCTT
TGGTTTATATAACCTTCTTGCCATTTGCTCTTCAGGTGTTCTTGCAGAACACTGGCTGCTTT
AATCTAGTTTAACTGTTGCTTGATTATTCTTAGGGATAAGATCTGAATAAACTTTTTGTGGC
TTTGCCAGACTTTAGCTTGGGCTTAGCTCCACATTAGCTTTTGCTGCCTTTTCTGTGAAGC
TATCAAGATCCTACTCAATGACATTAGCTGGGTGCAGGTGTACCAAATCCTGCTCTGTGGAA
CACATTGTCTGATGATACCGAAGGCAAACGTGAACTCAAAGAGGCACAGAGTTAAGAAGAAG
TCTGTGCAATTCAGAGGAAAAGCCAAAGTGGCCATTAGACACACTTCCATGCAGCATTTGC
CAGTAGGTTTCATATAAACTACAAAATGGAATAAACCCTACAAATGGGAAAAGCCTGATA
CTAGAATTTAAATATTACCCAGGCTCAAGGGGTGTTTCATGGAGTAATATCACTCTATAAA
AGTAGGGCAGCCAATTATTCACAGACAAAGCTTTTTTTTTTCTGTGCTGCAGTGTGTTTTT
CGGCTGATCCAGGGTTACTTATTGTGGGTCTGAGAGCTGAATGATTTCTCCTTGTGTCTATGT
TGGTGAAGGAGATATGGCCAGGGGGAGATGAGCATGTTCAAGAGGAAACGTTGCATTTTGGT
GGCTTGGGAGAAAGGTAGAACGATATCAGGTCCATAGTGTCACTAAGAGATCTGAAGGATGG
TTTTACAGAACAGTTGACTTGGCTGGGTGCAGGCTTGGCTGTAAATGGATGGAAGGATGGAC
AGATGGGTGGACAGAGATTTCTGTGCAGGAGATCATCTCCTGAGCTCGGTGCTTGACAGACT
GCAGATCCATCCCATAACCTTCTCCAGCATGAGAGCGCGGGGAGCTTTGGTACTGTTTCAGTC
TGCTGCTTGTGCTTCTGGGTGCACAGTGGTGATTTTCTTACTCACACAGGGCAAAAACCT
GAGCAGCTTCAAAGTGAACAGGTTGCTCTCATAGGCCATTACAGTTGTCAAGATGAGGTTTTT

GGTTTCTTGT TTTGTAAGGTGGGAAGAAGCACTGAAGGATCAGTTGCGAGGGCAGGGGTTTA
GCACTGTTCAAGAGAAGTCTTATTTTAACTCCTCTCATGAACAAAAAGAGATGCAGGTGCAGA
TTCTGGCAAGCATGCAGTGAAGGAGAAAGCCCTGAATTTCTGATATATGTGCAATGTTGGGC
ACCTAACATTCCCCGCTGAAGCACAGCAGCTCCAGCTCCATGCAGTACTCACAGCTGGTGCA
GCCCTCGGCTCCAGGGTCTGAGCAGTGCTGGGACTCACGAGGTTCATGTCTTTCACACTGA
TAATGGTCCAATTTCTGGAATGGGTGCCCATCCTTGGAGGTCCCCAAGGCCAGGCTGGCTGC
GTCTCCGAGCAGCCCCGATCTGGTGGTGAGTAGCCAGCCCATGGCAGGAGTTAGAGCCTGATG
GTCTTTAAGGTCCCTTCCAACCTAAGCCATCCTACGATTCTAGGAATCATGACTTGTGAGTG
TGTATTGCAGAGGCAATATTTTAAAGTTATAAATGTTTCTCCCCTTCCCTTGTGTCAAAG
TTATCTTGATCGCCTTATCAATGCTTTTGGAGTCTCCAGTCATTTTTCTTACAMCAAAAAGA
GGAGGAAGAATGAAGAGAATCATTTAATTTCTTGATTGAATAGTAGGATTCAGAAAGCTGTA
CGTAATGCCGTCTCTTTGTATCGAGCTGTAAGGTTTCTCATCATTTATCAGCGTGGTACATA
TCAGCACTTTTCCATCTGATGTGGAAAAAAATCCTTATCATCTACAGTCTCTGTACCTAA
ACATCGCTCAGACTCTTTACCAAAAAAGCTATAGGTTTAAAACTACATCTGCTGATAATTT
GCCTTGT TTTTAGCTCTTCTTCCATATGCTGCGTTTGTGAGAGGTGCGTGGATGGGCCTAAAC
TCTCAGCTGCTGAGCTTGATGGGTGCTTAAGATTGAAGCACTCACTGCTGAAACTGTTTTCA
TTTCACAGGAATGTTTTAGTGGCATTTGTTTTATAACTACATATTCCTCAGATAAATGAAAT
CCAGAAATAATTATGCAAACTCACTGCATCCGTTGCACAGGTCTTTATCTGCTAGCAAAGGA
AATAATTTGGGGATGGCAAAAACATTCCTTCAGACATCTATATTTAAAGGAATATAATCCTG
GTACCCACCCACTTCATCCCTCATTTATGTTACACTCAGAGATACTCATTTCTTGTGTGTA
TCATTTGATAGCGTTTTCTTTGGTTCCTTGGCACGCTCTGGGCTATGGCTGCACGCTCTGCA
CTGATCAGCAAGTAGATGCGAGGGAAGCAGCAGTAGAGAGGGGCTGCCCTCAGCTGGCACC
GCCGCTCAGCCTAGGAGGGGACCTTGCCCTTACCACAGCTGAGGTGCAGCCCTACAAGCTTA
CACGTGCTGCGAGCAGGTGAGCAAAGGGAGTCTTCATGGTGTGTTTTCTTGCTGCCCGGAAGC
AAAACTTTACTTTTCATTTCATTTCCCCTTGAAGAATGAGGAATGTTTGGAAACGGACTGCTTTA
CGTTCAATTTCTCTCTTCCCTTTAAGGCTCAGCCAGGGGCCATTGCTGAGGACGGCATCGGG
GCCCCCTGGACCAAATCTGTGGCACAGATGGTTTTCACTTACATCAGTGGATGTGGGATCTGC
GCCTGTAATGTGTCTTCTGAAGGAAGGAACGTGCCCTTCCAAGTGCCAGCCCCACAGCCCC
AGCCCCCTCCCTGTGCTGCTCCAATTCATCTCCTCTTCCCTTCCCTTCTGCTGTTTGTGTC
TCGGGTAGAAATCATGAAGATTTAGAAGAGAAAACAAATAACTGGAGTGGAAACCCAGGTG
ATGCAGTTTCATTTCAGCTGTCATAGGTTTGTGCTTGTATAGGTCTGTATCAGAGATGCTARC
ACCACTTTGCTGTCGGTGCTTAACTCGGGTGAACCTCCTTCACTCGCATCATTTGCGGGCC
TTATTTACATCCCCAGCATCCATCACCCCTCTGGGAAAATGGGCGCACTGGATCTCTAATGGA
AGACTTTCCCTCTTTTCAGAGCCTGTGGGATGTGCAGTGACAAGAAACGTGGAGGGGCTGAGC
AGCAGCACTGCCCCCAGGGAGCAGGAGCGGATGCCATCGGTGGCAGCATCCCAAATGATGTC
AGCGGATGCTGAGCAGGCAGCGGACGACGACGAGAGCGATGCGTACACCTTCTGTTGACA
TGGTATTTTGGCAGCGATTTAACACTCGCTTCCTAGTCTCCTGCTATTCTCCACAGGCTGCATTC
AAATGAACGAAGGGAAGGGAGGCAAAAAGATGCAAAATCCGAGACAAGCAGCAGAAATATTT
CTTCGCTACGGAAGCGTGCGCAAACAACCTTCTCCAACAGCACCAGAAGAGCACAGCGTAAC
CTTTTTCAAGACCAGAAAAGGAAATTCACAAAGCCTCTGTGGATACCAGCGGTTTCAGCTCT
CCTGATAGCAGATTTCTTGTGTCAGGTTGCGAATGGGGTATGGTGCCAGGAGGTGCAGGGACCA
TATGATCATATACAGCACAGCAGTCATTGTGCATGTATTAATATATATTGAGTAGCAGTGTT
ACTTTGCCAAAGCAATAGTTTCAGAGATGAGTCCTGCTGCATACCTCTATCTTAAACTA
TATAAATAGTAAAACCTTCTCAGTTTCAGCCACGTGCTCCTCTCTGTCAGCACCAATGGTGCT
TCGCCTGCACCCAGCTGCAAGGAATCAGCCCGTGATCTCATTAACACTCAGCTCTGCAGGAT
AAATTAGATTGTTCCACTCTCTTTTGTGTTAATTACGACGGAACAATTGTTTCAGTGCTGAT
GGTCCTAATTGTGTCAGCTACAGAAAACGTCTCCATGCAGTTCCTTCTGCGCCAGCAAACTGTC
CAGGCTATAGCACCGTGATGCATGCTACCTCTCACTCCATCCTTCTTCTTCTTCCCACAGG
GAGAGCTGTGTGTTTTCACTCTCAGCCACTCTGAACAATAACCAAACTGCTACGCATGCTC
CCTCGGAAAGAGAAATCCCCTTGTGCTTTTATTTATTTACAGGATCCTTCTTAAAGCAGACC
ATCATTCAGTGCAAAACCCAGAGCTTCATGCCTCTCCTTCCACAACCGAAAACAGCCGGCTTC
ATTTGTCTTTTTTAAATGCTGTTTTCCAGGTGAATTTTGGCCAGCGTGTTGGCTGAGATCCA
GGAGCACGTGTGCTGCTCTCATTTGCTCCTGTTCTGCATTGCCTCTTTCTGGGGTTT
CCAAGAGGGGGGGAGACTTTGCGCGGGGATGAGATAATGCCCTTTTCTTAGGGTGGCTGCT

GGGCAGCAGAGTGGGCTCTGGGTCACTGTGGCACCAATGGGAGGCACCAAGTGGGGGTGTGTTT
TGTGCAGGGGGGAAGCATTACAGAATGGGGCTGATCCTGAAGCTTGCAGTCCAAGGCTTTG
TCTGTGTACCCAGTGAAATCCTTCCTCTGTTACATAAAGCCCAGATAGGACTCAGAAATGTA
GTCATTCCAGCCCCCTCTTCCTCAGATCTGGAGCAGCACTTGTTCAGCCAGTCTCCCC
AAAATGCACAGACCTCGCCGAGTGGAGGGAGATGTAAACAGCGAAGGTTAATTACCTCCTTG
TCAAAAACACTTTTGTGGTCCATAGATGTTTCTGTCAATCTTACAAAACAGAACCGAGAGGCA
GCGAGCACTGAAGAGCGTGTTCCTATGCTGAGTTAATGAGACTTGGCAGCTCGCTGTGCAGA
GATGATCCCTGTGCTTCATGGGAGGCTGTAACCTGTCTCCCCATCGCCTTCACACCGCAGTG
CTGGCCTTGACACAAACCTCCATAAAGCTGTAGGATGCAGCTGCCAGGGATCAAGAGAACT
TTTCCTAAGGCTCTTAGGACTCATCTTTGCCGCTCAGTAGCGTGCAGCAATTACTCATCCCA
ACTATACTGAATGGGTTCCTGCCAGCTCTGCTTGTTCATTAAGCAATTTCTTCATTTTGC
CTCTAAGTTTCTCTCAGCAGCACCGCTCTGGGTGACCTGAGTGGCCACCTGGAACCCGAGGG
GCACAGCCACCACCTCCCTGTTGCTGCTGCTCCAGGGACTCATGTGCTGCTGGATGGGGGGA
AGCATGAAGTTCCTCACCCAGACACCTGGGTTCGAATGGCTGCAGCGTGTCTTCTTGGTAT
GCAGATTGTTTCCAGCCATTACTTGTAGAAATGTGCTGTGGAAGCCCTTTGTATCTCTTTCT
GTGGCCCTTGACACAAAGCTGTGGGAAAGCTGTAGGCTGCTTTCTTGGGTCTGGAGGAAT
TGATATGTTCCCTTCTTTAACAAAAATTATCCTTAGGAGAGAGCACTGTGCAAGCATTGTGCAC
ATAAAACAATTCAGGTTGAAAGGGCTCTCTGGAGGTTTCCAGCCTGACTACTGCTCGAAGCA
AGGCCAGGTTCAAAGATGGCTCAGGATGCTGTGTGCCTTCCTGATTATCTGTGCCACCAATG
GAGGAGATTACAGCCACTCTGCTTCCCGTGCCACTCATGGAGAGGAATATTCCTTATATTT
CAGATAGAATGTTATCCTTTAGCTCAGCCTTCCTATAACCCCATGAGGGAGCTGCAGATCC
CCATACTCTCCCCTTCTCTGGGGTGAAGGCCGTGTCCCCAGCCCCCTTCCACCTGTGC
CCTAAGCAGCCCGCTGGCCTCTGCTGGATGTGTGCCTATATGTCAATGCCTGTCTTGCAGT
CCAGCCTGGGACATTTAATTCATCACCAGGGTAATGTGGAACGTGTGTCATCTTCCCCTGCAG
GGTACAAAGTTCTGCACGGGGTCTTTTCGGTTCAGGAAAACCTTCACTGGTGCTACCTGAAT
CAAGCTCTATTTAATAAGTTCATAAGCACATGGATGTGTTTTCTTAGAGATACGTTTTAATG
GTATCAGTGATTTTTATTTGCTTTGTTGCTTACTTCAAACAGTGCCTTTGGGCAGGAGGTGA
GGGACGGGTCTGCCGTGGCTCTGCAGTGATTTCTCAGGCGTGTGGCTCAGGTCAGATAGT
GGTCACTCTGTGGCCAGAAGAAGGACAAAGATGGAATTCAGATTGAGTACGTTAAGCAG
GCATCTTGGATGATTTGAGGCAGTTTTCATGAAAGAGCTACGACCACTTATTTGTTTTCG
CCTTTTACAACAGAAGTTTTTCATCAAAATAACGTGGCAAAGCCCAGGAATGTTTGGGAAAAG
TGTAAGTTAAATGTTTTGTAATTCATTTGTGCGAGTGCTACCAGCTAAGAAAAAGTCTACC
TTTGGTATGGTAGTCCTGCAGAGAATACAACATCAATATTAGTTTGGAAAAAACACCACCA
CCACCAGAACTGTAATGGAAAATGTAAACCAAGAAATTCCTTGGGTAAAGAGAGAAAGGATG
TCGTATACTGGCCAAGTCCTGCCAGCTGTGAGCCTGCTGACCCTCTGCAGTTCAGGACCAT
GAAACGTGGCACTGTAAGACGTGTCCCCCTGCTTTGCTTGGCCACAGATCTCTGCCCTGTG
CTGACTCCTGCACACAAGAGCATTTCCCTGTAGCCAAACAGCGATTAGCCATAAGCTGCACC
TGACTTTGAGGATTAAGAGTTTGAATTAAGTGGATTGCAGCAGGAGATCAGTGGCAGGGTT
GCAGATGAAATCCTTTTCTAGGGGTAGCTAAGGGCTGAGCAACCTGTCTACAGCACAAGCC
AAACCAGCCAAGGGTTTTCTCTGTGCTGTTACAGAGGCAGGGCCAGCTGGAGCTGGAGGAGG
TTGTGCTGGGACCCTTCTCCCTGTGCTGAGAATGGAGTGATTTCTGGGTGCTGTTCTGTGG
CTTGCACTGAGCAGCTCAAGGGAGATCGGTGCTCCTCATGCAGTGCCAAAACCTCGTGTGTA
TGCAGAAAGATGGATGTGCACCTCCCTCCTGCTAATGCAGCCGTGAGCTTATGAAGGCAATG
AGCCCTCAGTGCAGCAGGAGCTGTAGTGCACCTCCTGTAGGTGCTAGGGAAAAATCTCTGGTTC
CCAGGGATGCATTCATAAGGGCAATATATCTTGAGGCTGCGCCAAATCTTTCTGAAATATTC
ATGCGTGTTCCCTTAATTTATAGAAACAAACACAGCAGAATAATTATTCGAATGCCTCCCCT
CGAAGGAAACCCATATTTCCATGTAGAAATGTAACCTATATACACACAGCCATGCTGCATCC
TTCAGAACGTGCCAGTGCTCATCTCCCATGGCAAAATACTACAGGTATTTCTCACTATGTTGG
ACCTGTGAAAGGAACCATGGTAAGAACTTCGGTTAAAGGTATGGCTGCAAAACTACTCATA
CCAAAACAGCAGAGCTCCAGACCTCCTCTTAGGAAAGAGCCACTTGGAGAGGGATGCTGTA
AGGCTGGAGGTGAGAGACAGAGCCTGTCCAGTTTCTCTGTCTCTATTTTCTGAAACGTTTG
CAGGAGGAAAGGACAACGTACTTTTCAAGCATAGCTGGTGGCCTCACGTAAATAAGTTCCCC
GAACTTCTGTGTCAATTTGTTCTTAAGATGCTTTGGCAGAACACTTTGAGTCAATTCGCTTAA
CTGTGACTAGGTCTGTAAATAAGTGCTCCCTGCTGATAAGGTTCAAGTGACATTTTTTAGTGG

AATTGACAGCATTACCTTGCCTTCAAGTCTTCTACCAAGCTCTTCTATACTTAAGCAGTG
 AAACCGCCAAGAAACCCCTTCCCTTTTATCAAGCTAGTGCTAAATACCATTAACTTCATAGGTT
 AGATACGGTGCTGCCAGCTTCACCTGGCAGTGGTTGGTCAGTTCTGCTGGTGACAAAGCCTC
 CCTGGCCTGTGCTTTTACCTAGAGGTGAATATCCAAGAATGCAGAAGTGCATGGAAAGCAGA
 GCTGACGGCAGCATGGTGCTGAGCCTTAGCTGCTTCTGCTGGGAGATGTGGATGCAGAGAC
 GAATGAAGGACCTGTCCCTTACTCCCTCAGCATTTCTGTGCTATTAGGGTTCTACAGAGT
 CCTTAAGAGGTTTTTTTTTTTTTTTGGTCCAAAAGTCTGTTTGTGTTGGTTTTGACCAC TGAGA
 GCATGTGACACTTGTCTCAAGCTATTAAACCAAGTGTCCAGCCAAAATCAATTGCCTGGGAGA
 CGCAGACCATTACCTGGAGGTGAGGACCTCAATAAATATTACCAGCCTCATTGTGCCGCTGA
 CAGATTGAGCTGGCTGCTCCGTGTTCCAGTCCAACAGTTCGGACGCCACGTTTGTATATATT
 TGCAGGCAGCCTCGGGGGGACCATCTCAGGAGCAGAGCACCAGCCGCGCTGCAGAGCCCGG
 GCAGTACCTCACCATTGGCTTTGACCTTTGCCTTACTGGTGGCTCTCCTGGTGCTGAGCTGCA
 AGAGCAGCTGCTCTGTGGGCTGCGATCTGCCTCAGACCCACAGCCTGGGCGAGCAGGAGGACC
 CTGATGCTGCTGGCTCAGATGAGGAGAATCAGCCTGTTTAGCTGCCTGAAGGATAGGCACGA
 TTTTGGCTTTTCTCAAGAGGAGTTTGGCAACCAGTTTCAGAAGGCTGAGACCATCCCTGTGC
 TGCACGAGATGATCCAGCAGATCTTTAACCCTGTTTAGCACCAAGGATAGCAGCGCTGCTTGG
 GATGAGACCTGCTGGATAAGTTTACACCGAGCTGTACCAGCAGCTGAACGATCTGGAGGC
 TTGCGTGATCCAGGGCGTGGGCGTGACCGGAGACCCCTCTGATGAAGGAGGATAGCATCCTGG
 CTGTGAGGAAGTACTTTCAGAGGATACCCCTGTACCTGAAGGAGAAGATACAGCCCCTGC
 GCTTGGGAAGTCGTGAGGGCTGAGATCATGAGGAGCTTTAGCCTGAGCACCAACCTGCAAGA
 GAGCTTGAGGTCTAAGGAGTAAAAAGTCTAGAGTCGGGGCGGCCGGCCGCTTCGAGCAGACA
 TGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGAAAAAATGCTTT
 ATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAAGT
 TAACAACAACAATTGCATTCAATTTTATGTTTTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTT
 AAAGCAAGTAAACCTCTACAAATGTGTGTTAAATCGATAACCTGCACCTCGACTGAGCAGCGG
 CACTAACATACGCTCTCCATCAAAACAAAACGAAACAAAACAAACTAGCAAAATAGGCTGTC
 CCCAGTGCAAGTGCAGGTGCCAGAACATTTCTCTATCGATAGGTACCGAGCTCTTACGCGTG
 CTAGCCCTCGAGCAGGATCTATACATTGAATCAATATTGGCAATTAGCCATATTAGTCATTG
 GTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATATCATAAT
 ATGTACATTTATATTGGCTCATGTCCAATATGACCGCCATGTTGACATTGATTATTGACTAG
 TTATTAATAGTAATCAATTACGGGGTCATTAGTTTCATAGCCCATATATGGAGTTCGCGCTTA
 CATAACTTACGGTAAATGGCCCGCTGGCTGACCGCCCAACGACCCCGCCCATGACGCTCA
 ATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGA
 GTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTCCGCCCC
 CTATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTACGG
 GACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTT
 TTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGA CTACGCGGGATTTCCAAGTCTCCACC
 CCATTGACGTCAATGGGAGTTTGTTTTGGCACAAAATCAACGGGACTTTCCAAAATGTGCT
 AACAACTCCGCCCCATTGACGCAAAATGGGCGGTAGGCGTGCATCGGTGGGAGGTCTATATAAG
 CAGAGCTCGTTTTAGTGAACCGTGCAGATCGCCTGGAGACGCGCATCCACGCTGTTTTGACCTCC
 ATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGAAGCTCGACTCTAGGGGCTCGAGATCT
 GCGATCTAAGTAAGCTTGCATGCCTGCAGGTGCGCCGCCACGACCGGTGCCGCCACCATCCC
 CTGACCCACGCCCCCTGACCCCTCACAAGGAGACGACCTTCCATGACCGAGTACAAGCCCACG
 GTGCGCCTCGCCACCCGCGACGACGCTCCCCCGGGCCGTACGCAACCCCTCGCCGCCGCGTTGCG
 CGACTACCCCGCCACGCGGCACACCGTCGACCCGCGGACCGCCACATCGAGCGGGTCAACGAGC
 TGAAGAAGTCTTCTCACGCGCGCTCGGGCTCGACATCGGCAAGGTGTGGTTCGCGGACGAC
 GGCGCCGCGGTGGCGGTCTGGACCACGCCGAGAGCGTGAAGCGGGGGCGGTGTTGCGCCGA
 GATCGGCCCGCGCATGGCCGAGTTGAGCGGTTCCCGGCTGGCCGCGCAGCAACAGATGGAAG
 GCCTCCTGGCGCGCACCGGCCCAAGGAGCCCGCGTGGTTCTTGCCACCGTCGGCGTCTCG
 CCGGACCCACAGGGCAAGGGTCTGGGCAGCGCCGTCGTGCTCCCCGGAGTGGAGGCGGCCGA
 GCGCGCCGGGGTCCCCGCTTCTTGAGACCTCCGCGCCCGGCAACCTCCCCCTTCTACGAGC
 GGCTCGGCTTACCCTCACCGCCAGCTCGAGTGCCCGCAAGGACCGCGACCTGGTGATG
 ACCCGCAAGCCCGGTGCTGACGCGCGCCCCACGACCCGACGCGCCGACCGAAGAGCGC
 ACGACCCCATGGCTCCGACCGAAGCCGACCCGGGCGGGCCCCGCGACCCCGCACCCGCCCC

GAGGCCCACCGACTCTAGAGTCGGGGCGGCCGCGCTTCGAGCAGACATGATAAGATACAT
TGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTT
GTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAGTTAACAACAACAAT
TGCATTCATTTTATGTTTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAAA
CCTCTACAAATGTGGTAAAATCGATAAGGATCAATTCGGCTTCAGGTACCGTCGACGATGTA
GGTCACGGTCTCGAAGCCGCGGTGCGGGTGCCAGGGCGTGCCCTTGGGCTCCCCGGGCGCGT
ACTCCACCTCACCCATCTGGTCCATCATGATGAACGGGTCGAGGTGGCGGTAGTTGATCCCG
GCGAACGCGCGGCGCACCCGGAAGCCCTCGCCCTCGAAACCGCTGGGCGCGGTGGTCACGGT
GAGCACGGGACGTGCGACGGCGTCGGCGGGTGCGGATACGCGGGGCAGCGTCAGCGGGTTCT
CGACGGTCACGGCGGGCATGTGACAGCCGAATTGATCCGTGACCGATGCCCTTGAGAGCC
TTCAACCCAGTCAGCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGAC
TGTCTTCTTTATCATGCAACTCGTAGGACAGGTGCCCGCAGC

Fig. 16

prSV-C31int (SEQ ID NO: 9)

CTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTTCGCTATTGGGCGCTCTTCC
GCTTCCCTCGCTCACTGACTCGCTGCGCTCGGTTCGGCTGCGGCGAGCGGTATCAGCT
CACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAATCATG
TGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTC
CATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGA
AACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCCTCT
CCTGTTCCGACCTGCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCGGAAGCGTG
GCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCTCCAAG
CTGGGCTGTGTGCACGAACCCCCCGTTACGCGGACCGCTGCGCTTATCCGGTAACTAT
CGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAAC
AGGATTAGCAGAGCGAGGTATGTAGGCGGTGTACAGAGTTCTTGAAGTGGTGGCCTAAC
TACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTC
GGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCCGCTGGTAGCGGTGGTTTT
TTTGTTCGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATC
TTTTCTACGGGGTCTGACGCTCAGTGGACGAAAACTCACGTTAAGGGATTTTGGTCATG
AGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAAATGAAGTTTTAAATCA
ATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCA
CCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAG
ATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAC
CCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGC
AGAAGTGGTCTGCAACTTTATCCGCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCT
AGAGTAAGTAGTTCCGCGAGTTAATAGTTTGGCGAACGTTGTTGCCATTGCTACAGGCATC
GTGGTGTACGCTCGTCTGTTGGTATGGTTCATTTCAGCTCCGCTTCCCAACGATCAAGG
CGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATC
GTTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAAT
TCTCTTACTGTCTATGCCATCCGTAAGATGCTTTTTCTGTGACTGGTGAGTACTCAACCAAG
TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCGGCGTCAATACGGGAT
AATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGAAAACGTTCTTCGGGG
CGAAAACCTCAAGGATCTTACCGCTGTTGAGATCCAGTTTCGATGTAAACCCACTCGTGCA
CCCAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGA
AGGCAAAATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTC
TTCCTTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATA
TTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTG
CCACCTGACGTCGACGGATCGGGAGATCTCCCGATCCCCTATGGTGCAGTCTCAGTACAA
TCTGCTCTGATGCCGATAGTTAAGCCAGTATCTGCTCCCTGCTTGTGTGTTGGAGTTCG
CTGAGTAGTGCGCGAGCAAAATTTAAGCTACAACAAGGCAAGGCTTGACCGACAATTGCA
TGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCGATGTACGGGCCAGATATA
CGCGTGCTAGGGGTCTAGGATCGATTCTAGGAATTCTCTAGCCGCGGTCTAGGGATCCCG
GCGCGTATGGTGCATCTCAGTACAATCTGCTCTGATGCCGATAGTTAAGCCAGTATCT
GCTCCCTGCTTGTGTGTTGGAGGTGCTGAGTAGTGCGCGAGCAAAATTTAAGCTACAAC
AAGGCAAGGCTTGACCGACAATTGCATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCT
GCTTCGCGATGTACGGGCCAGATATACGCGTATCTGAGGGGACTAGGGTGTGTTTAGGCG
AAAAGCGGGGCTTCGGTTGTACGCGGTTAGGAGTCCCCTCAGGATATAGTAGTTTCGCTT
TTGCATAGGGAGGGGGAAATGTAGTCTTATGCAATACACTTGTAGTCTTGCAACATGGTA
ACGATGAGTTAGCAACATGCCTTACAAGGAGAGAAAAAGCACCGTGATGCCGATTGGTG
GAAGTAAGGTGGTACGATCGTGCCTTATTAGGAAGGCAACAGACAGGTCTGACATGGATT
GGACGAACCACTGAATTCGCGATTCAGAGATAATTGTATTTAAGTGCCTAGCTCGATAC
AATAAACGCCATTTGACCATTACACCATTTGGTGTGACCTCCAAGCTTGATGCCTGCA
GGTACCGGTCCGGAATTCGCGGGTTCGACGAGCTCACTAGTCGTAGGGTTCGCGGACATGAC
ACAAGGGGTTGTGACCGGGGTGGACACGTACGCGGGTGCTTACGACCGTCAGTCGCGCGA
GCGCGAGAATTTCGAGCGCAGCAAGCCAGCGACACAGCGTAGCGCCAACGAAGACAAGGC
GGCCGACCTTCAGCGCGAAGTCGAGCGCGACGGGGGCCGTTTCAGGTTGCTCGGGCATT
CAGCGAAGCGCCGGGCACGTTCGGCTTCGGGACGGCGGAGCGCCCGAGTTTCAACGCAT

CCTGAACGAATGCCGCGCCGGGCGGGCTCAACATGATCATTGTCTATGACGTGTCGCGCTT
CTCGCGCCTGAAGGTCATGGACGCGATTCCGATTGTCTCGGAATTGCTCGCCCTGGGCGT
GACGATTGTTTTCCACTCAGGAAGGCGTCTTCCGGCAGGGAAACGTCATGGACCTGATTCA
CCTGATTATGCGGCTCGACGCGTCGCACAAAGAATCTTCGCTGAAGTCGGCGAAGATTCT
CGACACGAAGAACCTTCAGCGCGAATTGGGCGGGTACGTCGGCGGGAAGGCGCCTTACGG
CTTCGAGCTTGTTTCGGAGACGAAGGAGATCACGCGCAACGGCCGAATGGTCAATGTCGT
CATCAACAAGCTTGCGCACTCGACCACTCCCCTTACCGGACCCTTCGAGTTCGAGCCCCGA
CGTAATCCGGTGGTGGTGGCGTGAGATCAAGACGCACAAACACCTTCCCTTCAAGCCGGG
CAGTCAAGCCGCCATTCACCCGGGCAGCATCACGGGGCTTTGTAAGCGCATGGACGCTGA
CGCCGTGCCGACCCGGGGCGAGACGATTGGGAAGAAGACCGCTTCAAGCGCTGGGACCC
GGCAACCGTTATGCGAATCCTTCGGGACCCGCGTATTGCGGGCTTCGCCGCTGAGGTGAT
CTACAAGAAGAAGCCGGACGGCACGCCGACCACGAAGATTGAGGGTTACCGCATTACAGCG
CGACCCGATCACGCTCCGGCCGGTCGAGCTTGATTGCGGACCGATCATCGAGCCCCGCTGA
GTGGTATGAGCTTCAGGCGTGTTGGACGGCGCGGGGCGCGGCAAGGGGCTTTCCCGGGG
GAAGCCATTCTGTCCGCCATGGACAAGCTGTACTGCGAGTGTGGCGCCGTCATGACTTC
GAAGCGCGGGGAAGAATCGATCAAGGACTCTTACCGCTGCCGTCGCCGGAAGGTGGTCGA
CCCGTCCGCACCTGGGCAGCACGAAGGCACGTGCAACGTCAGCATGGCGGCACTCGACAA
GTTTCGTTGCGGAACGCATCTTCAACAAGATCAGGCACGCCGAAGGCGACGAAGAGACGTT
GGCGCTTCTGTGGGAAGCCGCCGACGCTTCGGCAAGCTCACTGAGGCGCCTGAGAAGAG
CGGCGAACGGGCGAACCTTGTTGCGGAGCGCGCCGACGCCCTGAACGCCCTTGAAGAGCT
GTACGAAGACCGCGCGGCAGGCGGTACGACGGACCCGTTGGCAGGAAGCACTTCCGGAA
GCAACAGGCAGCGCTGACGCTCCGGCAGCAAGGGGCGGAAGAGCGGCTTGCCGAACCTTGA
AGCCGCCGAAGCCCCGAAGCTTCCCCTTGACCAATGGTTCCCCGAAGACGCCGACGCTGA
CCCGACCGGCCCTAAGTCGTGGTGGGGGCGCGCGTCAGTAGACGACAAGCGCGTGTTCGT
CGGGCTCTTCGTAGACAAGATCGTTGTCACGAAGTCGACTACGGGCAGGGGGCAGGGAAC
GCCCATCGAGAAGCGCGCTTCGATCACGTGGGCGAAGCCGCCGACCGACGACGACGAAGA
CGACGCCCAGGACGGCACGGAAGACGTAGCGGCGTAGCGAGACACCCGGATCCCTCGAGG
GGCCCTATTCTATAGTGTACCTAAATGCTAGAGCTCGCTGATCAGCCTCGACTGTGCCT
TCTAGTTGCCAGCCATCTGTTGTTTGCCCCCTCCCCCGTGCCTTCCTTGACCCTGGAAGGT
GCCACTCCCCTGTCTTTCCTAATAAAATGAGGAAATTGCATCGCATTTGTCTGAGTAGG
TGTCATTCTATTCTGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGAC
AATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGCTTCTGAGGCGGAAGAACCAGG
TGCCAGTCATAGCCGAATAGCCTCTCCACCCAAGCGGCCGGAACCTGCGTGCAATCC
ACTGGGGGCGCG

Fig. 17

pCR-XL-TOPO-CMV-PUR-attB (SEQ ID NO: 10)

AGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTCAATTAATGCAGCTGGC
ACGACAGGTTTCCCGACTGGAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGC
TCACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAA
TTGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTAT
TTAGGTGACCGGTTAGAATACTCAAGCTATGCATCAAGCTTGGTACCGAGCTCGGATCCA
CTAGTAACGGCCGCCAGTGTGCTGGAATTCGCCCTTGGCCGCAATAAAATATCTTTATTT
TCATTACATCTGTGTGTTGGTTTTTTGTGTGAATCGATAGTACTAACATACGCTCTCCAT
CAAAACAAAACGAAACAAAACAACTAGCAAAATAGGCTGTCCCCAGTGCAAGTGCAGGT
GCCAGAACATTTCTCTATCGATAGGTACCGAGCTCTTACGCGTGCTAGCCCTCGAGCAGG
ATCTATACATTGAATCAATATTGGCAATTAGCCATATTAGTCATTGGTTATATAGCATAA
ATCAATATTGGCTATTGGCCATTGCATACGTTGTATCTATATCATAATATGTACATTTAT
ATTGGCTCATGTCCAATATGACCGCCATGTTGACATTGATTATTGACTAGTTATTAATAG
TAATCAATTACGGGGTTCATTAGTTCATAGCCCATATATGGAGTTCGCGCTTACATAACT
ACGGTAAATGGCCCGCTGGCTGACCGCCCAACGACCCCGCCCATTTGACGTCAATAATG
ACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTAT
TTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTCCGCCCCCT
ATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTACGG
GACTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATTGATGCGG
TTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCGAAGTCTC
CACCCCATTTGACGTCAATGGGAGTTTGTGTTTGGCACAAAATCAACGGGACTTTCCAAAA
TGTCGTAACAACCTCCGCCCCATTGACGCAATGGGCGGTAGGCGTGTACGGTGGGAGGTC
TATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGT
TTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGAAGCTCGACTCTAGG
GGCTCGAGATCTGCGATCTAAGTAAGCTTGCATGCCTGCAGGTGCGCCGCCACGACCGGT
GCCGCCACCATCCCCTGACCCACGCCCCCTGACCCCTCACAAGGAGACGACCTTCCATGAC
CGAGTACAAGCCCACGGTGCGCCCTCGCCACCCGCGACGACGTCCCCCGGGCCGTACGCAC
CCTCGCCGCCCGCTTCGCCGACTACCCCGCCACGCGCCACACCGTCGACCCGCGACCGCCA
CATCGAGCGGGTCACCGAGCTGCAAGAAGCTCTTCTCACGCGCGTCCGGCTCGACATCGG
CAAGGTGTGGGTGCGGGACGACGGCGCGCGGTGGCGGTCTGGACCACGCCGGAGAGCGT
CGAAGCGGGGGCGGTGTTGCGCGAGATCGGCCCCGCGCATGGCCGAGTTGAGCGGTTCCCG
GCTGGCCGCGCAGCAACAGATGGAAGGCTCTTGGCGCCGACCCGGCCCCAAGGAGCCCGC
GTGGTTCTTGGCCACCGTTCGGCGTCTCGCCCGACCAACAGGGCAAGGGTCTGGGCAGCGC
CGTCGTGCTCCCCGGAGTGGAGGCGGCCGAGCGCGCGGGGTGCCCGCTTCTGGAGAC
CTCCGCGCCCCGCAACCTCCCCCTTCTACGAGCGGCTCGGCTTACCCGTCACCGCCGACGT
CGAGGTGCCCCAAGGACCGCGCACCTGGTGCATGACCCGCAAGCCCGGTGCCTGACGCCC
GCCCCACGACCCGCGACGCGCCGACCGAAAGGAGCGCACGACCCCATGGCTCCGACCGAAG
CCGACCCGGGCGGCCCCCGCCGACCCCGCATGCAACGCCCCCGAGGCCACCGACTCTAGAGTC
GGGGCGGCCGCGGCTTCGAGCAGACATGATAAGATACATTGATGAGTTTGGACAAACCA
CAACTAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTAT
TTGTAACCATTATAAGCTGCAATAAACAAGTTAACAACAACAATTGCATTCATTTTATGT
TTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAGTAAAACCTCTACAAATGTG
GTAAATCGATAAGGATCAATTCGGCTTCAGGTACCGTCGACGATGTAGGTCACGGTCTC
GAAGCCGCGGTGCGGGTGCCAGGGCGTGCCTTGGGCTCCCCGGGCGCGTACTCCACCTC
ACCCATCTGGTCCATCATGATGAACGGGTGAGGTGGCGGTAGTTGATCCCGGCGAACGC
GCGGCGCACCGGGAAGCCCTCGCCCTCGAAACCGCTGGGCGCGGTGGTCACGGTGAGCAC
GGGACGTGCGACGGCGTTCGGCGGGTGCGGATACGCGGGGCGAGCGTCAGCGGGTTCTCGAC
GGTCACGGCGGGCATGTGACAGCCGAATTGATCCGTCGACCGATGCCCTTGAGAGCCCTT
CAACCCAGTCAGCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGAC
TGTCTTCTTTATCATGCAACTCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCTCGC
TCACTGACTCGCTGCGCTCGGTGCTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGG
CGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGAAGGCGCAAT
TCTGCAGATATCCATCACACTGGCGGCCGCTCGAGCATGCATCTAGAGGGCCCAATTCGC
CCTATAGTGAGTCGTATTACAATTCAGTGGCCGTGCTTTTACAACGTCGTGACTGGGAAA
ACCCTGGCGTTACCCAACCTTAATCGCCTTGCAGCACATCCCCCTTTCGCCAGCTGGCGTA
ATAGCGAAGAGGCCCCGACCGATCGCCCTTCCCAACAGTTGCGCAGCCTATACGTACGGC
AGTTTAAAGGTTTACACCTATAAAAGAGAGAGCCGTTATCGTCTGTTTGTGGATGTACAGA
GTGATATTATTGACACGCCGGGGCGACGGATGGTGATCCCCCTGGCCAGTGCACGTCTGC

TGTCAGATAAAGTCTCCCGTGAACCTTTACCCGGTGGTGTCATATCGGGGATGAAAGCTGGC
GCATGATGACCACCGATATGGCCAGTGTGCCGGTCTCCGTTATCGGGGAAGAAGTGGCTG
ATCTCAGCCACCGCGAAAATGACATCAAAAACGCCATTAACCTGATGTTCTGGGGAATAT
AAATGTGAGGCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTACGCTAGAAAAG
CCAGTCCGCAGAAACGGTGCTGACCCCGGATGAATGTCAGCTACTGGGCTATCTGGACAA
GGGAAAACGCAAGCGCAAAGAGAAAAGCAGGTAGCTTGCAGTGGGCTTACATGGCGATAGC
TAGACTGGGCGGTTTTATGGACAGCAAGCGAACCGGAATTGCCAGCTGGGGCGCCCTCTG
GTAAGGTTGGGAAGCCCTGCAAAGTAAACTGGATGGCTTTCTCGCCGCCAAGGATCTGAT
GGCGCAGGGGATCAAGCTCTGATCAAGAGACAGGATGAGGATCGTTTCGCATGATTGAAC
AAGATGGATTGCACGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTGCGCTATGACT
GGGCACAACAGACAATCGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTGAGCGCAGGGGC
GCCCCGTTCTTTTTGTCAAGACCGACCTGTCCGGTGCCCTGAATGAACTGCAAGACGAGG
CAGCGCGGCTATCGTGGCTGGCCACGACGGCGTTTCTTGCGCAGCTGTGCTCGACGTTG
TCACTGAAGCGGGAAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCAGGATCTCCTGT
CATCTCACCTTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGCGGCTGC
ATACGCTTGATCCGGCTACCTGCCCATTGACACCAAGCGAAACATCGCATCGAGCGAG
CACGTAATCGGATGGAAGCCGGTCTTGTCGATCAGGATGATCTGGACGAAGAGCATCAGG
GGCTCGCGCCAGCCGAACCTGTTGCCAGGCTCAAGGCGAGCATGCCGACGCGCAGGATC
TCGTCGTCGACCCATGGCGATGCCGTGCTTGCCGAATATCATGGTGGAATAATGCGCGCTTT
CTGGATTGATCGACTGTGGCCGGCTGGGTGTGGCGGACCGCTATCAGGACATAGCGTTGG
CTACCCGTGATATTGCTGAAGAGCTTGCGCGCGAATGGGCTGACCGCTTCTCTGCTGCTT
ACGGTATCGCCGCTCCCGATTGCGCAGCGCATCGCCTTCTATCGCCTTCTTGACGAGTTCT
TCTGAATTATTAACGCTTACAATTTCTGATGCGGTATTTTCTCCTTACGCATCTGTGCG
GTATTTTACACCGCATACAGGTGGCACTTTTCCGGGGAATGTGCGCGGAACCCCTATTTG
TTTATTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCGTGATAAAT
GCTTCAATAATAGCACGTGAGGAGGGCCACCATGGCCAAGTTGACCAAGTGCCGTTCCGGT
GCTCACCGCGCGCGACGTGCGCCGAGCGGTGAGATTCTGGACCGACCGGCTCGGGTTCTC
CCGGGACTTCTGAGGAGGACGACTTCGCGCGGTGTGGTCCGGGACGACGTGACCCGTGTT
CAGCGCGGTCCAGGACCAGGTGGTGCCGGAACAACCCCTGGCCTGGGTGTGGGTGCGCGG
CCTGGACGAGCTGTACGCCGAGTGGTCCGAGGTGCTGTCCACGAACCTCCGGGACGCTC
CGGGCCGGCCATGACCGAGATCGGCGAGCAGCCGTGGGGGCGGGAGTTGCGCCTGCGCGA
CCCGGCCGGCAACTGCGTGCACTTCTGCGCGAGGAGCAGGACTGACACGTGCTAAACT
TCATTTTAAATTTAAAGGATCTAGGTGAAGATCCTTTTGTATAATCTCATGACCAAAAT
CCCTTAACGTGAGTTTTCTGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATC
TTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGCAACAAAAAACCCGCT
ACCAGCGGTGGTTTTGTTTTGCCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAAGTGG
CTTCAGCAGAGCGCAGATACCAATACTGTCTTCTAGTGTAGCCGTAGTTAGGCCACCA
CTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAAGTGGC
TGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGA
TAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGGAGCGAAC
GACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGA
AGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAG
GGAGCTTCCAGGGGGAACGCCTGGTATCTTTATAGTCTGTCGGGTTTCGCCACCTCTG
ACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAG
CAACGCGGCCTTTTTACGTTTCTGGGCTTTTGCTGGCCTTTTGCTCACATGTTCTTTCC
TGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCGC
TCGCCGACCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAG

FIG. 18

SEQ ID NO: 11

GACTAGTACTGACGGACACACCGAAGCCCCGGCGGCAACCCTCAGCGGATGCCCCGGGGCTT
CACGTTTTCCCAGGTCAGAAGCGGTTTTTCGGGAGTAGTGCCCCAACTGGGGTAACCTTTGAG
TTCTCTCAGTTGGGGGCGTAGGGTCGCCGACATGACACAAGGGGTTGTGACCGGGGTGGACA
CGTACGCGGGTGCTTACGACCGTCAGTCGCGCGAGCGCGACTAGTACA

Fig. 19




Fig. 20


**Fig. 21**



Fig. 22




Fig. 23




Fig. 24





FIG. 25

**Fig. 26**

IgHeavy YAC
240 kb



Fig. 27A

IgLambda YAC
410 kb




Fig. 27B

025CIP7SEQ List.txt
SEQUENCE LISTING

<110> AviGeni cs, Inc

<120> ARTIFICIAL CHROMOSOMES IN AVIANS

<130> AVI-025CIP7PCT

<160> 16

<170> Patentln version 3.2

<210> 1

<211> 6230

<212> DNA

<213> Artificial Sequence

<220>

<223> Plasmid pCMV-31int

<400> 1

cattcgccat	tcaggctgcg	caactgttgg	gaagggcgat	cggtgcgggc	ctcttcgcta	60
ttacgccagc	caatacgcaa	accgcctctc	cccgcgcggt	ggccgattca	ttaatgcagg	120
atcgatccag	acatgataag	atacattgat	gagtttggac	aaaccacaac	tagaatgcag	180
tgaaaaaaat	gctttatttg	tgaaatttgt	gatgctattg	ctttatttgt	aaccattata	240
agctgcaata	aacaagttaa	caacaacaat	tgcattcatt	ttatgtttca	ggttcagggg	300
gaggtgtggg	agggttttta	aagcaagtaa	aacctctaca	aatgtgggat	ggctgattat	360
gatcatgaac	agactgtgag	gactgagggg	cctgaaatga	gccttgggac	tgtgaatcta	420
aaatacacaa	acaattagaa	tcactagctc	ctgtgtataa	tattttcata	aatcatactc	480
agtaagcaaa	actctcaagc	agcaagcata	tgcagctagt	ttaacacatt	atacacttaa	540
aaattttata	tttaccttag	agctttaaat	ctctgtaggt	agtttgtcca	attatgtcac	600
accacagaag	taaggttcct	tcacaaagat	ccaagctag	cttataatac	gactcactat	660
agggagagag	ctatgacgtc	gcatgcacgc	gtaagcttgg	gcccctcgag	ggatccgggt	720
gtctcgctac	gocgctacgt	cttccgtgcc	gtcctgggcg	tcgtcttcgt	cgctcgtcgt	780
cgggcggttc	gcccacgtga	tcgaagcgcg	cttctcgatg	ggcgttccct	gccccctgcc	840
cgtagtcgac	ttcgtgacaa	cgatcttgtc	tacgaagagc	ccgacgaaca	cgcgcttgtc	900
gtctactgac	gcgcgcccc	accacgactt	agggccgggc	gggtcagegt	cggcgtcttc	960
ggggaaccat	tgggtcaagg	gaagcttcgg	ggcttcggcg	gcttcaagtt	cggcaagccg	1020
ctcttcgcgc	ccttgctgcc	ggagcgtcag	cgctgcctgt	tgcttcggga	agtgcctcct	1080
gccaacgggt	ccgtcgtagc	cgctgcccgc	gcggtcttcg	tacagctctt	caagggcgtt	1140
cagggcgctc	gcgcgctccg	caacaagggt	cgcccgcttc	ccgctcttct	caggcgcttc	1200
agtgagcttg	ccgaagcgtc	gggcgggttc	ccacagaagc	gccaacgtct	cttcgctgcc	1260

025CIP7SEQ List.tXt

ttcggcgtgc	ctgatcttgt	tgaagatgcg	ttccgcaacg	aacttgtega	gtgccgccat	1320
gctgacgttg	cacgtgcctt	cgtgctgccc	aggtgcggac	gggtcgacca	ccttcgggcg	1380
acggcagcgg	taagagtcct	tgatcgattc	ttccccgcgc	ttcgaagtca	tgacggcgcc	1440
acactcgcag	tacagcttgt	ccatggcgga	cagaatggct	tgccccggg	aaagccccctt	1500
gccgcgcccc	ctgccgtcca	accacgcctg	aagctcatac	cactcagcgg	gctcgatgat	1560
cgggtccgcaa	tcaagctcga	ccggccggag	cgtgatcggg	tcgcgctgaa	tgcggttaacc	1620
ctcaatcttc	gtggctggcg	tgccgtccgg	cttcttcttg	tagatcacct	cagcggcgaa	1680
gcccgcataa	cgcgggtccc	gaaggattcg	cataacggtt	gccgggtccc	aggcgcttga	1740
agcggctcttc	ttcccaatcg	tctcgccccg	ggtcggcacg	gcgtcagcgt	ccatgcgctt	1800
acaaagcccc	gtgatgctgc	ccgggtgaat	ggcggttga	ctgcccggt	tgaagggaag	1860
gtgtttgtgc	gtcttgatct	cacgccacca	ccaccggatt	acgtcgggct	cgaactcgaa	1920
gggtccggta	aggggagtg	tcgagtgcgc	aagcttggtg	atgacgacat	tgaccattcg	1980
gccgttgccg	gtgatctcct	tcgtctccga	aacaagctcg	aagccgtaag	gcgccttccc	2040
gccgacgtac	ccgccaatt	cgcgctgaag	gttcttcgtg	tcgagaatct	tcgccgactt	2100
cagcgaagat	tctttgtg	acgcgtcgag	ccgcataatc	aggtgaatca	ggtccatgac	2160
gtttccctgc	cggaagacgc	cttcctgagt	ggaacaatc	gtcacgcca	gggcgagcaa	2220
ttccgagaca	atcggaatcg	cgtccatgac	cttcaggcgc	gagaagcgcg	acacgtcata	2280
gacaatgate	atgttgagcc	gcccggcgcg	gcattcggtt	aggatgcgtt	cgaactccgg	2340
gcgctccgcc	gtcccgaacg	ccgacgtgcc	cggcgcttcg	ctgaaatgcc	cgacgaacct	2400
gaaccggccc	ccgtcgcgct	cgacttcgcg	ctgaaggctg	gccgccttgt	cttcgttggc	2460
gctacgctgt	gtcgtggggc	ttgctgcgct	cgaattctcg	cgctcgcgcg	actgacggtc	2520
gtaagcaccc	gcgtacgtgt	ccaccccggt	cacaaccct	tgtgtcatgt	cggcgaccct	2580
acgactagt	agctcgtcga	cccgggaatt	ccggaccggt	acctgcaggc	gtaccttcta	2640
tagtgtcacc	taaatagctt	tttgcaaaag	cctaggctag	agtcgggagg	ctggatcggt	2700
cccgggtgtct	tctatggagg	tcaaaacagc	gtggatggcg	tctccaggcg	atctgacggt	2760
tcactaaacg	agctctgctt	atatagacct	cccaccgtac	acgcctaccg	cccatttgcg	2820
tcaatggggc	ggagttgtta	cgacattttg	gaaagtcccg	ttgatttttg	tgccaaaaca	2880
aactcccatt	gacgtcaatg	gggtggagac	ttggaaatcc	ccgtgagtca	aaccgctatc	2940
cacgcccatt	gatgtactgc	caaaaccgca	tcaccatggg	aatagcgatg	actaatacgt	3000
agatgtactg	ccaagtagga	aagtcccata	aggtcatgta	ctgggcataa	tgccaggcgg	3060
gccatttacc	gtcattgacg	tcaatagggg	gcgtacttgg	catatgatac	acttgatgta	3120

				025CIP7SEQ	List.tXt	
ctgccaaagtg	ggcagtttac	cgtaaatact	ccacccattg	acgtcaatgg	aaagtccta	3180
ttggcgttac	tatgggaaca	tacgtcatta	ttgacgtcaa	tgggcggggg	tcgttgggcg	3240
gtcagccagg	cgggccattt	accgtaagtt	atgtaacgac	ctgcacgatg	ctgtttcctg	3300
tgtgaaattg	ttatccgctc	acaattccac	acattatacg	agccggaagc	tataaagtgt	3360
aaagcctggg	gtgcctaagt	agtgaaaggg	cctcgataac	gcctattttt	ataggttaat	3420
gtcatgataa	taatggtttc	ttagacgtca	ggtggcactt	ttcggggaaa	tgtgcgcgga	3480
acccctattt	gtttattttt	ctaaatacat	tcaaatatgt	atccgctcat	gagacaataa	3540
ccctgataaa	tgcttcaata	atattgaaaa	acgcgcgaat	tgcaagctct	gcattaatga	3600
atcggccaac	gcgcggggag	aggcggtttg	cgtattgggc	gctcttcgcg	ttcctcgctc	3660
actgactcgc	tgcgctcggt	cgttcggctg	cggcgagcgg	tatcagctca	ctcaaaggcg	3720
gtaatacggg	tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	agcaaaaggc	3780
cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	taggctcgcg	3840
ccccctgacg	agcatcacia	aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgacagga	3900
ctataaagat	accaggcggt	tccccctgga	agctccctcg	tgcgctctcc	tgttccgacc	3960
ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcaa	4020
tgtcacgct	gtaggtatct	cagttcggtg	taggtcgttc	gctccaagct	gggctgtgtg	4080
cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	4140
aaccgcgtaa	gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	4200
gcgaggatat	taggcggtgc	tacagagttc	ttgaagtggg	ggcctaacta	cggctacact	4260
agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagagtt	4320
ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	gtggtttttt	tgtttgcaag	4380
cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	4440
tctgacgctc	agtggaacga	aaactcacgt	taagggaattt	tggatcatgcc	ataacttcgt	4500
atagcataca	ttatacgaag	ttatggcatg	agattatcaa	aaaggatctt	cacctagatc	4560
cttttaaatt	aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	aacttgggtc	4620
gacagttacc	aatgcttaat	cagtgaggca	cctatctcag	cgatctgtct	atttcgttca	4680
tccatagttg	cctgactccc	cgtcgtgtag	ataactacga	tacgggaggg	cttaccatct	4740
ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	cggctccaga	tttatcagca	4800
ataaaccagc	cagccggaag	ggccgagcgc	agaagtgggc	ctgcaacttt	atccgcctcc	4860
atccagtcta	ttaattggtg	ccgggaagct	agagtaagta	gttcgccagt	taatagtttg	4920
cgaacgttg	ttgccattgc	tacaggcatc	gtggtgtcac	gctcgtcgtt	tggatggct	4980
tcattcagct	ccggttccca	acgatcaagg	cgagttacat	gatcccccat	gttgtgcaaa	5040

025CIP7SEQ List.tXt

```

aaagcgggta gtccttcggt tcctccgata gttgtcagaa gtaagttggc cgcagtgtta 5100
tactcatggg ttatggcagc actgcataat tctcttactg tcatgccatc cgtaagatgc 5160
ttttctgtga ctggtgagta ctcaaccaag tcattctgag aatagtgtat gcggcgaccg 5220
agttgctctt gcccggcgtc aatacgggat aataccgcgc cacatagcag aactttaaaa 5280
gtgctcatca ttggaaaacg ttcttcgggg cgaaaactct caaggatctt accgctgttg 5340
agatccagtt cgatgtaacc cactcgtgca cccaactgat cttcagcatc ttttactttc 5400
accagcgttt ctgggtgagc aaaaacagga aggcaaaatg ccgcaaaaaa gggaataagg 5460
gcgacacgga aatgttgaat actcatactc ttcccttttc aatattattg aagcatttat 5520
caggggttatt gtctcatgcc aggggtgggc acacatatatt gataccagcg atccctacac 5580
agcacataat tcaatgcgac ttccctctat cgcacatctt agacctttat tctccctcca 5640
gcacacatcg aagctgccga gcaagccgtt ctcaccagtc caagacctgg catgagcgga 5700
tacatatattg aatgtattta gaaaaataaa caaatagggg ttccgcgcac atttccccga 5760
aaagtgccac ctgaaattgt aaacgttaat attttggtta aattcgcgtt aaatttttgt 5820
taaatacagct cattttttaa ccaataggcc gaaatcggca aaatccctta taaatcaaaa 5880
gaatagaccg agatagggtt gagtgttggt ccagtttgga acaagagtcc actattaaag 5940
aacgtggact ccaacgtcaa agggcgaaaa accgtctatc agggcgatgg cccactacgt 6000
gaaccatcac cctaatacaag ttttttgggg tcgaggtgcc gtaaagcact aaatcggaac 6060
cctaaaggga gccccgatt tagagcttga cggggaaagc cggcgaaacgt ggcgagaaag 6120
gaagggaaga aagcgaaaag agcgggcgct agggcgctgg caagtgtagc ggtcacgctg 6180
cgcgtaacca ccacacccgc cgcgcttaat gcgcgcgtac agggcgcgtc 6230

```

```

<210> 2
<2U> 5982
<212> DNA
<213> Artificial sequence

```

```

<220>
<223> Plasmid pCMV-luc-attB

```

```

<400> 2
ctctatcgat aggtaccgag ctcttacgcg tgctagccct cgagcaggat ctatacattg 60
aatcaatatt ggcaattagc catattagtc attggttata tagcataaat caatattggc 120
tattggccat tgcatacgtt gtatctatat cataatatgt acatttatat tggctcatgt 180
ccaatatgac cgccatgttg acattgatta ttgactagtt attaatagta atcaattacg 240
gggtcattag ttcatagccc atatatggag ttccgcgtta cataacttac ggtaaatggc 300
ccgcctggct gaccgcccaa cgaccccgcc ccattgacgt caataatgac gtatgttccc 360

```

O25CIP7SEQ List.tXt

atagtaacgc	caatagggac	tttccattga	cgtcaatggg	tggagtattt	acggtaaaact	420
gcccacttgg	cagtacatca	agtgtatcat	atgccaaagtc	cgeccccctat	tgacgtcaat	480
gacggtaaat	ggcccgctg	gcattatgcc	cagtacatga	ccttacggga	ctttcctact	540
tggcagtaca	tctacgtatt	agtcatcgct	attaccatgg	tgatgcgggt	ttggcagtac	600
atcaatgggc	gtggatagcg	gtttgactca	cggggatttc	caagtctcca	ccccattgac	660
gtcaatggga	gtttgttttg	gcacccaaaat	caacgggact	ttccaaaatg	tcgtaacaac	720
tccgccccat	tgacgcaaat	gggcggtagg	cgtgtacggg	gggagggtcta	tataagcaga	780
gctcgtttag	tgaaccgtca	gatcgctgg	agacgccatc	cacgctgttt	tgacctccat	840
agaagacacc	gggaccgatc	cagcctcccc	tcgaagctcg	actctagggg	ctcgagatct	900
gcgatctaag	taagcttggc	attccggtac	tgttggtaaa	gccaccatgg	aagacgccaa	960
aaacataaag	aaaggccccg	cgccattcta	tccgctggaa	gatggaaccg	ctggagagca	1020
actgcataag	gctatgaaga	gatacgccct	ggttccctgga	acaattgctt	ttacagatgc	1080
acatatcgag	gtggacatca	cttacgctga	gtacttcgaa	atgtccgttc	ggttggcaga	1140
agctatgaaa	cgatatgggc	tgaatacaaaa	tcacagaatc	gtcgtatgca	gtgaaaaactc	1200
tcttcaattc	tttatgccgg	tgttggggcg	gttattttatc	ggagttgcag	ttgcgccccg	1260
gaacgacatt	tataatgaac	gtgaattgct	caacagtatg	ggcatttcgc	agcctaccgt	1320
ggtgttcggt	tccaaaaagg	ggttgcaaaa	aattttgaac	gtgcaaaaaa	agctcccaat	1380
catccaaaaa	attattatca	tggattctaa	aacggattac	cagggatttc	agtcgatgta	1440
cacgttcgct	acatctcctc	tacctcccg	ttttaatgaa	tacgattttg	tgccagagtc	1500
cttcgatagg	gacaagacaa	ttgcactgat	catgaactcc	tctggatcta	ctgggtctgcc	1560
taaaggtgtc	gctctgcctc	atagaactgc	ctgcgtgaga	ttctcgcgatg	ccagagatcc	1620
tattttttggc	aatcaaatca	ttccggatac	tgcgatttta	agtgttggtc	cattccatca	1680
cggtttttgg	atgtttacta	cactcggata	tttgatatgt	ggatttcgag	tcgtcttaaat	1740
gtatagattt	gaagaagagc	tgtttctgag	gagccttcag	gattacaaga	ttcaaagtgc	1800
gctgctggtg	ccaaccctat	tctccttctt	cgccaaaagc	actctgattg	acaaatacga	1860
tttatctaata	ttacacgaaa	ttgcttctgg	tggcgctccc	ctctctaagg	aagtccgggga	1920
agcggttgcc	aagagggtcc	atctgccagg	tatcaggcaa	ggatatgggc	tactgagac	1980
tacatcagct	attctgatta	cacccgaggg	ggatgataaa	ccgggcgcgg	tcggtaaagt	2040
tgttccattt	tttgaagcga	aggttgtgga	tctggatacc	gggaaaacgc	tgggcgttaa	2100
tcaaagaggc	gaactgtgtg	tgagagggtcc	tatgattatg	tccggttatg	taaacaatcc	2160
ggaagcgacc	aacgccttga	ttgacaagga	tggatggcta	cattctggag	acatagctta	2220
ctgggacgaa	gacgaacaact	tcttcatcgt	tgaccgcctg	aagtctctga	ttaagtacaa	2280

025CIP7SEQ List.tXt

aggetatcag	gtggctccccg	ctgaattgga	atccatcttg	ctccaacacc	ccaacatctt	2340
cgacgcaggt	gtcgcaggtc	ttccccgacg	tgacgcgggt	gaacttccccg	ccgccgttgt	2400
tgttttgag	cacggaaaga	cgatgacgga	aaaagagatc	gtggattacg	tcgccagtca	2460
agtaacaacc	gcgaaaaagt	tgcgcggagg	agttgtgttt	gtggacgaag	taccgaaagg	2520
tcttaccgga	aaactcgacg	caagaaaaat	cagagagatc	ctcataaagg	ccaagaaggg	2580
cggaaagatc	gccgtgtaat	tctagagtcg	gggcggccgg	ccgcttcgag	cagacatgat	2640
aagatacatt	gatgagtttg	gacaaaccac	aactagaatg	cagtgaaaaa	aatgctttat	2700
ttgtgaaatt	tgtgatgcta	ttgctttatt	tgttaaccatt	ataagctgca	ataaacaagt	2760
taacaacaac	aattgcattc	attttatgtt	tcaggttcag	ggggagggtgt	gggagggtttt	2820
ttaaagcaag	taaaacctct	acaaatgtgg	taaaatcgat	aaggatcaat	tcggcttcag	2880
gtaccgtcga	cgatgtaggt	cacggtctcg	aagccgcgggt	gcgggtgccca	gggcgtgccc	2940
ttgggctccc	cgggcgcgta	ctccacctca	cccatctggt	ccatcatgat	gaacgggtcg	3000
aggtggcgggt	agttgatccc	ggcgaacgcg	cggcgcaccg	ggaagccctc	gccctcgaaa	3060
ccgctggggcg	cgggtggtcac	ggtgagcacg	ggacgtgcga	cggcgtcggc	gggtgcggat	3120
acgcggggca	gcgtcagcgg	gttctcgacg	gtcacggcgg	gcatgtcgac	agccgaattg	3180
atccgtcgac	cgatgccctt	gagagccttc	aaccacgtca	gctccttcgg	gtgggcgcgg	3240
ggcatgacta	tcgtcgccgc	acttatgact	gtcttcttta	tcatgcaact	cgtaggacag	3300
gtgccggcag	cgtctcttcg	cttctctgct	cactgactcg	ctgcgctcgg	tcgttcgggt	3360
gcggcgagcg	gtatcagctc	actcaaaggc	ggtaatacgg	ttatccacag	aatcagggga	3420
taacgcagga	aagaacatgt	gagcaaaagg	ccagcaaaag	gccaggaacc	gtaaaaaggc	3480
cgcgttgctg	gcgtttttcc	ataggctccg	ccccctgac	gagcatcaca	aaaatcgacg	3540
ctcaagtcag	aggtggcgaa	acccgacagg	actataaaga	taccaggcgt	ttccccctgg	3600
aagctccctc	gtgcgctctc	ctgttccgac	cctgcgcgtt	accggatacc	tgtccgcctt	3660
tctcccttcg	ggaagcgtgg	cgttttctca	atgctcacgc	tgtaggtatc	tcagttcgggt	3720
gtaggtcggt	cgtcccaagc	tgggctgtgt	gcacgaaccc	cccgttcagc	ccgaccgctg	3780
cgccttatcc	ggtaactatc	gtcttgagtc	caacccggtg	agacacgact	tatcgccact	3840
ggcagcagcc	actggtaaca	ggattagcag	agcgagggtat	gtaggcggtg	ctacagagtt	3900
cttgaagtgg	tggcctaact	acggctacac	tagaaggaca	gtatttggtg	tctgcgctct	3960
gctgaagcca	gttaccttcg	gaaaaagagt	tggtagctct	tgatccggca	aacaaaccac	4020
cgcgtggtagc	ggtgggtttt	ttgtttgcaa	gcagcagatt	acgcgcagaa	aaaaaggatc	4080
tcaagaagat	cctttgatct	tttctacggg	gtctgacgct	cagtggaacg	aaaactcacg	4140

			025CIP7SEQ	List.tXt		
ttaagggatt	ttgggtcatga	gattatcaaa	aaggatccttc	acctagatcc	ttttaaatta	4200
aaaatgaagt	tttaaatacaa	tctaaagtat	atatgagtaa	acttgggtctg	acagttacca	4260
atgcttaatc	agtgaggcac	ctatctcagc	gatctgtcta	tttcgttcat	ccatagttgc	4320
ctgactcccc	gtcgtgtaga	taactacgat	acgggagggc	ttaccatctg	gccccagtgc	4380
tgcaatgata	ccgcgagacc	cacgctcacc	ggctccagat	ttatcagcaa	taaaccagcc	4440
agccggaagg	gccgagcgca	gaagtgggtcc	tgcaacttta	tccgcctcca	tccagtctat	4500
taattgttgc	cggaagcta	gagtaagtag	ttcgccagtt	aatagtttgc	gcaacgttgt	4560
tgccattgct	acaggcatcg	tggtgtcacg	ctcgtcgttt	ggatatggctt	cattcagctc	4620
cggttcccaa	cgatcaaggc	gagttacatg	atccccatg	ttgtgcaaaa	aagcggttag	4680
ctccttcggt	cctccgatcg	ttgtcagaag	taagttggcc	gcagtgttat	cactcatggt	4740
tatggcagca	ctgcataatt	ctcttactgt	catgccatcc	gtaagatgct	tttctgtgac	4800
tggtgagtac	tcaaccaagt	cattctgaga	atagtgtatg	cggcgaccga	gttgctcttg	4860
cccggcgta	atacgggata	ataccgcgcc	acatagcaga	actttaaaag	tgtcatcat	4920
tggaacacgt	tcttcggggc	gaaaactctc	aaggatctta	ccgctgttga	gatccagttc	4980
gatgtaaccc	actcgtgcac	ccaactgatc	ttcagcatct	tttactttca	ccagcgtttc	5040
tgggtgagca	aaaacaggaa	ggcaaaatgc	cgcaaaaaag	ggaataaggg	cgacacggaa	5100
atgttgaata	ctcatactct	tcctttttca	atattattga	agcatttatc	agggttattg	5160
tctcatgagc	ggatacatat	ttgaatgtat	ttagaaaaat	aaacaaatag	gggttccgcg	5220
cacatttccc	cgaaaagtgc	cacctgacgc	gccctgtagc	ggcgcattaa	gcgcggcggg	5280
tgtggtggtt	acgcgcagcg	tgaccgctac	acttgccagc	gccctagcgc	ccgctccttt	5340
cgttttcttc	ccttcctttc	tcgccacggt	cgccggcttt	ccccgtcaag	ctctaaatcg	5400
ggggctccct	ttagggttcc	gatttagtgc	tttacggcac	ctcgaccca	aaaaacttga	5460
ttagggtgat	ggttcacgta	gtgggccatc	gccctgatag	acggtttttc	gccctttgac	5520
gttgaggtcc	acgttcttta	atagtggact	cttgttccaa	actggaacaa	cactcaaccc	5580
tatctcggtc	tattcttttg	atctataagg	gattttgccg	atttcggcct	attggttaaa	5640
aatgagctg	atttaacaaa	aatttaacgc	gaattttaac	aaaatattaa	cgtttacaat	5700
ttcccatcgc	ccattcaggc	tgcgcaactg	ttgggaaggg	cgatcggtgc	gggcctcttc	5760
gctattacgc	cagcccaagc	taccatgata	agtaagtaat	attaaggtac	gggaggtact	5820
tggagcggcc	gcaataaaaat	atctttatct	tcattacatc	tgtgtgttgg	ttttttgtgt	5880
gaatcgatag	tactaacata	cgtctctccat	caaaacaaaa	cgaaacaaaa	caaaactagca	5940
aaataggctg	tccccagtgc	aagtgcaggt	gccagaacat	tt		5982

025CIP7SEQ Li st.tXt

<210> 3
 <211> 5924
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Plasmid pCMV-luc-attP

<400> 3
 ctctatcgat aggtaccgag ctcttacgcg tgctagccct cgagcaggat ctatacattg 60
 aatcaatatt ggcaattagc catattagtc attggttata tagcataaat caatattggc 120
 tattggccat tgcatacggt gtatctatat cataatatgt acatttatat tggctcatgt 180
 ccaatatgac cgccatgttg acattgatta ttgactagtt attaatagta atcaattacg 240
 gggtcattag ttcatagccc atatatggag ttccgcgtta cataacttac ggtaaattggc 300
 ccgcctggct gaccgcccac cgacccccgc ccattgacgt caataatgac gtatgttccc 360
 atagtaacgc caatagggac tttccattga cgtcaatggg tggagtattt acggtaaaact 420
 gccacttggt cagtacatca agtgtatcat atgccaagtc cgccccctat tgacgtcaat 480
 gacggtaaat ggccgccttg gcattatgcc cagtacatga ccttacggga ctttcctact 540
 tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcgggt ttggcagtac 600
 atcaatgggc gtggatagcg gtttgactca cggggatttc caagtctcca ccccatlgac 660
 gtcaatggga gtttggtttg gcacaaaaat caacgggact ttccaaaatg tcgtaacaac 720
 tccgccccat tgacgcaaat gggcggtagg cgtgtacggg gggagggtcta tataagcaga 780
 gctcgttttag tgaaccgtca gatcgccctg agacgccatc cacgctgttt tgacctccat 840
 agaagacacc gggaccgatc cagcctcccc tcgaagctcg actctagggg ctcgagatct 900
 gcgatctaag taagcttggc attccggtac tgttggtaaa gccaccatgg aagacgcccac 960
 aacataaaag aaaggcccg gcccattcta tccgctggaa gatggaaccg ctggagagca 1020
 actgcataag gctatgaaga gatacgccct ggttcctgga acaattgctt ttacagatgc 1080
 acatategag gtggacatca cttacgctga gtacttcgaa atgtccgttc gggttggcaga 1140
 agctatgaaa cgatatgggc tgaatacaaa tcacagaatc gtcgtatgca gtgaaaactc 1200
 tcttcaattc tttatgccgg tgttggggcg gttatttata ggagttgcag ttgcgccccg 1260
 gaacgacatt tataatgaac gtgaattgct caacagtatg ggcatttcgc agcctaccgt 1320
 ggtgttcgtt tccaaaaagg ggttgcaaaa aattttgaac gtgcaaaaaa agctcccaat 1380
 catccaaaaa attattatca tggattctaa aacggattac cagggttttc agtcgatgta 1440
 cacgttcgtc acatctcatc tacctcccg ttttaatgaa tacgattttg tgccagagtc 1500
 cttcgatagg gacaagacaa ttgcactgat catgaactcc tctggatcta ctggtctgcc 1560
 taaagggtgtc gctctgcctc atagaactgc ctgcgtgaga ttctcgcatg ccagagatcc 1620

			025CIP7SEQ	List.tXt		
tatTTTTTggc	aatcaaatca	ttccggatac	tgcgatttta	agtgttggtc	cattccatca	1680
cggTTTTTgga	atgtttacta	cactcggata	tttgatatgt	ggatttcgag	tcgtcttaat	1740
gtatagatTTT	gaagaagagc	tgTTTctgag	gagccttcag	gattacaaga	ttcaaagtgc	1800
gctgctggTg	ccaacctat	tctccttctt	cgccaaaagc	actctgattg	acaaatacga	1860
TTtatctaT	ttacacgaaa	ttgcttctgg	tggcgctccc	ctctctaagg	aagtcgggga	1920
agcggTtgcc	aagaggTtcc	atctgccagg	tatcaggcaa	ggatatgggc	tactgagac	1980
tacatcagct	attctgatta	cacccgaggg	ggatgataaa	ccgggcgcgg	tcggtaaagt	2040
tgTtccattT	TTTgaagcga	aggTtTgTga	tctggatacc	gggaaaacgc	tgggcgttaa	2100
tcaaagaggc	gaactgtgtg	tgagaggTcc	tatgattatg	tccggttatg	taaacaatcc	2160
ggaagcgacc	aacgcctTga	ttgacaagga	tggatggcta	cattctggag	acatagctta	2220
ctgggacgaa	gacgaacact	tcttcatcgt	tgaccgcctg	aagtctctga	ttaagtacaa	2280
aggctatcag	gtggctcccg	ctgaattTga	atccatcttg	ctccaacacc	ccaacatctt	2340
cgacgcaggt	gtcgcaggtc	ttcccgacga	tgacgcgggt	gaacttcccg	ccgccgttgt	2400
tgTTTTggag	cacggaaaga	cgatgacgga	aaaagagatc	gtggattacg	tcgccagtca	2460
agtaacaacc	gcgaaaaagt	tgcgcggagg	agTtgtgttt	gtggacgaag	taccgaaagg	2520
tcttaccgga	aaactcgacg	caagaaaaat	cagagagatc	ctcataaagg	ccaagaaggg	2580
cggaaagatc	gccgtgtaat	tctagagTcg	gggcggccgg	ccgcttcgag	cagacatgat	2640
aagatacatt	gatgagTttg	gacaaaccac	aactagaatg	cagtgaaaaa	aatgctttat	2700
ttgtgaaatt	tgtgatgcta	ttgctttatt	tgtaacatt	ataagctgca	ataaacaagt	2760
taacaacaac	aattgcattc	atTTtatgtt	tcaggTtcag	ggggaggTgt	gggaggTttt	2820
TTaaagcaag	Taaaacctct	acaaatgtgg	Taaaatcgat	aaggatcaat	tcggcttcga	2880
ctagtactga	cggacacacc	gaagccccgg	cggcaaccct	cagcggatgc	cccggggctt	2940
cacgtTTTcc	caggtcagaa	gcggTTTTcg	ggagtagTgc	cccaactggg	gtaacctTTg	3000
agTtctctca	gttgggggcg	tagggTcgcc	gacatgacac	aaggggtTgt	gaccggggTg	3060
gacacgtacg	cggtgctta	cgaccgtcag	tcgcgcgagc	gcgactagta	caagccgaat	3120
Tgatccgtcg	accgatgccc	ttgagagcct	tcaaccagT	cagTccttc	cggTgggcgc	3180
ggggcatgac	tatcgtcgcc	gcacttatga	ctgtcttctt	tatcatgcaa	ctcgtaggac	3240
aggTgccggc	agcgtcttc	cgcttcctcg	ctcactgact	cgctgcgctc	ggtcgTtcgg	3300
ctgcggcgag	cggtatcagc	tactcaaag	gcggtaatac	ggttatccac	agaatcaggg	3360
gataacgcag	gaaagaacat	gtgagcaaaa	ggccagcaaa	aggccaggaa	ccgTaaaaag	3420
gccgcgtTgc	tggcgTTTTT	ccataggTct	cgccccctg	acgagcatca	caaaaatcga	3480
cgtcaagTc	agaggTggcg	aaaccgcaca	ggactataaa	gataaccaggc	gtttccccct	3540

025CIP7SEQ List.txt

ggaagctccc	tcgtgcgctc	tcctgttccg	accctgccgc	ttaccggata	cctgtccgcc	3600
tttctccctt	cgggaagcgt	ggcgctttct	caatgctcac	gctgtaggta	tctcagttcg	3660
gtgtaggtcg	ttcgctccaa	gctgggctgt	gtgcacgaac	cccccgttca	gcccgaaccgc	3720
tgcgccttat	ccggtaacta	tcgtcttgag	tccaacccgg	taagacacga	cttatcgcca	3780
ctggcagcag	ccactggtaa	caggattagc	agagcgaggt	atgtaggcgg	tgctacagag	3840
ttcttgaagt	ggtggcctaa	ctacggctac	actagaagga	cagtatttgg	tatctgcgct	3900
ctgctgaagc	cagttacctt	cggaaaaaga	gttggtagct	cttgatccgg	caaaaaaacc	3960
accgctggta	gcggtggttt	ttttgtttgc	aagcagcaga	ttacgcgcag	aaaaaaagga	4020
tctcaagaag	atcctttgat	cttttctacg	gggtctgacg	ctcagtggaa	cgaaaaactca	4080
cgtaagggga	ttttggtcac	gagattatca	aaaaggatct	tcacctagat	ccttttaaat	4140
taaaaaatgaa	gttttaaatc	aatctaaagt	atatatgagt	aaacttggtc	tgacagttac	4200
caatgcttaa	tcagtgaggc	acctatctca	gcgatctgtc	tatttcgttc	atccatagtt	4260
gcctgactcc	ccgtcgtgta	gataactacg	atacgggagg	gcttaccatc	tggtcccgagt	4320
gctgcaatga	taccgcgcaga	cccacgctca	ccggctccag	atttatcagc	aataaaccag	4380
ccagccggaa	gggcccagcg	cagaagtggc	cctgcaactt	tatccgcctc	catccagtct	4440
attaattggt	gccgggaagc	tagagtaagt	agttcgccag	ttaatagttt	gcgcaacggt	4500
gttgccattg	ctacaggcat	cgtggtgtca	cgctcgtcgt	ttggtatggc	ttcattcagc	4560
tccggttccc	aacgatcaag	gcgagttaca	tgatccccc	tggtgtgcaa	aaaagcggtt	4620
agctccttcg	gtcctccgat	cgttgtcaga	agtaagttgg	ccgcagtgtt	atcactcatg	4680
gttatggcag	cactgcataa	ttctcttact	gtcatgccat	ccgtaagatg	cttttctgtg	4740
actggtgagt	actcaaccaa	gtcattctga	gaatagtgt	tgccggcgacc	gagttgctct	4800
tgcccggcgt	caatacggga	taataccgcg	ccacatagca	gaactttaaa	agtgtcatc	4860
attggaanaac	gttcttcggg	gcgaaaactc	tcaaggatct	taccgctggt	gagatccagt	4920
tcgatgtaac	ccactcgtgc	acccaactga	tcttcagcat	cttttacttt	caccagcgtt	4980
tctgggtgag	caaaaacagg	aaggcaaat	gccgcaaaaa	agggataaag	ggcgacacgg	5040
aaatgttgaa	tactcatact	cttccttttt	caatattatt	gaagcattta	tcagggttat	5100
tgtctcatga	gcggatacat	atttgaatgt	atttagaaaa	ataaacaat	aggggttccg	5160
cgcacatttc	cccgaanaag	gccacctgac	gcgccttgta	gcggcgcat	aagcgcgcg	5220
ggtgtggtgg	ttacgcgcag	cgtgaccgct	acacttgcca	gcgccttagc	gcccgctcct	5280
ttcgctttct	tcccttcctt	tctcgccacg	ttcgccggct	ttccccgtca	agctctaaat	5340
cgggggctcc	ctttagggtt	ccgatttagt	gctttacggc	acctcgacct	caaaaaactt	5400

O25CIP7SEQ List.txt

gattaggggtg	atgggttcacg	tagtggggcca	tcgccctgat	agacgggtttt	tcgccctttg	5460
acgttggagt	ccacgttctt	taatagtgga	ctcttggtcc	aaactggaac	aacactcaac	5520
cctatctcgg	tctattcttt	tgatttataa	gggattttgc	cgatttcggc	ctattggtta	5580
aaaaatgagc	tgatttaaca	aaaatttaac	gcgaatttta	acaaaatatt	aacgtttaca	5640
atttcccatt	cgccattcag	gctgcgcaac	tgttgggaag	ggcgatcggg	gcgggcctct	5700
tcgctattac	gccagcccaa	gctaccatga	taagtaagta	atattaaggt	acgggaggta	5760
cttgagcgg	ccgcaataaa	atatctttat	tttcattaca	tctgtgtgtt	ggttttttgt	5820
gtgaatcgat	agtactaaca	tacgctctcc	atcaaaacaa	aacgaaacaa	aacaaactag	5880
caaaataggc	tgtccccagt	gcaagtgcag	gtgccagaac	attt		5924

<210> 4
 <211> 5101
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Plasmid pCMV-pur-attB

<400> 4

ctagagtcgg	ggcggccggc	cgcttcgagc	agacatgata	agatacattg	atgagtttgg	60
acaaaccaca	actagaatgc	agtgaaaaaa	atgctttatt	tgtgaaattt	gtgatgctat	120
tgctttat	gtaaccatta	taagctgcaa	taaacaagtt	aacaacaaca	attgcattca	180
ttttatgt	cagggtcagg	gggaggtgtg	ggagggtttt	taaagcaagt	aaaacctcta	240
caaagtgt	aaaatcgata	aggatcaatt	cggttcagg	taccgtcgac	gatgtaggtc	300
acggtctcga	agccgcgggtg	cggttgccag	ggcgtgccct	tgggctcccc	gggcgcgtac	360
tccacctcac	ccatctggtc	catcatgatg	aacgggtcga	ggtggcggtg	gttgatcccg	420
gcgaacgcgc	ggcgcaccgg	gaagccctcg	ccctcgaaac	cgctgggcgc	ggtggtcacg	480
gtgagcacgg	gacgtgcgac	ggcgtcggcg	ggtgcggata	cgcggggcag	cgtcagcggg	540
ttctcgacgg	tcacggcggg	catgtcgaca	gccgaattga	tccgtcgacc	gatgcccttg	600
agagccttca	accagtcag	ctccttcgg	tgggcgcggg	gcatgactat	cgtcgccgca	660
cttatgactg	tcttctttat	catgcaactc	gtaggacagg	tgccggcagc	gctcttcgc	720
ttcctcgctc	actgactcgc	tgcgctcgg	cgcttcggctg	cgcgagcgg	tatcagctca	780
ctcaaaggcg	gtaatacgg	tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	840
agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	900
taggctccgc	ccccctgacg	agcatcacia	aaatcgacgc	tcaagtcaga	ggtggcgaaa	960
cccgacagga	ctataaagat	accaggcggt	tccccctgga	agctccctcg	tgcgctctcc	1020
tgttccgacc	ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	gaagcgtggc	1080

O25CIP7SEQ List.txt

gctttctcaa	tgtcacgct	gtaggtatct	cagttcgggtg	taggtcgttc	gctccaagct	1140
gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	1200
tcttgagtcc	aacccggtaa	gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	1260
gattagcaga	gcgaggtatg	taggcgggtgc	tacagagttc	ttgaagtgg	ggcctaacta	1320
cggctacact	agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	1380
aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	gtggtttttt	1440
tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	1500
ttctacgggg	tctgacgctc	agtggaacga	aaactcacgt	taagggattt	tggtcatgag	1560
attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	ttaaatcaat	1620
ctaaagtata	tatgagtaaa	cttggtctga	cagttaccaa	tgcttaatca	gtgaggcacc	1680
tatctcagcg	atctgtctat	ttcgttcatc	catagttgcc	tgactccccg	tcgtgtagat	1740
aactacgata	cgggagggct	taccatctgg	ccccagtgc	gcaatgatac	cgcgagaccc	1800
acgtccaccg	gctccagatt	tatcagcaat	aaaccagcca	gccggaagg	ccgagcgcag	1860
aagtggctct	gcaactttat	ccgcctccat	ccagtctatt	aattgttgcc	gggaagctag	1920
agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	gccattgcta	caggcatcgt	1980
gggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	ggttcccaac	gatcaaggcg	2040
agttacatga	tccccatgt	tgtgcaaaaa	agcggttagc	tccttcggtc	ctccgatcgt	2100
tgtcagaagt	aagttggccg	cagtgttatc	actcatgggt	atggcagcac	tgcataattc	2160
tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	ggtgagtact	caaccaagtc	2220
attctgagaa	tagtgatg	ggcgaccgag	ttgctcttgc	ccggcgtcaa	tacgggataa	2280
taccgcgcca	catagcagaa	ctttaaaagt	gtcatcatt	ggaaaacggt	cttcggggcg	2340
aaaactctca	aggatcttac	cgtgttgtag	atccagttcg	atgtaacca	ctcgtgcacc	2400
caactgatct	tcagcatctt	ttactttcac	cagcgtttct	gggtgagcaa	aaacaggaag	2460
gcaaaatgcc	gcaaaaaaag	gaataagggc	gacacggaaa	tggtgaatac	tcatactctt	2520
cctttttcaa	tattattgaa	gcatttatca	gggttattgt	ctcatgagcg	gatacatatt	2580
tgaatgtatt	tagaaaaata	aacaaatagg	ggttccgcgc	acatttcccc	gaaaagtgcc	2640
acctgacgcg	ccctgtagcg	gcgcattaag	cgcggcggg	gtggtgggta	cgcgcagcgt	2700
gaccgctaca	cttgccagcg	ccctagcgcc	cgtcctttc	gctttcttcc	cttcctttct	2760
cgccacgttc	gccggctttc	cccgtcaagc	tctaaatcgg	gggctccctt	tagggttccg	2820
atttagtgct	ttacggcacc	tcgaccccaa	aaaacttgat	tagggtgatg	gttcacgtag	2880
tgggccatcg	ccctgataga	cggtttttctg	ccctttgaag	ttggagtcca	cgttctttaa	2940

			025CIP7SEQ	List.txt		
tagtggactc	ttgttccaaa	ctggaacaac	actcaaccct	atctcgggtct	attctttttga	3000
tttataaggg	attttgccga	tttcggccta	ttggttaaaa	aatgagctga	tttaacaaaa	3060
atttaacgcg	aattttaaca	aaatattaac	gtttacaatt	tcccattcgc	cattcaggct	3120
gcgcaactgt	tgggaagggc	gacgggtgcg	ggcctcttcg	ctattacgcc	agcccaagct	3180
accatgataa	gtaagtaata	ttaagggtacg	ggaggtactt	ggagcggccg	caataaaata	3240
tctttatfff	cattacatct	gtgtgtttgt	tttttgtgtg	aatcgatagt	actaacatac	3300
gctctccatc	aaaacaaaa	gaaacaaaa	aaactagcaa	aataggctgt	ccccagtgca	3360
agtgcagggtg	ccagaacatt	tctctatcga	taggtaccga	gctcttacgc	gtgctagccc	3420
tcgagcagga	tctatacatt	gaatcaatat	tggcaattag	ccatattagt	cattgggttat	3480
atagcataaa	tcaatattgg	ctattggcca	ttgcatacgt	tgtatctata	tcataaatatg	3540
tacatttata	ttggctcatg	tccaatatga	ccgccatgtt	gacattgatt	attgactagt	3600
tattaatagt	aatcaattac	ggggtcatta	gttcatagcc	catatatgga	gttcgcggtt	3660
acataactta	cggtaaatgg	cccgcctggc	tgaccgcca	acgacccccg	cccattgacg	3720
tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	ctttccattg	acgtcaatgg	3780
gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	aagtgtatca	tatgccaaagt	3840
ccgcccccta	ttgacgtcaa	tgacggtaaa	tggcccgcct	ggcattatgc	ccagtacatg	3900
accttacggg	actttcctac	ttggcagtac	atctacgtat	tagtcacgc	tattaccatg	3960
gtgatgcgggt	tttggcagta	catcaatggg	cgtggatagc	ggtttgactc	acggggattt	4020
ccaagtctcc	accccatgga	cgtcaatggg	agtttgtttt	ggcaccaaaa	tcaacggggac	4080
tttccaaaat	gtcgtataca	ctccgcccc	ttgacgcaa	tgggcggtag	gcgtgtacgg	4140
tgggaggtct	atataagcag	agctcgttta	gtgaaccgtc	agatcgccctg	gagacgccat	4200
ccacgctgtt	ttgacctcca	tagaagacac	cgggaccgat	ccagcctccc	ctcgaagctc	4260
gactctaggg	gctcgagatc	tgcgatctaa	gtaagcttgc	atgcctgcag	gtcggccgcc	4320
acgaccggtg	ccgccaccat	cccctgacct	acgcccctga	cccctcaca	ggagacgacc	4380
ttccatgacc	gagtacaagc	ccacgggtgcg	cctcgccacc	cgcgacgacg	tccccggggc	4440
cgtacgcacc	ctcgccgcgc	cgttcgcgca	ctaccccgcc	acgcgccaca	ccgtcgacct	4500
ggaccgccac	atcgagcggg	tcaccgagct	gcaagaactc	ttcctcacgc	gcgtcgggct	4560
cgacatcggc	aagggtgtggg	tgcgggacga	cggcgccgcg	gtggcggtct	ggaccacgcc	4620
ggagagcgtc	gaagcggggg	cgggtgttcgc	cgagatcggc	ccgcgcatgg	ccgagttgag	4680
cggttccccg	ctggccgcgc	agcaacagat	ggaaggcctc	ctggcgccgc	accggcccaa	4740
ggagcccgcg	tggttcctgg	ccaccgtcgg	cgtctcgccc	gaccaccagg	gcaagggtct	4800
gggcagcgcc	gtcgtgctcc	ccggagtggg	ggcgcccgag	cgcgcggggg	tgcgcgcctt	4860

O25CIP/SEQ List.txt

cctggagacc	tccgcgcccc	gcaacctccc	cttctacgag	cggctcggct	tcaccgtcac	4920
cgccgacgtc	gaggtgcccc	aaggaccgcg	cacctggtgc	atgaccgcga	agcccgggtgc	4980
ctgacgcccc	ccccacgacc	cgcagcgccc	gaccgaaagg	agcgcacgac	cccatggctc	5040
cgaccgaagc	cgacccgggc	ggccccgccc	accccgccacc	cgcccccgag	gcccaccgac	5100
t						5101

<210> 5
 <211> 5043
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Pl asmid pCMV-pur-attP

<400> 5	
ctagagtcgg	ggcggccggc cgcttcgagc agacatgata agatacattg atgagtttgg 60
acaaaccaca	actagaatgc agtgaaaaaa atgcttttatt tgtgaaattt gtgatgctat 120
tgcttttatt	gtaaccatta taagctgcaa taaacaagtt aacaacaaca attgcattca 180
ttttatgttt	caggttcagg gggaggtgtg ggaggttttt taaagcaagt aaaacctcta 240
caaatgtggt	aaaatcgata aggatcaatt cggcttcgac tagtactgac ggacacaccg 300
aagccccggc	ggcaaccctc agcggatgcc ccggggcttc acgttttccc aggtcagaag 360
cggttttcgg	gagtagtgcc ccaactgggg taacctttga gttctctcag ttgggggctg 420
agggtcgccc	acatgacaca aggggttggt accgggggtgg acacgtacgc ggggtgcttac 480
gaccgtcagt	cgcgcgagcg cgactagtac aagccgaatt gatccgtcga ccgatgccct 540
tgagagcctt	caaccacgac agtccttcc ggtgggcgcg gggcatgact atcgctcgccg 600
cacttatgac	tgtcttcttt atcatgcaac tcgtaggaca ggtgccggca gcgctcttcc 660
gcttcctcgc	tcactgactc gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct 720
cactcaaagg	cggtaatacg gttatccaca gaatcagggg ataacgcagg aaagaacatg 780
tgagcaaaag	gccagcaaaa ggccagggaac cgtaaaaagg ccgcgttgct ggcgtttttc 840
cataggctcc	gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 900
aaccgcacag	gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct 960
cctgttccga	ccctgccgct taccggatac ctgtccgcct ttctcccttc gggaagcgtg 1020
gcgctttctc	aatgctcacg ctgtaggtat ctcagttcgg tgtaggtegt tcgctccaag 1080
ctgggctgtg	tgcacgaacc ccccgttcag ccgaccgct gcgccttata cggtaactat 1140
cgtcttgagt	ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac 1200
aggattagca	gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 1260

			025CIP7SEQ	List.tXt			
tacggctaca	ctagaaggac	agtatttggg	atctgcgctc	tgctgaagcc	agttaccttc		1320
ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cggtaggtttt		1380
tttgtttgca	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc		1440
ttttctacgg	ggctctgacgc	tcagtggaac	gaaaactcac	gttaagggat	tttggtcatg		1500
agattatcaa	aaaggatctt	cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca		1560
atctaaagta	tatatgagta	aacttggtct	gacagttacc	aatgcttaat	cagtgaggca		1620
cctatctcag	cgatctgtct	atttcgttca	tccatagttg	cctgactccc	cgctcgttag		1680
ataactacga	tacggggagg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac		1740
ccacgctcac	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc		1800
agaagtgggc	ctgcaacttt	atccgcctcc	atccagtcta	ttaattgttg	ccgggaagct		1860
agagtaagta	gttcgccagt	taatagtttg	cgcaacgttg	ttgccattgc	tacaggcatc		1920
gtggtgtcac	gctcgtcggt	tggtatggct	tcattcagct	ccgggtccca	acgatcaagg		1980
cgagttacat	gatcccccat	gttgtgcaaa	aaagcggtta	gtccttcggg	tcctccgatc		2040
gttgtcagaa	gtaagttggc	cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat		2100
tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	ctgggtgagta	ctcaaccaag		2160
tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgctc	aatacgggat		2220
aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg		2280
cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca		2340
cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga		2400
aggcaaaatg	ccgcaaaaaa	gggaataagg	gcgacacgga	aatggtgaat	actcatactc		2460
ttcctttttc	aatattattg	aagcatttat	caggggttatt	gtctcatgag	cggatacata		2520
tttgaatgta	tttagaaaaa	taaacaaata	ggggttccgc	gcacatttcc	ccgaaaagtg		2580
ccacctgacg	cgcctgtag	cggcgcatta	agcgcggcgg	gtgtgggtgg	tacgcgcagc		2640
gtgaccgcta	cacttgccag	cgcctagcgc	cccgtctctt	tcgttttctt	cccttccttt		2700
ctcgccacgt	tcgcgggctt	tccccgtcaa	gctctaaatc	gggggctccc	tttagggttc		2760
cgatttagtg	ctttacggca	cctcgacccc	aaaaaacttg	attaggggtga	tggttcacgt		2820
agtgggccat	cgcctgata	gacggttttt	cgccctttga	cgttggagtc	cacgttcttt		2880
aatagtggac	tcttgttcca	aactggaaca	acactcaacc	ctatctcggt	ctattctttt		2940
gatttataag	ggattttgcc	gatttcggcc	tattggttaa	aaaatgagct	gatttaacaa		3000
aaatttaacg	cgaattttta	caaaatatta	acgtttacaa	tttccattc	gccattcagg		3060
ctgcgcaact	gttgggaagg	gcgatcggtg	cgggcctctt	cgctattacg	ccagcccaag		3120
ctaccatgat	aagtaagtaa	tattaaggta	cgggaggtac	ttggagcggc	cgcaataaaa		3180

025CIP7SEQ List.tXt

tatctttatt	ttcattacat	ctgtgtgttg	gttttttgtg	tgaatcgata	gtactaacat	3240
acgctctcca	tcaaaacaaa	acgaaacaaa	acaaactagc	aaaataggct	gtccccagtg	3300
caagtgcagg	tgccagaaca	tttctctatc	gataggtacc	gagctcttac	gcgtgctagc	3360
cctcgagcag	gatctataca	ttgaatcaat	attggcaatt	agccatatta	gtcattgggt	3420
atatagcata	aatcaatatt	ggctattggc	cattgcatac	gttgatatcta	tatcataata	3480
tgtacattta	tattggctca	tgtccaatat	gaccgccatg	ttgacattga	ttattgacta	3540
gttattaata	gtaatcaatt	acgggggtcat	tagttcatag	cccatatatg	gagttccgog	3600
ttacataact	tacggtaaat	ggccccgectg	gctgaccgcc	caacgacccc	cgcgccattga	3660
cgtcaataat	gacgtatggt	cccatagtaa	cgccaatagg	gactttccat	tgacgtcaat	3720
gggtggagta	tttacggtaa	actgcccact	tggcagtaca	tcaagtgtat	catatgccaa	3780
gtccgcccc	tattgacgtc	aatgacggta	aatggccgc	ctggcattat	gccagtgaca	3840
tgaccttacg	ggactttcct	acttggcagt	acatctacgt	attagtcac	gctattacca	3900
tggatgatgcg	gttttggcag	tacatcaatg	ggcgtggata	gcggtttgac	tcacggggat	3960
ttccaagtct	ccaccccat	gacgtcaatg	ggagtttgtt	ttggcaccaa	aatcaacggg	4020
actttccaaa	atgtcgtaac	aactccgcc	cattgacgca	aatgggcggt	aggcgtgtac	4080
ggtgggaggt	ctatataagc	agagctcggt	tagtgaaccg	tcagatcgcc	tggagacgcc	4140
atccacgctg	ttttgacctc	catagaagac	accgggaccg	atccagcctc	ccctcgaagc	4200
tcgactctag	gggctcgaga	tctgcgatct	aagtaagctt	gcatgcctgc	aggtcggccg	4260
ccacgaccgg	tgccgccacc	atcccctgac	ccacgcccct	gacccctcac	aaggagacga	4320
ccttccatga	cagagtacaa	gcccacgggtg	cgctcgcga	cccgcgacga	cgtcccccg	4380
gccgtacgca	ccctcgccgc	cgcgttcgcc	gactaccccg	ccacgcgcca	caccgtcgac	4440
ccggaccgcc	acatcgagcg	ggtcaccgag	ctgcaagaac	tcttctctac	gcgcgtcggg	4500
ctcgacatcg	gcaaggtgtg	ggtcgcggac	gacggcgccg	cgggtggcggt	ctggaccacg	4560
ccggagagcg	tcgaagcggg	ggcgggtgtc	gccgagatcg	gcccgcgcac	ggccgagttg	4620
agcggttccc	ggctggccgc	gcagcaacag	atggaaggcc	tcctggcgcc	gcaccggccc	4680
aaggagcccc	cgtggttcc	ggccaccgtc	ggcgtctcgc	ccgaccacca	gggcaagggt	4740
ctgggcagcg	ccgtcgtgct	ccccggagtg	gaggcgccg	agcgcgcggg	ggtgcccgcc	4800
ttcctggaga	cctccgcgcc	ccgcaacctc	cccttctacg	agcggctcgg	cttcaccgtc	4860
accgccgacg	tcgaggtgcc	cgaaggaccg	cgcacctggt	gcatgaccgc	caagcccggt	4920
gcctgacgcc	cgcgccacga	ccgcgagcgc	ccgaccgaaa	ggagcgcacg	accccatggc	4980
tccgaccgaa	gccgaccggg	gcggccccgc	cgaccccgca	ccgccccccg	aggcccaccg	5040

act 025CIP7SEQ List.tXt 5043

<210> 6
 <211> 5041
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Plasmid pCMV-EGFP-attB

<400> 6
 ctagagtcgg ggcggccggc cgcttcgagc agacatgata agatacattg atgagtttgg 60
 acaaaccaca actagaatgc agtgaaaaaa atgcttttatt tgtgaaattt gtgatgctat 120
 tgcttttattt gtaaccatta taagctgcaa taaacaagtt aacaacaaca attgcattca 180
 ttttatgttt cagggttcagg gggaggtgtg ggaggttttt taaagcaagt aaaacctcta 240
 caaatgtggg aaaatcgata aggatcaatt cggcttcagg taccgtcgac gatgtaggtc 300
 acggtctcga agccgcgggtg cgggtgccag ggcgtgccct tgggctcccc ggcgcgtac 360
 tccacctcac ccatctgggc catcatgatg aacgggtcga ggtggcggta gttgatcccc 420
 gcgaacgcgc ggcgcaccgg gaagccctcg cctcgaaac cgctgggcgc ggtggtcacg 480
 gtgagcacgg gacgtgcgac ggcgtcggcg ggtgcggata cgcggggcag cgtcagcggg 540
 ttctcgacgg tcacggcggg catgtcgaca gccgaattga tccgtcgacc gatgcccttg 600
 agagccttca acccagtcag ctccctccgg tgggcgcggg gcatgactat cgtcgccgca 660
 cttatgactg tcttctttat catgcaactc gtaggacagg tgccggcagc gctcttccgc 720
 ttctctcgctc actgactcgc tgcgctcggc cgttcggctg cggcgagcgg tatcagctca 780
 ctcaaaggcg gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg 840
 agcaaaaggc cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgtttttcca 900
 taggctccgc cccctgacg agcatcacia aaatcgacgc tcaagtcaga ggtggcgaaa 960
 cccgacagga ctataaagat accaggcggt tccccctgga agctccctcg tgcgctctcc 1020
 tgttccgacc ctgcgcgtta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc 1080
 gctttctcaa tgctcacgct gtaggtatct cagttcgggtg taggtcgttc gctccaagct 1140
 gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg 1200
 tcttgagtcc aacccggtta gacacgactt atcgccactg gcagcagcca ctggtaacag 1260
 gattagcaga gogaggtatg taggcggtgc tacagagttc ttgaagtggg ggcctaacta 1320
 cggctacact agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg 1380
 aaaaagagtt ggtagctctt gatccggcaa acaaaccacc gctggtagcg gtgggttttt 1440
 tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt 1500
 ttctacgggg tctgacgctc agtggaacga aaactcacgt taagggattt tggatcatgag 1560

025CIP7SEQ List.txt

attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	ttaaataaat	1620
ctaaagtata	tatgagtaaa	cttgggtctga	cagttaccaa	tgcttaataca	gtgaggcacc	1680
tatctcagcg	atctgtctat	ttcgttcac	catagttgcc	tgactccccg	tcgtgtagat	1740
aactacgata	cgggagggct	taccatctgg	ccccagtgc	gcaatgatac	cgcgagaccc	1800
acgctcaccg	gctccagatt	tatcagcaat	aaaccagcca	gccggaagg	ccgagcgcag	1860
aagtggctct	gcaactttat	cgcctccat	ccagtctatt	aattgttgcc	gggaagctag	1920
agtaagtagt	tcgccagtta	atagtttg	caacgttg	gccattgcta	caggcatcgt	1980
gggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	ggttcccaac	gatcaaggcg	2040
agttacatga	ttccccatgt	tgtgcaaaaa	agcggtttag	tccttcggtc	ctccgatcgt	2100
tgtcagaagt	aagttggccg	cagtgttatc	actcatgggt	atggcagcac	tgcataattc	2160
tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	ggtagagtact	caaccaagtc	2220
attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	ccggcgtaaa	tacgggataa	2280
taccgcgcca	catagcagaa	ctttaaaagt	gctcatcatt	ggaaaacgtt	cttcggggcg	2340
aaaactctca	aggatcttac	cgtgtgtgag	atccagttcg	atgtaacca	ctcgtgcacc	2400
caactgatct	tcagcatctt	ttactttcac	cagcgtttct	gggtgagcaa	aaacaggaag	2460
gcaaaatgcc	gcaaaaaagg	gaataagggc	gacacggaaa	tggtgaatac	tcatactctt	2520
cctttttcaa	tattattgaa	gcatttatca	gggttattgt	ctcatgagcg	gatacatatt	2580
tgaatgtatt	tagaaaaata	aacaaatagg	ggttccgcgc	acattttccc	gaaaagtgcc	2640
acctgacgcg	ccctgtagcg	gcgcattaag	cgcggcgggt	gtgggtggta	cgcgcagcgt	2700
gaccgctaca	cttgccagcg	ccctagcgcc	cgtctctttc	gctttcttcc	cttcctttct	2760
cgccacgttc	gcgggctttc	ccgtcaagc	tctaaatcgg	gggtctccct	taggggtccg	2820
atttagtgct	ttacggcacc	tcgaccccaa	aaaacttgat	taggggtgatg	gttcacgtag	2880
tgggccatcg	ccctgataga	cggtttttcg	cccttgacg	ttggagtcca	cgttctttaa	2940
tagtggactc	ttgttccaaa	ctggaacaac	actcaaccct	atctcggtct	attcttttga	3000
tttataaggg	attttgccga	tttcggccta	ttgggttaaaa	aatgagctga	tttaacaaaa	3060
atttaacgcg	aattttaaca	aaatattaac	gtttacaatt	tcccatctgc	cattcaggct	3120
gcgcaactgt	tgggaagggc	gatcggtgcg	ggcctcttcg	ctattacgcc	agcccaagct	3180
accatgataa	gtaagtaata	ttaaggtagc	ggaggtactt	ggagcggccg	caataaaata	3240
tctttatatt	cattacatct	gtgtgttggt	ttttgtgtg	aatcgatagt	actaacatac	3300
gctctccatc	aaaacaaaac	gaaacaaaac	aaactagcaa	aataggctgt	ccccagtgca	3360
agtgcaggtg	ccagaacatt	tctctatcga	taggtaccga	gctcttacgc	gtgctagccc	3420

025CIP/SEQ List.tXt

tcgagcagga	tctatacatt	gaatcaatat	tggaatttag	ccatattagt	cattgggttat	3480
atagcataaa	tcaatattgg	ctattggcca	ttgcatacgt	tgtatctata	tcataaatatg	3540
tacattttata	ttggctcatg	tccaatatga	ccgccatggt	gacattgatt	attgactagt	3600
tattaatagt	aatcaattac	ggggtcatta	gttcatagcc	catatatgga	gttccgcgtt	3660
acataactta	cggtaaattg	ccgcctggc	tgaccgccc	acgacccccg	cccattgacg	3720
tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	ctttccattg	acgtcaatgg	3780
gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	aagtgtatca	tatgccaaagt	3840
ccgcccccta	ttgacgtcaa	tgacggtaaa	tgggccgcct	ggcattatgc	ccagtacatg	3900
accttacggg	actttcctac	ttggcagtac	atctacgtat	tagtcacgc	tattaccatg	3960
gtgatgcggt	tttggcagta	catcaatggg	cgtggatagc	ggtttgactc	acggggattt	4020
ccaagtctcc	acccattga	cgtcaatggg	agtttgtttt	ggcaccaaaa	tcaacgggac	4080
tttccaaaat	gtcgtaacaa	ctccgcccc	ttgacgcaaa	tgggcggtag	gcgtgtacgg	4140
tgggaggtct	atataagcag	agctcgttta	gtgaaccgtc	agatcgctg	gagacgccat	4200
ccacgtgtgt	ttgacctcca	tagaagacac	cgggaccgat	ccagcctccc	ctcgaagctc	4260
gactctaggg	gctcgagatc	cccgggtacc	ggtcgccacc	atggtgagca	agggcgagga	4320
gctgttcacc	ggggtggtgc	ccatcctggg	cgagctggac	ggcgacgtaa	acggccacaa	4380
gttcagcgtg	tccggcgagg	gcgagggcga	tgccacctac	ggcaagctga	ccctgaagtt	4440
catctgcacc	accggcaagc	tgcccggtgc	ctggcccacc	ctcgtgacca	ccctgacct	4500
cggcgtgcag	tgcttcagcc	gctaccccga	ccacatgaag	cagcacgact	tcttcaagtc	4560
cgccatgccc	gaaggctacg	tccaggagcg	caccatcttc	ttcaaggacg	acggcaacta	4620
caagacccgc	gccgaggtga	agttcgaggg	cgacaccctg	gtgaaccgca	tcgagctgaa	4680
gggcatcgac	ttcaaggagg	acggcaacat	cctggggcac	aagctggagt	acaactacaa	4740
cagccacaa	gtctatatca	tggccgacaa	gcagaagaac	ggcatcaagg	tgaacttcaa	4800
gatccgccac	aacatcgagg	acggcagcgt	gcagctcgcc	gaccactacc	agcagaacac	4860
ccccatcggc	gacggccccg	tgctgctgcc	cgacaaccac	tacctgagca	cccagtcgcg	4920
cctgagcaaa	gaccccaacg	agaagcgcca	tcacatggtc	ctgctggagt	tcgtgaccgc	4980
cgcggggatc	actctcggca	tggacgagct	gtacaagtaa	agcggccgct	cgagcatgca	5040
t						5041

<210> 7
 <211> 18116
 <212> DNA
 <213> Artificial Sequence
 <220>

O25CIP7SEQ List.tXt
 <223> Plasmid pl2 .01ys-LSPIPNMM-CMV-pur-attB

<400> 7

```

gggctgcagg aattcgattg ccgccttctt tgatattcac tctgttgat ttcattctctt      60
cttgccgatg aaaggatata acagtctgta taacagtctg tgaggaaata cttgggtatctt    120
cttctgatca gtgtttttat aagtaatgtt gaatattgga taaggctgtg tgtcctttgt      180
cttgggagac aaagcccaca gcagggtggtg gttgggggtgg tggcagctca gtgacaggag    240
aggttttttt gcctgttttt tttttttttt ttttttttaa gtaagggtgtt cttttttctt    300
agtaaatttt ctactggact gtatgttttg acaggtcaga aacatttctt caaaagaaga    360
accttttgga aactgtacag cccttttctt tcattccctt tttgctttct gtgccaatgc    420
ctttggttct gattgcatta tggaaaacgt tgatcggaac ttgaggtttt tatttatagt    480
gtggcttgaa agcttgata gctgttggtt cacgagatac cttattaagt ttaggccagc    540
ttgatgcttt attttttccc tttgaagtag tgagcgttct ctggtttttt tcctttgaaa    600
ctggtgaggc ttagattttt ctaatgggat tttttacctg atgatctagt tgcataccca    660
aatgcttgta aatgttttcc tagttaacat gttgataact tcggatttac atgttgata      720
tacttgatcat ctgtgtttct agtaaaaata tatggcattt atagaaatac gtaattcctg    780
atttcctttt tttttatctc tatgctctgt gtgtacaggc caaacagact tcactcctat    840
ttttatttat agaattttat atgcagtctg tcgttggttc ttgtgttgta aggatacagc    900
cttaaatttc ctagagcgat gctcagtaag gcgggttggt acatgggttc aaatgtaaaa    960
cgggcacggt tggctgctgc cttcccgaga tccaggacac taaactgctt ctgcactgag   1020
gtataaatcg cttcagatcc cagggaagtg cagatccacg tgcataattct taaagaagaa   1080
tgaatacttt ctaaaatatt ttggcatagg aagcaagctg catggatttg tttgggactt   1140
aaattatttt ggtaacggag tgcatagggt ttaaacacag ttgcagcatg ctaacgagtc   1200
acagcgttta tgcagaagtg atgcctggat gcctgttgca gctgtttacg gcactgcctt   1260
gcagtgagca ttgcagatag ggggtgggtg ctttgtgtcg tgttcccaca cgctgccaca   1320
cagccacctc ccggaacaca tctcacctgc tgggtacttt tcaaaccatc ttagcagtag   1380
tagatgagtt actatgaaac agagaagttc ctcagttgga tattctcatg ggatgtcttt   1440
tttcccatgt tgggcaaagt atgataaagc atctctatct gtaaattatg cacttgtagt   1500
ttcctgaatc ctttctatag caccacttat tgcagcaggt gtaggctctg gtgtggcctg   1560
tgtctgtgct tcaatctttt aaagcttctt tggaaataca ctgacttgat tgaagtctct   1620
tgaagatagt aaacagtact tacctttgat cccaatgaaa tcgagcattt cagttgtaaa   1680
agaattccgc ctattcatac catgtaatgt aattttacac cccagtgct gacactttgg   1740
aatatattca agtaatagac tttggcctca cctcttgttg tactgtatct tgtaatagaa   1800

```

025CIP7SEQ List.txt

aataatttttaa	actgtgcata	tgattattac	attatgaaag	agacattctg	ctgatcttca	1860
aatgtaagaa	aatgaggagt	gcgtgtgctt	ttataaatac	aagtgattgc	aaattagtgc	1920
aggtgtcctt	aaaaaaaaaa	aaaaaaagta	atataaaaag	gaccaggtgt	tttacaagtg	1980
aaatacattc	ctatttggtg	aacagttaca	tttttatgaa	gattaccage	gctgctgact	2040
ttctaaacat	aaggctgtat	tgtcttcctg	taccattgca	tttcctcatt	cccaatttgc	2100
acaaggatgt	ctgggtaaac	tattcaagaa	atggctttga	aatacagcat	gggagcttgt	2160
ctgagttgga	atgcagagtt	gcactgcaaa	atgtcaggaa	atggatgtct	ctcagaatgc	2220
ccaactccaa	aggattttat	atgtgtatat	agtaagcagt	ttcctgattc	cagcaggcca	2280
aagagtctgc	tgaatgttgt	gttgccggag	acctgtatct	ctcaacaagg	taagatggta	2340
tcctagcaac	tgcggatttt	aatacatttt	cagcagaagt	acttagttaa	tctctacctt	2400
tagggatcgt	ttcatcattt	ttagatgtta	tacttgaaat	actgcataac	ttttagcttt	2460
catgggttcc	tttttttcag	cctttaggag	actgttaagc	aatttgctgt	ccaacttttg	2520
tgttggctct	aaactgcaat	agtagtttac	cttgtattga	agaaataaag	accattttta	2580
tattaaaaaa	tacttttgtc	tgtcttcatt	ttgacttgtc	tgatatacct	gcagtgccca	2640
ttatgtcagt	tctgtcagat	attcagacat	caaaacttaa	cgtgagctca	gtggagttac	2700
agctgcgggt	ttgatgctgt	tattatttct	gaaactagaa	atgatgttgt	cttcactctgc	2760
tcatacaaca	cttcacgcag	agtgtgaagg	tagtgagaaa	tgcatatact	tattgatact	2820
tttttaaaagt	caacttttta	tcagattttt	ttttcatttg	gaaatatatt	gttttctaga	2880
ctgcatagct	tctgaatctg	aatgacagtc	tgattggcat	gaagaagcac	agcactcttc	2940
atcttactta	aacttcattt	tggaatgaag	gaagttaagc	aagggcacag	gtccatgaaa	3000
tagagacagt	gcgctcagga	gaaagtgaac	ctggatttct	ttggctagtg	ttctaaatct	3060
gtagtgagga	aagtaacacc	cgattccttg	aaagggctcc	agctttaatg	cttccaaatt	3120
gaaggtggca	ggcaacttgg	ccactgggta	tttactgcat	tatgtctcag	tttcgcagct	3180
aacctggctt	ctccactatt	gagcatggac	tatagcctgg	cttcagaggc	caggtgaagg	3240
ttgggatggg	tgggaaggagt	gctgggctgt	ggctgggggg	actgtgggga	ctccaagctg	3300
agcttggggg	gggcagcaca	gggaaaagtg	tgggtaacta	tttttaagta	ctgtgttgca	3360
aacgtctcat	ctgcaaatac	gtagggtgtg	tactctcgaa	gattaacagt	gtgggttcag	3420
taatatatgg	atgaattcac	agtggaagca	ttcaagggtg	gatcatctaa	cgacaccaga	3480
tcataaagct	atgattggaa	gcggtatcag	aagagcgagg	aaggtaagca	gtcttcatat	3540
gttttccctc	cacgtaaagc	agtctgggaa	agtagcaccc	cttgagcaga	gacaaggaaa	3600
taattcagga	gcatgtgcta	ggagaacttt	cttgcgtgaat	tctacttgca	agagctttga	3660
tgcttggctt	ctggtgcctt	ctgcagcacc	tgcaaggccc	agagcctgtg	gtgagctgga	3720

O25CIP7SEQ List.tXt

gggaaagatt	ctgctcaagt	ccaagcttca	gcaggtcatt	gtctttgctt	cttccccag	3780
cactgtgcag	cagagtggaa	ctgatgtcga	agcctcctgt	ccactacctg	ttgctgcagg	3840
cagactgctc	tcagaaaaag	agagctaact	ctatgccata	gtctgaaggt	aaaatggggt	3900
ttaaaaaaga	aaacacaaag	gcaaaaccgg	ctgccccatg	agaagaaagc	agtggtaaac	3960
atggtagaaa	aggtgcagaa	gccccaggc	agtgtgacag	gcccctcctg	ccacctagag	4020
gcgggaacaa	gcttccctgc	ctagggctct	gcccgcgaag	tgcgtgtttc	tttgggtgggt	4080
tttgtttggc	gtttgggttt	gagatttaga	cacaagggaa	gcctgaaagg	aggtgttggg	4140
cactattttg	gtttgtaaag	cctgtacttc	aaatatatat	tttgtgaggg	agtgtagcga	4200
attggccaat	ttaaaataaa	gttgcaagag	attgaaggct	gagtagttga	gagggttaaca	4260
cgtttaatga	gatcttctga	aactactgct	tctaaacact	tgtttgagtg	gtgagacctt	4320
ggataggtga	gtgctcttgt	tacatgtctg	atgcacttgc	ttgtcctttt	ccatccacat	4380
ccatgcattc	cacatccacg	catttgtcac	ttatcccata	tctgtcatat	ctgacatacc	4440
tgtctcttcg	tcacttggtc	agaagaaaca	gatgtgataa	tcccagccg	ccccaaagttt.	4500
gagaagatgg	cagttgcttc	tttccctttt	tctgtctaag	taaggatttt	ctcctggcct	4560
tgacacctca	cgaaatagtc	ttcctgcctt	acattctggg	cattatttca	aatatctttg	4620
gagtgcgctg	ctctcaagtt	tgtgtcttcc	tactcttaga	gtgaatgctc	ttagagtga	4680
agagaaggaa	gagaagatgt	tggccgcagt	tctctgatga	acacacctct	gaataatggc	4740
caaaggtggg	tgggtttctc	tgaggaacgg	gcagcgtttg	cctctgaaag	caaggagctc	4800
tgcgagattg	cagttatttt	gcaactgatg	gtggaactgg	tgcttaaagc	agattcccta	4860
ggttccctgc	tacttctttt	ccttcttggc	agtcagttta	tttctgacag	acaaacagcc	4920
acccccactg	caggcttaga	aagtatgtgg	ctctgcctgg	gtgtgttaca	gctctgccct	4980
ggtgaaaggg	gattaaaacg	ggcaccattc	atcccaaaca	ggatcctcat	tcatggatca	5040
agctgtaagg	aacttgggct	ccaacctcaa	aacattaatt	ggagtacgaa	tgtaatataa	5100
actgcattct	cgcattccta	agtcatttag	tctggactct	gcagcatgta	ggtcggcagc	5160
tcccactttc	tcaaagacca	ctgatggagg	agtagtaaaa	atggagaccg	attcagaaca	5220
accaacggag	tgttgccgaa	gaaactgatg	gaaataatgc	atgaattgtg	tgggtggacat	5280
ttttttttaa	tacataaact	acttcaaagt	aggtcggaga	aggtcagtgt	tttattagca	5340
gccataaaac	caggtgagcg	agtaccattt	ttctctacaa	gaaaaacgat	tctgagctct	5400
gcgtaagtat	aagttctcca	tagcggctga	agctcccccc	tggctgcctg	ccatctcagc	5460
tggagtgcag	tgccatttcc	ttggggtttc	tctcacagca	gtaatgggac	aatacttcac	5520
aaaaattctt	tcttttctctg	tcatgtggga	tccctactgt	gccctcctgg	ttttacgtta	5580

			O25CIP7SEQ	List.tXt		
ccccctgact	gttccattca	gcggttttgg	aagagaaaaa	gaattttggaa	ataaaacatg	5640
tctacgttat	cacctcctcc	agcatttttg	tttttaatta	tgtcaataac	tggettagat	5700
ttggaaatga	gagggggttg	ggtgtattac	cgaggaacaa	aggaaggctt	atataaactc	5760
aagtctttta	tttagagaac	tggcaagctg	tcaaaaacaa	aaaggcctta	ccaccaaatt	5820
aagtgaatag	ccgctatagc	cagcagggcc	agcacgaggg	atggtgcact	gctggcacta	5880
tgccacggcc	tgcttgtgac	tctgagagca	actgcttttg	aaatgacagc	acttggtgca	5940
atttcctttg	tttcagaatg	cgtagagcgt	gtgcttggcg	acagtttttc	tagttaggcc	6000
acttcttttt	tccttctctc	ctcattctcc	taagcatgtc	tccatgctgg	taatcccagt	6060
caagtgaacg	ttcaaacaat	gaatccatca	ctgtaggatt	ctcgtggtga	tcaaactctt	6120
gtgtgaggtc	tataaaatat	ggaagcttat	ttatttttcg	ttcttccata	tcagtcttct	6180
ctatgacaat	tcacatccac	cacagcaa	taaagggtga	ggaggctggt	gggatgaaga	6240
gggtcttcta	gctttacgtt	cttccttgca	aggccacagg	aaaatgctga	gagctgtaga	6300
atacagcctg	gggtaagaag	ttcagtctcc	tgctgggaca	gctaaccgca	tcttataacc	6360
ccttctgaga	ctcatcttag	gaccaa	ggtctatctg	gggtttttgt	tcctgctgtt	6420
cctcctggaa	ggctatctca	ctatttctact	gtctccacgg	ttacaaacca	aagatacagc	6480
ctgaattttt	tctaggccac	attacataaa	tttgacctgg	taccaatatt	gttctctata	6540
tagttatttc	cttccccact	gtgtttaacc	ccttaaggca	ttcagaacaa	ctagaatcat	6600
agaatggttt	ggattggaag	gggccttaaa	catcatccat	ttccaaccct	ctgccatggg	6660
ctgcttgcca	cccactggct	caggctgccc	agggccccat	ccagcctggc	cttgagcacc	6720
tccagggatg	gggcacccac	agcttctctg	ggcagcctgt	gccaacacct	caccactctc	6780
tgggtaaaga	attctctttt	aacatcta	ctaaatctct	tctcttttag	tttaaagcca	6840
ttcctctttt	tcccgttgct	atctgtccaa	gaaatgtgta	ttggtctccc	tcctgcttat	6900
aagcaggaag	tactggaag	ctgcagttag	gtctccccac	agccttctct	tctccaggct	6960
gaacaagccc	agctccttca	gcctgtcttc	gtaggagatc	atcttagtgg	ccctcctctg	7020
gacccattcc	aacagttcca	cggctttctt	gtggagcccc	aggtctggat	gcagtacttc	7080
agatggggcc	ttacaaaggc	agagcagatg	gggacaatcg	cttaccctc	cctgctggct	7140
gccccgtttt	tgatgcagcc	cagggtagctg	ttggcctttc	aggtcccag	acccttgct	7200
gatttgtgtc	aagcttttca	tccaccagaa	cccacgcttc	ctggttaata	cttctgcctt	7260
cacttctgta	agcttgtttc	aggagacttc	cattcttttag	gacagactgt	gttacaccta	7320
cctgccctat	tcttgcatat	atacatttca	gttcatgttt	cctgtaacag	gacagaatat	7380
gtattcctct	aacaaaaata	catgcagaat	tcctagtgcc	atctcagtag	ggttttcatg	7440
gcagtattag	cacatagtca	atttgctgca	agtaccttcc	aagctgcggc	ctcccataaa	7500

025CIP7SEQ List.tXt

tcctgtat	ttt	gggatcag	tt	accttttggg	gtaagctttt	gtatctgcag	agacctggg	7560
ggttctgatg	tgttcagct	ctgctctgtt	ctgactgcac	cattttctag	atcaccag	7620		
tgttcctgta	caacttcctt	gtcctccatc	ctttccagc	ttgtatcttt	gacaaataca	7680		
ggcctat	ttt	tgtgtttgct	tcagcagcca	tttaattctt	cagtgtcatc	ttgttctgtt	7740	
gatgccactg	gaacaggatt	ttcagcagtc	ttgcaaagaa	catctagctg	aaaactttct	7800		
gccattcaat	attcttacca	gttcttcttg	tttgagggtga	gccataaatt	actagaactt	7860		
cgtcactgac	aagtttatgc	attttattac	ttctattatg	tacttacttt	gacataacac	7920		
agacacgcac	atattttgct	gggatttcca	cagtgtctct	gtgtccttca	catggtttta	7980		
ctgtcatact	tccgttataa	ccttggcaat	ctgccagct	gcccatcaca	agaaaagaga	8040		
ttcctttttt	attacttctc	ttcagccaat	aaacaaaatg	tgagaagccc	aaacaagaac	8100		
ttgtggggca	ggctgccatc	aaggagaga	cagctgaagg	gttgtgtagc	tcaatagaat	8160		
taagaaataa	taaagctgtg	tcagacagtt	ttgcctgatt	tatacaggca	cgccccaagc	8220		
cagagaggct	gtctgccaa	gccaccttgc	agtccttgg	ttgtaagata	agtcataggt	8280		
aacttttctg	gtgaattg	tgagaaatca	tgatggcag	tcttgctgtt	tactatggta	8340		
agatgctaaa	ataggagaca	gcaaagtaac	acttgctgct	gtaggtgctc	tgctatccag	8400		
acagcgatgg	cactgcaca	ccaagatgag	ggatgctccc	agctgacgga	tgctggggca	8460		
gtaacagtgg	gtcccatgct	gcctgctcat	tagcatcacc	tcagccctca	ccagcccatc	8520		
agaaggatca	tcccaagctg	aggaaagttg	ctcatcttct	tcacatcacc	aaacctttgg	8580		
cctgactgat	gcctcccgg	tgtttaa	atg	tggtcactga	catctttatt	tttctatgat	8640	
ttcaagtcag	aacctccgg	tcaggaggga	acacatagtg	ggaatgtacc	ctcagctcca	8700		
aggccagatc	ttccttcaat	gatcatgcat	gctacttagg	aagggtgtgtg	tgtgtgaatg	8760		
tagaattgcc	tttgttat	tttcttctg	ctgtcaggaa	catttttgaat	accagagaaa	8820		
aagaaaagt	g	ctcttcttgg	catgggagga	gttgtcacac	ttgcaaaata	aaggatgcag	8880	
tcccaa	atgt	tcataatctc	agggctgaa	ggaggatcag	aaactgtgta	tacaatttca	8940	
ggcttctctg	aatgcagctt	ttgaaagctg	ttcctggccg	aggcagtact	agtcagaacc	9000		
ctcgga	aaca	ggaacaaatg	tcttcaaggt	gcagcaggag	gaaacacctt	gcccatcatg	9060	
aaagtgaata	accactgccg	ctgaaggaat	ccagctcctg	tttgagcagg	tgctgcacac	9120		
tcccacactg	aaacaacagt	tcatttttat	aggacttcca	ggaaggatct	tcttcttaag	9180		
cttcttaatt	atggtacatc	tccagttggc	agatgactat	gactactgac	aggagaatga	9240		
ggaactagct	gggaatat	ctgtttgacc	accatggagt	caccat	ttt	actggta	9300	
tttgga	aaata	ataattctga	attgcaaagc	aggagttagc	gaagatcttc	atttcttcca	9360	

			025CIP7SEQ	List.tXt			
tggttggtgac	agcacagttc	tggttatgaa	agtctgctta	caaggaagag	gataaaaatc		9420
atagggataa	taaatctaag	tttgaagaca	atgaggtttt	agctgcattt	gacatgaaga		9480
aattgagacc	tctactggat	agctatggta	tttacgtgtc	tttttgctta	gttacttatt		9540
gaccccagct	gaggtcaagt	atgaactcag	gtctctcggg	ctactggcat	ggattgatta		9600
catacaactg	taattttagc	agtgatttag	ggtttatgag	tacttttgca	gtaaatcata		9660
gggttagtaa	tgttaatctc	agggaaaaaa	aaaaaaagcc	aaccctgaca	gacatcccag		9720
ctcaggtgga	aatcaaggat	cacagctcag	tgcggtccca	gagaacacag	ggactcttct		9780
cttaggacct	ttatgtacag	ggcctcaaga	taactgatgt	tagtcagaag	actttccatt		9840
ctggccacag	ttcagctgag	gcaatcctgg	aattttctct	ccgtgcaca	gttccagtca		9900
tcccagtttg	tacagttctg	gcactttttg	ggtcaggccg	tgatccaagg	agcagaagtt		9960
ccagctatgg	tcagggagtg	cctgaccgtc	ccaactcact	gcactcaaac	aaaggcgaaa		10020
ccacaagagt	ggcttttggt	gaaattgcag	tgtggcccag	aggggctgca	ccagtactgg		10080
attgaccacg	aggcaacatt	aatcctcagc	aagtgcattt	tgcagccatt	aaattgaact		10140
aactgatact	acaatgcaat	cagtatcaac	aagtggtttg	gcttggaaga	tggagtctag		10200
gggctctaca	ggagtagcta	ctctctaatt	gagttgcatt	ttgaagcagg	acactgtgaa		10260
aagctggcct	cctaaagagg	ctgctaaaca	ttaggggtcaa	ttttccagtg	cactttctga		10320
agtgtctgca	gttccccatg	caaagctgcc	caaacatagc	acttccaatt	gaatacaatt		10380
atatgcaggc	gtactgcttc	ttgccagcac	tgtccttctc	aaatgaactc	aacaaacaat		10440
ttcaaagtct	agtagaaagt	aacaagcttt	gaatgtcatt	aaaaagtata	tctgctttca		10500
gtagttcagc	ttattttatgc	ccactagaaa	catcttgtag	aagctgaaca	ctggggctcc		10560
agattagtgg	taaaacctac	tttatacaat	catagaatca	tagaatggcc	tgggttgga		10620
gggaccccaa	ggatcatgaa	gatccaacac	ccccgccaca	ggcagggcca	ccaacctcca		10680
gatctggtac	tagaccaggc	agcccagggc	tccatccaac	ctggccatga	acacctccag		10740
ggatggagca	tccacaacct	ctctgggcag	cctgtgccag	cacctacca	ccctctctgt		10800
gaagaacttt	tccctgacat	ccaatctaag	ccttcctccc	ttgaggtag	atccactccc		10860
ccttggtgcta	tcactgtcta	ctcttgtaaa	aagttgatcc	tcctcctttt	tggaagggtg		10920
caatgaggtc	tccttgagc	cttcttctct	tctgcaggat	gaacaagccc	agctccctca		10980
gcctgtcttt	ataggagagg	tgtccagccc	ctctgatcat	ctttgtggcc	ctcctctgga		11040
cccgtccaa	gagctccaca	tctttcctgt	actgggggcc	ccaggcctga	atgcagtact		11100
ccagatgggg	cctcaaaaga	gcagagtaaa	gagggacaat	caccttctc	accctgctgg		11160
ccagccctct	tctgatggag	ccctggatac	aactggcttt	ctgagctgca	acttctcctt		11220
atcagttcca	ctattaaaaa	aggaacaata	caacaggtgc	tgatggccag	tgcagagttt		11280

025CIP7SEQ List.tXt

ttcacacttc	ttcatttcgg	tagatcttag	atgaggaacg	ttgaagttgt	gcttctgcgt	11340
gtgcttcttc	ctcctcaaat	actcctgcct	gatacctcac	cccacctgcc	actgaatggc	11400
tccatggccc	cctgcagcca	gggccctgat	gaacccggca	ctgcttcaga	tgctgtttaa	11460
tagcacagta	tgaccaagtt	gcacctatga	atacacaaac	aatgtgttgc	atccttcagc	11520
acttgagaag	aagagccaaa	tttgcatgtg	caggaaatgg	tttagtaatt	ctgccaatta	11580
aaacttgttt	atctaccatg	gctgttttta	tggctgttag	tagtggtaca	ctgatgatga	11640
acaatggcta	tgcaagtaaa	tcaagactgt	agatattgca	acagactata	aaattcctct	11700
gtggcttagc	caatgtggta	cttcccacat	tgtataagaa	atgttgcaag	tttagagcaa	11760
tgtttgaaat	gttgggaaat	ttctgtatac	tcaagagggc	gtttttgaca	actgtagaac	11820
agaggaatca	aaaggggggtg	ggaggaagtt	aaaagaagag	gcaggtgcaa	gagagcttgc	11880
agtcccgtcg	tgtgtacgac	actggcaaca	tgaggtcttt	gctaattctg	gtgctttgct	11940
tccctgcccct	ggctgcctta	gggtgcgatac	tgccctcagac	ccacagcctg	ggcagcagga	12000
ggaccctgat	gctgctggct	cagatgagga	gaatcagcct	gtttagctgc	ctgaaggata	12060
ggcacgattt	tggctttcct	caagaggagt	ttggcaacca	gtttcagaag	gctgagacca	12120
tccctgtgct	gcacgagatg	atccagcaga	tctttaacct	gtttagcacc	aaggatagca	12180
gcgctgcttg	ggatgagacc	ctgctggata	agtttttacac	cgagctgtac	cagcagctga	12240
acgatctgga	ggcttgctg	atccagggcg	tgggcgtgac	cgagaccct	ctgatgaagg	12300
aggatagcat	cctggctgtg	aggaagtact	ttcagaggat	caccctgtac	ctgaaggaga	12360
agaagtacag	cccctgcgct	tgggaagtgc	tgagggctga	gatcatgagg	agcttttagcc	12420
tgagcaccaa	cctgcaagag	agcttgaggt	ctaaggagta	aaaagtctag	agtcggggcg	12480
gccggccgct	tcgagcagac	atgataagat	acattgatga	gtttggacaa	accacaacta	12540
gaatgcagtg	aaaaaaatgc	tttatttgtg	aaatttgtga	tgctattgct	ttatttgtaa	12600
ccattataag	ctgcaataaa	caagttaaca	acaacaattg	cattcatttt	atgtttcagg	12660
ttcaggggga	ggtgtgggag	gtttttttaa	gcaagtaaaa	cctctacaaa	tgtggtaaaa	12720
tcgataagga	tccgtcgacc	gatgcccttg	agagccttca	accagtcag	ctccttcggg	12780
tgggcgcggg	gcatgactat	cgtcgcgcga	cttatgactg	tcttctttat	catgcaactc	12840
gtaggacagg	tgccggcagc	gctcttcgcg	ttcctcgcgc	actgactcgc	tgcgctcggt	12900
cgttcggtcg	cggcgagcgg	tatcagctca	ctcaaaggcg	gtaatacggg	tatccacaga	12960
atcaggggat	aacgcaggaa	agaacatgtg	agcaaaaagg	cagcaaaaagg	ccaggaaccg	13020
taaaaaggcc	gcgttgctgg	cgtttttcca	taggctccgc	ccccctgacg	agcatcacia	13080
aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgacagga	ctataaagat	accaggcggt	13140

O25CIP7SEQ List.tXt

tccccctgga	agctccctcg	tgcgctctcc	tgttccgacc	ctgccgctta	cgggatacct	13200
gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcaa	tgctcacgct	gtaggtatct	13260
cagttcgggtg	taggtcgttc	gctccaagct	gggctgtgtg	cacgaacccc	cgttcagcc	13320
cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtcc	aacccggtaa	gacacgactt	13380
atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtgc	13440
tacagagttc	ttgaagtggg	ggcctaacta	cggctacact	agaaggacag	tatttggtat	13500
ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	13560
acaaaccacc	gctggtagcg	gtgggtttttt	tggttgcaag	cagcagatta	cgcgcagaaa	13620
aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	tctgacgctc	agtggaacga	13680
aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	aggatcttca	cctagatcct	13740
tttaaattaa	aatgaagtt	ttaaataaat	ctaaagtata	tatgagtaaa	cttggtctga	13800
cagttaccaa	tgtttaatca	gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcac	13860
catagttgcc	tgactccccg	tcgtgtagat	aactacgata	cgggagggct	taccatctgg	13920
ccccagtgt	gcaatgatac	cgcgagaccc	acgtcacccg	gtccagatt	tatcagcaat	13980
aaaccagcca	gccggaagg	cgcgagcag	aagtggctct	gcaactttat	ccgcctccat	14040
ccagtctatt	aattgttgcc	gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	14100
caacgttggt	gccattgcta	caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	14160
attcagctcc	ggttcccaac	gatcaaggcg	agttacatga	tccccatgt	tgtgcaaaaa	14220
agcggtagc	tccttcgggtc	ctccgatcgt	tgtcagaagt	aagttggccg	cagtgttatc	14280
actcatggtt	atggcagcac	tgcataattc	tcttactgtc	atgccatccg	taagatgctt	14340
ttctgtgact	ggtgagtact	caaccaagtc	attctgagaa	tagtgatgc	ggcgaccgag	14400
ttgctcttgc	ccggcgtcaa	tacgggataa	taccgcgcca	catagcagaa	ctttaaaagt	14460
gctcatcatt	ggaaaacgtt	cttcggggcg	aaaactctca	aggatcttac	cgtgttgag	14520
atccagttcg	atgtaacca	ctcgtgcacc	caactgatct	tcagcatctt	ttactttcac	14580
cagcgtttct	gggtgagcaa	aaacaggaag	gcaaaatgcc	gcaaaaaagg	gaataagggc	14640
gacacggaaa	tgttgaatac	tcatactctt	cctttttcaa	tattattgaa	gcatttatca	14700
gggttattgt	ctcatgagcg	gatacatatt	tgaatgtatt	tagaaaaata	aacaaatagg	14760
ggttccgcgc	acatttcccc	gaaaagtgcc	acctgacgcg	ccctgtagcg	gcgcattaag	14820
cgcggcgggt	gtggtggtta	cgcgcagcgt	gaccgctaca	cttgccagcg	ccctagcgcc	14880
cgtcctttc	gctttcttcc	cttcctttct	cgcacgcttc	gccggtttc	cccgtcaagc	14940
tctaaatcgg	gggtccctt	tagggttccg	atttagtgct	ttacggcacc	tcgaccccaa	15000
aaaacttgat	tagggtagtg	gttcacgtag	tgggccatcg	ccctgataga	cgggtttctg	15060

025CIP7SEQ List.txt

ccctttgacg	ttggagtcca	cgttctttta	tagtggactc	ttgttccaaa	ctggaacaac	15120
actcaaccct	atctcggtct	attcttttga	tttataaggg	attttgccga	tttcggccta	15180
ttggttaaaa	aatgagctga	tttaacaaaa	atttaacgcg	aattttaaca	aaatattaac	15240
gtttacaatt	tcccatctgc	cattcaggct	gcgcaactgt	tgggaagggc	gatcggtgcg	15300
ggcctcttcg	ctattacgcc	agcccaagct	accatgataa	gtaagtaata	ttaaggtacg	15360
ggaggtactt	ggagcggccg	ctctagaact	agtggatccc	cgggccgcaa	taaaatatct	15420
ttattttcat	tacatctgtg	tgttggtttt	ttgtgtgaat	cgatagtact	aacatacgct	15480
ctccatcaaa	acaaaacgaa	acaaaacaaa	ctagcaaaat	aggctgtccc	cagtgcgaagt	15540
gcaggtgcca	gaacatttct	ctatcgatag	gtaccgagct	cttacgcgtg	ctagccctcg	15600
agcaggatct	atacattgaa	tcaatattgg	caattagcca	tattagtcac	tggttatata	15660
gcataaatca	atattggcta	ttggccattg	catacgttgt	atctatatca	taatatgtac	15720
atttatattg	gctcatgtcc	aatatgaccg	ccatgttgac	attgattatt	gactagttat	15780
taatagtaat	caattacggg	gtcattagtt	catagcccat	atatggagtt	ccgcggtaca	15840
taacttacgg	taaatggccc	gcctggctga	ccgcccacgc	acccccgccc	attgacgtca	15900
ataatgacgt	atgttcccat	agtaacgcca	atagggactt	tccattgacg	tcaatgggtg	15960
gagtattttac	ggtaaactgc	ccacttgcca	gtacatcaag	tgtatcatat	gccaagtccg	16020
ccccctattg	acgtcaatga	cggtaaattg	cccgctggc	attatgcca	gtacatgacc	16080
ttacgggact	ttcctacttg	gcagtacatc	tacgtattag	tcacgctat	taccatgggtg	16140
atgcggtttt	ggcagtacat	caatgggctg	ggatagcggg	ttgactcacg	gggattttcca	16200
agtctccacc	ccattgacgt	caatgggagt	ttgttttggc	acaaaaatca	acgggacttt	16260
ccaaaaatgtc	gtaacaactc	cgccccattg	acgcaaatgg	gcggtaggcg	tgtacgggtg	16320
gaggtctata	taagcagagc	tcgttttagtg	aaccgtcaga	tcgcctggag	acgccatcca	16380
cgctgttttg	acctccatag	aagacaccgg	gaccgatcca	gcctcccctc	gaagctcgac	16440
tctaggggct	cgagatctgc	gatctaagta	agcttgcatg	cctgcaggtc	ggccgccacg	16500
accggtgccg	ccaccatccc	ctgaccacgc	ccctgacccc	ctcacaagga	gacgaccttc	16560
catgaccgag	tacaagccca	cgggtgcgct	cgccacccgc	gacgacgtcc	cccgggcccgt	16620
acgcaccctc	gccgccgcgt	tcgccgacta	ccccgccacg	cgccacaccg	tcgacccgga	16680
ccgccacatc	gagcgggtca	ccgagctgca	agaactcttc	ctcacgcgcg	tcgggctcga	16740
catcgccaag	gtgtgggtcg	cggacgacgg	cgccgcgggtg	gcggtctgga	ccacgccgga	16800
gagcgtcgaa	gcgggggcgg	tgttcgccga	gatcggcccc	cgcatggccg	agttgagcgg	16860
ttcccggctg	gccgcgcagc	aacagatgga	aggcctcctg	gcgccgcacc	ggcccaagga	16920

025CIP7SEQ List.tXt

gcccgcgtgg	ttcctggcca	cgtcgggcgt	ctcgcccgac	caccagggca	aggggtctggg	16980
cagcgccgtc	gtgctccccg	gagtggagge	ggccgagcgc	gccgggggtgc	ccgccttctt	17040
ggagacctcc	gcgccccgca	acctccccctt	ctacgagcgg	ctcggtttca	ccgtcacccgc	17100
cgacgtcgag	gtgcccgaag	gaccgcgcac	ctgggtgcatg	acccgcaagc	ccgggtgcctg	17160
acgcccgcgc	cacgaccgcg	agcgcccgac	cgaaaggagc	gcacgacccc	atggctccga	17220
ccgaagccga	cccgggcggc	cccgcgcgac	ccgcacccgc	ccccgaggcc	caccgactct	17280
agagtcgggg	cggccggccg	cttcgagcag	acatgataag	atacattgat	gagtttggac	17340
aaaccacaac	tagaatgcag	tgaaaaaaat	gctttatttg	tgaattttgt	gatgctattg	17400
ctttatttgt	aaccattata	agctgcaata	aacaagttaa	caacaacaat	tgcattcatt	17460
ttatgtttca	ggttcagggg	gaggtgtggg	aggtttttta	aagcaagtaa	aacctctaca	17520
aatgtggtaa	aatcgataag	gatcaattcg	gcttcaggta	ccgtcgacga	tgtaggtcac	17580
ggtctcgaag	ccgcgggtgc	ggtgccaggg	cgtgcccttg	ggctccccgg	gcgcgtactc	17640
cacctcacc	atctggtcca	tcatgatgaa	cgggtcgagg	tggcggtagt	tgatcccggc	17700
gaacgcgcgg	cgcaccggga	agccctcgcc	ctcgaaaccg	ctgggcgcgg	tggtcacggg	17760
gagcacggga	cgtgcgacgg	cgtcggcggg	tgccgatacg	cggggcagcg	tcagcggggt	17820
ctcgacggtc	acggcgggca	tgtcgacagc	cgaattgatc	cgtcgaccga	tgcccttgag	17880
agccttcaac	ccagtcagct	ccttcgggtg	ggcgcggggc	atgactatcg	tcgccgcact	17940
tatgactgtc	ttctttatca	tgcaactcgt	aggacaggtg	ccggcagcgc	tcttcgcgtt	18000
cctcgctcac	tgactcgctg	cgtcggtcgt	ttcggtcgtg	gcgagcggtg	tcagctcact	18060
caaaggcggg	aatacggtta	tccacagaat	caggggataa	cgcaggaaag	aacatg	18116

<210> 8
 <211> 17402
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Plasmid pOM1FN-lns-CMV-pur-attB

<400> 8						
ggccgccacc	gcgggtggagc	tccaattcgc	cctatagtga	gtcgtattac	aattcactgg	60
ccgtcgtttt	acaacgtcgt	gactgggaaa	accctggcgt	tacccaactt	aatcgccctg	120
cagcacatcc	cccttttcgc	agctggcgta	atagcgaaga	ggcccgccac	gacgcgccct	180
cccaacagtt	gcgcagcctg	aatggcgaat	gggacgcgcc	ctgtagcggc	gcattaagcg	240
cggcggggtg	ggtggttacg	cgcagcgtga	ccgtacact	tgccagcgcc	ctagcgcccg	300
ctcctttcgc	tttcttccct	tcctttctcg	ccacgttcgc	cggttttccc	cgtcaagctc	360
taaatcgggg	gctcccttta	gggttcgcgt	ttagtgcttt	acggcacctc	gacccccaaa	420

025CIP7SEQ List.tXt

aacttgatta	gggtgatggt	tcacgtagt	ggccatcgcc	ctgatagacg	gtttttcgcc	480
ctttgacgtt	ggagtccacg	ttctttaata	gtggactctt	gttccaaact	ggaacaacac	540
tcaaccctat	ctcgggtctat	tcttttgatt	tataagggat	tttgccgatt	tcggcctatt	600
ggttaaaaaa	tgagctgatt	taacaaaaat	ttaacgcgaa	ttttaacaaa	atattaacgc	660
ttacaattta	ggtggcactt	ttcggggaaa	tgtgcgcgga	acccctat	gtttat	720
ctaaatacat	tcaaataatgt	atccgctcat	gagacaataa	ccctgataaa	tgettcaata	780
atattgaaaa	aggaagagta	tgagtattca	acatttcctg	gtcgcctt	ttccct	840
tgcggcattt	tgccttcctg	tttttgctca	cccagaaacg	ctggtgaaag	taaaagatgc	900
tgaagatcag	ttgggtgcac	gagtgggtta	catcgaactg	gatctcaaca	gcggtaagat	960
ccttgagagt	tttcgccccg	aagaacgttt	tccaatgatg	agcactttta	aagtctctgct	1020
atgtggcgcg	gtattatccc	gtattgacgc	cgggcaagag	caactcggtc	gccgcataca	1080
ctattctcag	aatgacttgg	ttgagtactc	accagtcaca	gaaaagcatc	ttacggatgg	1140
catgacagta	agagaattat	gcagtgctgc	cataaccatg	agtgataaca	ctgcggccaa	1200
cttacttctg	acaacgatcg	gaggaccgaa	ggagctaacc	gcttttttgc	acaacatggg	1260
ggatcatgta	actcgccttg	atcgttggga	accggagctg	aatgaagcca	taccaaacga	1320
cgagcgtgac	accacgatgc	ctglagcaat	ggcaacaacg	ttgcgcaaac	tattaactgg	1380
cgaactactt	actctagctt	cccggcaaca	attaatagac	tggatggagg	cggataaagt	1440
tgcaggacca	cttctgcgct	cggcccttcc	ggctggctgg	tttattgctg	ataaatctgg	1500
agccggtgag	cgtgggtctc	gcggtatcat	tgcagcactg	gggccagatg	gtaagccctc	1560
ccgtatcgta	gttatctaca	cgacggggag	tcaggcaact	atggatgaac	gaaatagaca	1620
gatcgctgag	ataggtgcct	cactgattaa	gcattggtaa	ctgtcagacc	aagtttactc	1680
atatatactt	tagattgatt	taaaacttca	tttttaattt	aaaaggatct	aggtgaagat	1740
cctttttgat	aatctcatga	ccaaaatccc	ttaacgtgag	ttttcgttcc	actgagcgtc	1800
agaccccgta	gaaaagatca	aaggatcttc	ttgagatcct	tttttctgc	gcgtaatctg	1860
ctgcttgcaa	acaaaaaaac	caccgctacc	agcggtggtt	tgtttgccgg	atcaagagct	1920
accaactctt	tttcgaagg	taactggctt	cagcagagcg	cagataccaa	atactgtcct	1980
tctagtgtag	ccgtagttag	gccaccactt	caagaactct	gtagcacgcg	ctacatacct	2040
cgctctgcta	atcctgttac	cagtggctgc	tgccagtggc	gataagtcgt	gtcttaccgg	2100
gttggactca	agacgatagt	taccggataa	ggcgcagcgg	tcgggctgaa	cgggggggttc	2160
gtgcacacag	cccagcttgg	agcgaacgac	ctacaccgaa	ctgagatacc	tacagcgtga	2220
gctatgagaa	agcgcacgcg	ttcccgaagg	gagaaaggcg	gacaggtatc	cggtaagcgg	2280

			025CIP7SEQ	List.tXt			
caggggtcggga	acagggagagc	gcacgagggga	gcttccaggg	ggaaacgcct	ggtatcttta		2340
tagtcctgtc	gggttttcgcc	acctctgact	tgagcgtcga	tttttgtgat	gctcgtcagg		2400
ggggcgggagc	ctatggaaaa	acgccagcaa	cgcggccttt	ttacggttcc	tggccttttg		2460
ctggcctttt	gctcacatgt	tctttcctgc	gttatccct	gattctgtgg	ataaccgtat		2520
taccgccttt	gagtgagctg	ataccgctcg	ccgcagccga	acgaccgagc	gcagcgagtc		2580
agtgagcgag	gaagcgggaag	agcgcccaat	acgcaaaccg	cctctccccg	cgcgttggcc		2640
gattcattaa	tgcagctggc	acgacaggtt	tcccgactgg	aaagcgggca	gtgagcgcaa		2700
cgaattaat	gtgagttagc	tcactcatta	ggcaccccag	gctttacact	ttatgcttcc		2760
ggctcgtatg	ttgtgtggaa	ttgtgagcgg	ataacaattt	cacacaggaa	acagctatga		2820
ccatgattac	gccaagctcg	aaattaaccc	tcactaaagg	gaacaaaagc	tgggtaccgg		2880
gccccccctc	gactagaggg	acagcccccc	cccaaagccc	ccagggatgt	aattacgtcc		2940
ctcccccgct	aggggggcagc	agcgagccgc	ccggggctcc	gctccgggtcc	ggcgtccccc		3000
ccgcatcccc	gagccggcag	cgtgcgggga	cagcccgggc	acggggaagg	tggcacggga		3060
tcgctttcct	ctgaacgctt	ctcgtctgtc	tttgagcctg	cagacacctg	gggggatacg		3120
gggaaaaagc	tttaggctga	aagagagatt	tagaatgaca	gaatcataga	acggcctggg		3180
ttgcaaagga	gcacagtgtc	catccagatc	caaccccctg	ctatgtgcag	ggcatcaac		3240
cagcagccca	ggctgcccag	agccacatcc	agcctggcct	tgaatgcctg	cagggatggg		3300
gcateccacag	cctccttggg	caacctgttc	agtgcgtcac	cacctctctg	gggaaaaact		3360
gcctcctcat	atccaaccca	aacctcccct	gtctcagtgt	aaagccattc	ccccttgtcc		3420
tatcaagggg	gagtttgctg	tgacattgtt	ggtctggggg	gacacatgtt	tgccaattca		3480
gtgcatcacg	gagaggcaga	tcttggggat	aagggaagtgc	aggacagcat	ggacgtggga		3540
catgcagggtg	ttgagggctc	tgggacactc	tccaagtcac	agcgttcaga	acagccttaa		3600
ggataagaag	ataggataga	aggacaaaga	gcaagttaaa	accagcatg	gagaggagca		3660
caaaaaggcc	acagacactg	ctggtcctctg	tgtctgagcc	tgcattgttg	atggtgtctg		3720
gatgcaagca	gaaggggtgg	aagagcttgc	ctggagagat	acagctgggt	cagtaggact		3780
gggacaggca	gctggagaat	tgccatgtag	atgttcatac	aatcgtcaaa	tcatgaaggc		3840
tggaaaagcc	ctccaagatc	ccaagacca	accccaaccc	accacccgtg	ccactggcc		3900
atgtccctca	gtgccacatc	cccacagttc	ttcatcacct	ccagggacgg	tgaccccccc		3960
acctccgtgg	gcagctgtgc	cactgcagca	ccgctctttg	gagaaggtaa	atcttgctaa		4020
atccagcccc	accctcccct	ggcacaacgt	aaggccatta	tctctcatcc	aactccagga		4080
cggagtcagt	gaggatgggg	ctctagtcga	ggtcgacggg	atcgataagc	ttgattaggc		4140
agagcaatag	gactctcaac	ctcgtgagta	tggcagcatg	ttaactctgc	actggagtcc		4200

025CIP7SEQ List.tXt

agcgtgggaa	acaatctgcc	ttgcacatga	gtcttcgtgg	gccaatattc	cccaacgggt	4260
ttccttcagc	ttgtcttgtc	tcctaagctc	tcaaaacacc	tttttggtga	ataaactcac	4320
ttggcaacgt	ttatctgtct	taccttagtg	tcacgtttca	tcctattcc	cctttctcct	4380
cctccgtgtg	gtacacagtg	gtgcacactg	gttcttctgt	tgatgttctg	ctctgacagc	4440
caatgtgggt	aaagtcttc	ctgccacgtg	tctgtgtgt	tttcaactca	aaaagggtcc	4500
tgggctcccc	ttggagctct	caggcatttc	cttaatcatc	acagtcacgc	tggcaggatt	4560
agtccctcct	aaaccttaga	atgacctgaa	cgtgtgctcc	ctctttgtag	tcagtgcagg	4620
gagacgtttg	cctcaagatc	aggggccatc	tcacccacag	ggccattccc	aagatgaggt	4680
ggatgggttta	ctctcacaaa	aagttttctt	atgtttggct	agaaaggaga	actcactgcc	4740
tacctgtgaa	ttcccctagt	cctgggttctg	ctgccactgc	tgccctgtgca	gcctgtccca	4800
tggagggggc	agcaactgct	gtcacaaagg	tgatcccacc	ctgtctccac	tgaaatgacc	4860
tcagtgccac	gtgttgata	gggtataaag	tacgggaggg	ggatgcccg	ctcccttcag	4920
ggttgacagag	cagaagtgtc	tgtgtataga	gtgtgtctta	atctattaat	gtaacagaac	4980
aacttcagtc	ctagtgtttt	gtgggctgga	attgcccatg	tggtaggggac	aggcctgcta	5040
aatcaactgca	atcgctatg	ttctgaagg	atttgggaaa	gaaagggtt	tgggggattg	5100
cctgtgattg	gctttaattg	aatggcaaat	cacaggaaa	cagttctgct	caacagttgg	5160
ttgtttcagc	caattcttgc	agccaaagag	ccgggtgccc	agcgatataa	tagttgtcac	5220
ttgtgtctgt	atggatgaca	gggaggtagg	gtgacctgag	gaccaccctc	cagcttctgc	5280
tagcgtaggt	acagtcacca	cctccagctc	cacacgagtc	ccatcgtggt	ttaccaaaga	5340
aacacaatta	tttgaccag	tttgaaagt	caccgctga	attgtgaggc	tagattaata	5400
gagctgaaga	gcaaatgttc	ccaacttgga	gatactagtt	ggtattagta	tcagagggaac	5460
agggccatag	cacctccatg	ctattagatt	ccggctggca	tgtacttttc	aagatgattt	5520
gtaactaaca	atggcttatt	gtgcttgtct	taagtctgtg	tcctaatagt	aatgttctct	5580
tggtttatat	aaccttcttg	ccatttgctc	ttcagggtgt	cttgcagaac	actggctgct	5640
ttaatctagt	ttaactgttg	cttgattatt	cttagggata	agatctgaat	aaactttttg	5700
tggctttggc	agacttttagc	ttgggcttag	ctcccacatt	agcttttgct	gccttttctg	5760
tgaagctatc	aagatcctac	tcaatgacat	tagctgggtg	cagggtgtacc	aaatcctgct	5820
ctgtggaaca	cattgtctga	tgataccgaa	ggcaaactg	aactcaaaga	ggcacagagt	5880
taagaagaag	tctgtgcaat	tcagaggaaa	agccaaagt	gccattagac	acactttcca	5940
tgcagcattt	gccagtaggt	ttcatataaa	actacaaaat	ggaataaacc	actacaaatg	6000
ggaaaagcct	gatactagaa	tttaaatatt	caccaggct	caagggtgt	ttcatggagt	6060

025CIP7SEQ List.txt

aatatcactc	tataaaagta	gggcagccaa	ttattcacag	acaaagcttt	tttttttctg	6120
tgctgcagtg	ctgttttttcg	gctgatccag	ggttacttat	tgtgggtctg	agagctgaat	6180
gattttctcct	tgtgtcatgt	tggatgaagga	gatatggcca	gggggagatg	agcatgttca	6240
agaggaaacg	ttgcatttttg	gtggccttggg	agaaaggtag	aacgatataca	ggtccatagt	6300
gtcactaaga	gatctgaagg	atggtttttac	agaacagttg	acttggtctg	gtgcaggctt	6360
ggctgtaaat	ggatggaagg	atggacagat	gggtggacag	agattttctgt	gcaggagatc	6420
atctcctgag	ctcgggtgctt	gacagactgc	agatccatcc	cataaccttc	tccagcatga	6480
gagcgcgggg	agcttttgta	ctgttcagtc	tgtgtcttgt	tgttctctgg	gtgcacagtg	6540
gtgattttct	tactcacaca	gggcaaaaac	ctgagcagct	tcaaagtga	caggttgctc	6600
tcataggcca	ttcagttgtc	aagatgaggt	ttttggtttc	ttgttttgta	aggtgggaag	6660
aagcactgaa	ggatcagttg	cgagggcagg	ggtttagcac	tgttcagaga	agtcttattt	6720
taactcctct	catgaacaaa	aagagatgca	ggtgcagatt	ctggcaagca	tgcagtgaag	6780
gagaaagccc	tgaattttctg	atatatgtgc	aatgttgggc	acctaacatt	ccccgctgaa	6840
gcacagcagc	tccagctcca	tgcagtactc	acagctgggtg	cagccctcgg	ctccagggtc	6900
tgagcagtgc	tgggactcac	gaggttccat	gtctttcaca	ctgataatgg	tccaatttct	6960
ggaatgggtg	cccatccttg	gaggtcccca	aggccaggct	ggctgcgtct	ccgagcagcc	7020
cgatctggtg	gtgagtagcc	agcccatggc	aggagttaga	gcctgatggt	ctttaaggtc	7080
ccttccaacc	taagccatcc	tacgattcta	ggaatcatga	cttgtgagtg	tgtattgcag	7140
aggcaatatt	ttaaagttat	aaatgttttc	tccccttcct	tgtttgtcaa	agttatcttg	7200
atcgcttat	caatgctttt	ggagtctcca	gtcatttttc	ttacamcaaa	aagaggagga	7260
agaatgaaga	gaatcattta	atttcttgat	tgaatagtag	gattcagaaa	gctgtacgta	7320
atgcggtctc	tttgatatga	gctgtaaggt	ttctcatcat	ttatcagcgt	ggtacatatc	7380
agcacttttc	catctgatgt	ggaaaaaaa	atccttatca	tctacagtct	ctgtacctaa	7440
acatcgctca	gactctttac	caaaaaagct	ataggtttta	aaactacatc	tgctgataat	7500
ttgccttggt	ttagctcttc	ttccatatgc	tgcgtttgtg	agaggtgcgt	ggatgggcct	7560
aaactctcag	ctgctgagct	tgatgggtgc	ttaagaatga	agcactcact	gctgaaactg	7620
ttttcatttc	acaggaatgt	tttagtggca	ttgtttttat	aactacatat	tcctcagata	7680
aatgaaatcc	agaaataatt	atgcaaactc	actgcacccg	ttgcacaggt	ctttatctgc	7740
tagcaaagga	aataatttgg	ggatggcaaa	aacattcctt	cagacatcta	tatttaaagg	7800
aatataatcc	tggtagccac	ccacttcac	cctcattatg	ttcacactca	gagatactca	7860
ttctcttggt	gttatcatth	gatagcgtht	tctttggtht	tttgccacgc	tctgggctat	7920
ggctgcacgc	tctgcactga	tcagcaagta	gatgcgaggg	aagcagcagt	gagaggggct	7980

O25CIP7SEQ List.tXt

gccctcagct	ggcaccacgc	cgctcagcct	aggaggggac	cttgcctttc	caccagctga	8040
ggtgcagccc	tacaagctta	cacgtgctgc	gagcaggtga	gcaaagggag	tcttcatggt	8100
gtgtttcttg	ctgcccgga	gcaaaacttt	actttcattc	atcccccttg	aagaatgagg	8160
aatgtttgga	aacggactgc	tttacgttca	atttctctct	tccctttaag	gctcagccag	8220
gggccattgc	tgaggacggc	atcggggccc	cctggacca	atctgtggca	cagatggttt	8280
cacttacatc	agtggatgtg	ggatctgcgc	ctgtaatgtg	tccttctgaa	ggaaggaacg	8340
tgctttccaa	gtgccagccc	cacagccccc	agccctctcc	tgtgctgctc	caattcatct	8400
cctcttctct	cttctccctt	tgtgttttgt	gctcgggtag	aaatcatgaa	gatttagaag	8460
agaaaacaaa	ataactggag	tggaaaccca	ggtgatgcag	ttcattcagc	tgtcatagggt	8520
ttgtcgttgc	tataggtctg	tatcagagat	gctarcacca	ctttgctgtc	ggtgcttaac	8580
tcgggtgaac	tctccttcac	tcgcattcatt	tgcgggcctt	atttacatcc	ccagcatcca	8640
tcaccctctg	ggaaaatggg	cgcactggat	ctctaattga	agactttccc	tctttcagag	8700
cctgtgggat	gtgcagtgac	aagaaacgtg	gaggggctga	gcagcagcac	tgcccccagg	8760
gagcaggagc	ggatgccatc	ggtggcagca	tcccaaata	tgtcagcgga	tgctgagcag	8820
gcagcggacg	aacggacaga	agcgatgcgt	acaccttctg	ttgacatggt	atttggcagc	8880
gatttaacac	tcgcttctca	gtcctgctat	tctccacagg	ctgcattcaa	atgaacgaag	8940
ggaagggagg	caaaaagatg	caaaatccga	gacaagcagc	agaaatattt	cttcgctacg	9000
gaagcgtgcg	caaacaacct	tctccaacag	caccagaaga	gcacagcgta	acctttttca	9060
agaccagaaa	aggaaattca	caaagcctct	gtggatacca	gcgcggttcag	ctctcctgat	9120
agcagatttc	ttgtcagggt	gcgaatgggg	tatggtgcc	ggaggtgcag	ggaccatatg	9180
atcatataca	gcacagcagt	cattgtgcac	gtattaatat	atattgagta	gcagtgttac	9240
tttgccaaag	caatagttca	gagatgagtc	ctgctgcata	cctctatctt	aaaactaact	9300
tataaatagt	aaaaccttct	cagttcagcc	acgtgctcct	ctctgtcagc	accaatgggtg	9360
cttcgcctgc	accagctgc	aaggaatcag	cccgatgatc	cattaacact	cagctctgca	9420
ggataaatta	gattgtttca	ctctcttttg	ttgttaatta	cgacggaaca	attgttcagt	9480
gctgatggtc	ctaattgtca	gctacagaaa	acgtctccat	gcagtctcct	ctgcgccagc	9540
aaactgtcca	ggctatagca	cogtgatgca	tgctacctct	cactccatcc	ttcttctctt	9600
tcccaccagg	gagagctgtg	tggttttca	ctcagccact	ctgaacaata	ccaaaactgt	9660
acgcactgcc	tccctcgga	agagaatccc	cttggtgctt	ttttatttac	aggatccttc	9720
ttaaaaagca	gaccatcatt	cactgcaaac	ccagagcttc	atgcctctcc	ttccacaacc	9780
gaaaacagcc	ggcttcattt	gtctttttta	aatgctgttt	tccaggtgaa	ttttggccag	9840

025CIP7SEQ List.tXt						
cgtgttgget	gagatccagg	agcacgtgtc	agcttttctgc	tctcattgct	cctgtttctgc	9900
attgcctctt	tctgggggtt	ccaagagggg	gggagacttt	gcgcggggat	gagataatgc	9960
cccttttctt	aggggtggctg	ctgggcagca	gagtggctct	gggtcactgt	ggcaccaatg	10020
ggaggcacca	gtgggggtgt	gttttgtgca	ggggggaagc	attcacagaa	tggggctgat	10080
cctgaagctt	gcagtccaag	gctttgtctg	tgtacccagt	gaaatccttc	ctctgttaca	10140
taaagcccag	ataggactca	gaaatgtagt	cattccagcc	cccctcttcc	tcagatctgg	10200
agcagcactt	gtttgcagcc	agtccctccc	aaaatgcaca	gacctcgccg	agtggagggga	10260
gatgtaaaca	gcgaaggtta	attacctctt	tgtcaaaaac	actttgtggg	ccatagatgt	10320
ttctgtcaat	cttacaaaac	agaaccgaga	ggcagcgagc	actgaagagc	gtgttcccat	10380
gctgagttaa	tgagacttgg	cagctcgctg	tgcagagatg	atccctgtgc	ttcatgggag	10440
gctgtaacct	gtctcccat	cgccttcaca	ccgcagtgtc	gtcctggaca	cctcaccctc	10500
cataagctgt	aggatgcagc	tgcccagggg	tcaagagact	tttcctaagg	ctcttaggac	10560
tcctctttgc	cgctcagtag	cgtgcagcaa	ttactcatcc	caactatact	gaatgggttt	10620
ctgccagctc	tgcttggttg	tcaataagca	tttcttcatt	ttgcctctaa	gtttctctca	10680
gcagcaccgc	tctgggtgac	ctgagtggcc	acctggaacc	cgagggggcac	agccaccacc	10740
tcctgtgtgc	tgctgctcca	gggactcatg	tgctgctgga	tgggggggaag	catgaagttc	10800
ctcaccaga	cacctgggtt	gcaatggctg	cagcgtgtct	ttcttggtat	gcagattggt	10860
tccagccatt	acttgtagaa	atgtgctgtg	gaagcccttt	gtatctcttt	ctgtggccct	10920
tcagcaaaag	ctgtgggaaa	gctctgaggg	tgctttcttg	ggtcgtggag	gaattgtatg	10980
ttccttcttt	aacaaaaatt	atccttagga	gagagcactg	tgcaagcatt	gtgcacataa	11040
aacaattcag	gttgaaaggg	ctctctggag	gtttccagcc	tgactactgc	tcgaagcaag	11100
gccaggttca	aagatggctc	aggatgctgt	gtgccttcct	gattatctgt	gccaccaatg	11160
gaggagattc	acagccactc	tgcttcccg	gccactcatg	gagaggaata	ttcccttata	11220
ttcagataga	atgttatcct	ttagctcagc	cttccctata	accccatgag	ggagctgcag	11280
atccccatac	tctccccttc	tctgggggtga	aggccgtgtc	cccagcccc	ccttcccacc	11340
ctgtgcccta	agcagcccgc	tggcctctgc	tggtatgtgtg	cctatatgtc	aatgcctgtc	11400
cttgcagtcc	agcctgggac	atttaattca	tcaccagggt	aatgtggaac	tgtgtcatct	11460
tcccctgcag	ggtacaaagt	tctgcacggg	gtcctttcgg	ttcaggaaaa	ccttccactgg	11520
tgctacctga	atcaagctct	atttaataag	ttcataagca	catggatgtg	ttttcctaga	11580
gatacgtttt	aatgggtatca	gtgattttta	tttgctttgt	tgcttacttc	aaacagtgcc	11640
tttgggcagg	aggtgagggg	cgggtctgcc	gttggctctg	cagtgatttc	tccaggcgtg	11700
tggctcaggt	cagatagtgg	tcactctgtg	gccagaagaa	ggacaaagat	ggaaattgca	11760

025CIP7SEQ List.tXt

gattgagtca	cgtaagcag	gcatcttgga	gtgatttgag	gcagtttcat	gaaagagcta	11820
cgaccactta	ttgttgTTTT	ccccttttac	aacagaagtt	ttcatcaaaa	taacgtggca	11880
aagcccagga	atgtttggga	aaagtgtagt	taaatgtttt	gtaattcatt	tgtcggagtg	11940
ctaccagcta	agaaaaaagt	cctacctttg	gtatggtagt	cctgcagaga	atacaacatc	12000
aatattagtt	tggaaaaaaa	caccaccacc	accagaaaact	gtaatggaaa	atgtaaacca	12060
agaaattcct	tgggtaagag	agaaaggatg	tcgatactg	gccaagtcct	gcccagctgt	12120
cagcctgctg	accctctgca	gttcaggacc	atgaaacgtg	gcactgtaag	acgtgtcccc	12180
tgcctttgct	tgccacaga	tctctgccct	tgtgctgact	cctgcacaca	agagcatttc	12240
cctgtagcca	aacagcgatt	agccataagc	tgcacctgac	tttgaggatt	aagagtttgc	12300
aattaagtgg	attgcagcag	gagatcagtg	gcaggggtgc	agatgaaatc	cttttctagg	12360
ggtagctaag	ggctgagcaa	cctgtcctac	agcacaagcc	aaaccagcca	agggttttcc	12420
tgtgctgttc	acagaggcag	ggccagctgg	agctggagga	ggttgtgctg	ggacccttct	12480
ccctgtgctg	agaatggagt	gatttctggg	tgtgttcct	gtggcttgca	ctgagcagct	12540
caagggagat	cggtgctcct	catgcagtgc	caaaactcgt	gtttgatgca	gaaagatgga	12600
tgtgcacctc	cctcctgcta	atgcagccgt	gagcttatga	aggcaatgag	ccctcagtgc	12660
agcaggagct	gtagtgcact	cctgtagggtg	ctagggaaaa	tctctggttc	ccagggatgc	12720
attcataagg	gcaatatatc	ttgaggctgc	gccaaatcct	tctgaaatat	tcatgcgtgt	12780
tcccttaatt	tatagaaaca	aacacagcag	aataattatt	ccaatgcctc	ccctcgaagg	12840
aaacccatat	ttccatgtag	aaatgtaacc	tatatacaca	cagccatgct	gcatecttca	12900
gaacgtgcc	gtgctcatct	cccatggcaa	aatactacag	gtattctcac	tatgttggac	12960
ctgtgaaagg	aaccatggta	agaaacttcg	gttaaaggta	tggctgcaaa	actactcata	13020
ccaaaacagc	agagctccag	acctcctctt	aggaaagagc	cacttgagga	gggatgggtg	13080
gaaggctgga	ggtgagagac	agagcctgtc	ccagttttcc	tgtctctatt	ttctgaaacg	13140
tttgcaggag	gaaaggacaa	ctgtactttc	aggcatagct	ggtgccctca	cgtaaataag	13200
ttccccgaac	ttctgtgtca	tttgttctta	agatgctttg	gcagaacact	ttgagtcaat	13260
tgccttaact	gtgactaggt	ctgtaaataa	gtgtccctg	ctgataaggt	tcaagtgaca	13320
tttttagtgg	tatttgacag	catttacctt	gctttcaagt	cttctaccaa	gctcttctat	13380
acttaagcag	tgaaacccgc	aagaaacctt	tccttttatc	aagctagtgc	taaataccat	13440
taacttcata	ggttagatac	ggtgctgcca	gcttcacctg	gcagtgggtg	gtcagttctg	13500
ctggtgacaa	agcctccctg	gcctgtgctt	ttacctagag	gtgaatatcc	aagaatgcag	13560
aactgcatgg	aaagcagagc	tgcaggcacg	atgggtgctga	gccttagctg	cttcctgctg	13620

025CIP7SEQ List.tXt

ggagatgtgg	atgcagagac	gaatgaagga	cctgtccctt	actcccctca	gcattctgtg	13680
ctatttaggg	ttctaccaga	gtccttaaga	ggtttttttt	tttttttggtc	caaaagtctg	13740
tttgtttggt	tttgaccact	gagagcatgt	gacacttgct	tcaagctatt	aaccaagtgt	13800
ccagccaaaa	tcaattgcct	gggagacgca	gaccattacc	tggaggtcag	gacctcaata	13860
aatattacca	gcctcattgt	gccgctgaca	gattcagctg	gctgctccgt	gttccagtcc	13920
aacagttcgg	acgccacgtt	tgtatatatt	tgcaggcagc	ctcgggggga	ccatctcagg	13980
agcagagcac	cggcagccgc	ctgcagagcc	gggcagtacc	tcaccatggc	tttgaccttt	14040
gccttactgg	tggctctcct	ggtgctgagc	tgcaagagca	gctgctctgt	gggctgcgat	14100
ctgcctcaga	cccacagcct	gggcagcagg	aggaccctga	tgctgctggc	tcagatgagg	14160
agaatcagcc	tgttttagctg	cctgaaggat	aggcacgatt	ttggctttcc	tcaagaggag	14220
tttggcaacc	agtttcagaa	ggctgagacc	atccctgtgc	tgcacgagat	gatccagcag	14280
atctttaacc	tgttttagcac	caaggatagc	agcgtctgctt	gggatgagac	cctgctggat	14340
aagttttaca	ccgagctgta	ccagcagctg	aacgatctgg	aggcttgctg	gatccagggc	14400
gtgggctgta	ccgagacccc	tctgatgaag	gaggatagca	tcctggctgt	gaggaagtac	14460
tttcagagga	tcacctgtga	cctgaaggag	aagaagtaca	gccctgcgc	ttgggaagtc	14520
gtgagggtg	agatcatgag	gagctttagc	ctgagcacca	acctgcaaga	gagcttgagg	14580
tctaaggagt	aaaaagtcta	gagtcggggc	ggccggccgc	ttcgagcaga	catgataaga	14640
tacattgatg	agtttgagca	aaccacaact	agaatgcagt	gaaaaaaatg	ctttatattgt	14700
gaaatttggtg	atgctattgc	tttatattgta	accattataa	gctgcaataa	acaagttaac	14760
aacaacaatt	gcattcattt	tatgtttcag	gttcaggggg	aggtgtggga	ggttttttta	14820
agcaagtaaa	acctctacaa	atgtggtaaa	atcgataccg	tcgacctcga	ctagagcggc	14880
cactaacata	cgctctccat	caaaacaaaa	cgaacaaaaa	caaactagca	aaataggctg	14940
tccccagtgc	aagtgcaggt	gccagaacat	ttctctatcg	ataggtaccg	agctcttacg	15000
cgtgctagcc	ctcgagcagg	atctatacat	tgaatcaata	ttggcaatta	gccatattag	15060
tcattgggta	tatagcataa	atcaatattg	gctattggcc	attgcatacg	ttgtatctat	15120
atcataatat	gtacatttat	attggctcat	gtccaatatg	accgccatgt	tgacattgat	15180
tattgactag	ttattaatag	taatcaatta	cggggctcatt	agttcatagc	ccatatatgg	15240
agttccgcgt	tacataactt	acggtaaatg	gcccgcctgg	ctgaccgccc	aacgaccccc	15300
gccattgac	gtcaataatg	acgtatgttc	ccatagtaac	gccaataggg	actttccatt	15360
gacgtcaatg	ggtggagtat	ttacggtaaa	ctgccactt	ggcagtacat	caagtgtatc	15420
atatgccaag	tccgccccct	attgacgtca	atgacggtaa	atggcccgc	tggcattatg	15480
cccagtacat	gaccttacgg	gactttccta	cttggcagta	catctacgta	ttagtcatcg	15540

025CIP7SEQ List.tXt

ctattacat	ggtgatgcgg	ttttggcagt	acatcaatgg	gcgtggatag	cggtttgact	15600
cacggggatt	tccaagtctc	caccccatg	acgtcaatgg	gagtttgttt	tggcaccaaa	15660
atcaacggga	ctttccaaaa	tgtcgtaaca	actccgcccc	attgacgcaa	atgggcggtg	15720
ggcgtgtacg	gtgggaggtc	tatataagca	gagctcgttt	agtgaaccgt	cagatcgctt	15780
ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	tccagcctcc	15840
cctcgaagct	cgactctagg	ggctcgagat	ctgcgatcta	agtaagcttg	catgcctgca	15900
ggtcggccgc	cacgaccggt	gccgccacca	ccccctgacc	cacgccccctg	acccttcaca	15960
aggagacgac	cttccatgac	cgagtacaag	cccacgggtgc	gcctcgccac	ccgcgacgac	16020
gtcccccg	cggtacgcac	cctcgccgcc	gcgttcgccg	actaccccg	cacgcgccac	16080
accgtcgacc	cggaccgcca	catcgagcgg	gtcaccgagc	tgcaagaact	cttcctcacg	16140
cgcgtcgggc	tcgacatcgg	caaggtgtgg	gtcgcggacg	acggcgccgc	ggtggcggtc	16200
tggaccacgc	cggagagcgt	cgaagcgggg	gcgggtgttcg	ccgagatcgg	cccgcgcatg	16260
gccgagttga	gcggttcccc	gctggccgcg	cagcaacaga	tggaaggcct	cctggcgccg	16320
caccggccca	aggagcccgc	gtggttcctg	gccaccgtcg	gcgtctcgcc	cgaccaccag	16380
ggcaagggtc	tgggcagcgc	cgtcgtgctc	cccggagtgg	aggcggccga	gcgcgccggg	16440
gtgcccgcct	tcctggagac	ctccgcgccc	cgcaacctcc	ccttctacga	gcggctcggc	16500
ttcaccgtea	ccgccgacgt	cgaggtgccc	gaaggaccgc	gcacctggtg	catgaccgcg	16560
aagcccgggtg	cctgacgccc	gccccacgac	ccgcagcgcc	cgaccgaaag	gagcgcacga	16620
ccccatggct	ccgaccgaag	ccgaccgggg	cggccccgcc	gacccccgcac	ccgcccccg	16680
ggccacacga	ctctagagtc	ggggcgggcg	gccgcttcga	gcagacatga	taagatacat	16740
tgatgagttt	ggacaaacca	caactagaat	gcagtgaaaa	aaatgcttta	tttgtgaaat	16800
ttgtgatgct	attgctttat	ttgtaacct	tataagctgc	aataaacaag	ttaacaacaa	16860
caattgcatt	catttttatgt	ttcagggttc	gggggaggtg	tgggaggttt	tttaaagcaa	16920
gtaaaacctc	tacaaatgtg	gtaaaatcga	taaggatcaa	ttcggttc	ggtaccgtcg	16980
acgatgtagg	tcacggtctc	gaagccgcgg	tgcgggtgcc	agggcggtgc	cttgggctcc	17040
ccgggcgcgt	actccacctc	accatctg	tccatcatga	tgaacgggtc	gaggtggcgg	17100
tagttgatcc	cggcgaacgc	gcggcgacc	gggaagccct	cgccctcgaa	accgtggggc	17160
gcgggtggtc	cgggtgagcac	gggacgtgcg	acggcgctcg	cgggtgcgga	tacgcggggc	17220
agcgtcagcg	ggttctcgac	ggtcacggcg	ggcatgtcga	cagccgaatt	gatccgtcga	17280
ccgatgccct	tgagagcctt	caaccacgtc	agctccttcc	ggtgggcgcg	gggcatgact	17340
atcgtcgcgc	cacttatgac	tgtcttcttt	atcatgcaac	tcgtaggaca	ggtgccggca	17400

gc

025CIP7SEQ List.tXt

17402

<210> 9
<211> 5172
<212> DNA
<213> Artificial Sequence

<220>
<223> Plasmid pRSV-Int

<400> 9
ctgcattaat gaatcggcca acgcgcgggg agaggcgggt tgcgtattgg gcgctcttcc 60
gcttcctcgc tactgactc gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct 120
cactcaaagg cggtaatagc gttatccaca gaatcagggg ataacgcagg aaagaacatg 180
tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc 240
cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 300
aaccgcagag gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct 360
cctgttccga ccctgccgct taccggatac ctgtccgcct ttctcccttc gggaaagcgtg 420
gcgctttctc aatgctcacg ctgtagggtat ctgagttcgg tgtagggtcgt tcgctccaag 480
ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct gcgccttata cggtaactat 540
cgtcttgagt ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac 600
aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 660
tacggctaca ctagaaggac agtatttggt atctgcgctc tgetgaagcc agttaccttc 720
ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggtttt 780
tttgtttgca agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc 840
ttttctacgg ggtctgacgc tcagtggaac gaaaactcac gttaagggat tttggtcatg 900
agattatcaa aaaggatctt cacctagatc cttttaaat aaatgaag ttttaaatca 960
atctaaagta tatatgagta aacttgggtc gacagttacc aatgcttaat cagtgaggca 1020
cctatctcag cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag 1080
ataactacga tacgggaggg cttacatct ggccccagtg ctgcaatgat accgcgagac 1140
ccacgctcac cggctccaga tttatcagca ataaaccagc cagccggaag ggccgagcgc 1200
agaagtggtc ctgcaacttt atccgcctcc atccagtcta ttaattgttg ccgggaagct 1260
agagtaagta gttcgccagt taatagtttg cgcaacgttg ttgccattgc tacaggcatc 1320
gtggtgtcac gctcgtcgtt tgggtatggc tcattcagct ccggttccca acgatcaagg 1380
cgagttacat gatcccccac gttgtgcaaa aaagcgggta gtccttcgg tctccgatac 1440
gttgtcagaa gtaagttggc cgcagtggtt tcaactcatg ttatggcagc actgcataat 1500
tctcttactg tcatgccatc cgtaagatgc ttttctgtga ctggtgagta ctcaaccaag 1560

025CIP7SEQ List.tXt

tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgctc	aatacgggat	1620
aataccgcgc	cacatagcag	aacttttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	1680
cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	1740
cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	1800
aggcaaaatg	cgcacaaaaa	gggaataagg	gcgacacgga	aatgttgaat	actcatactc	1860
ttcctttttc	aatattattg	aagcatttat	cagggttatt	gtctcatgag	cggatacata	1920
tttgaatgta	tttagaaaaa	taaacaaata	ggggttccgc	gcacatttcc	ccgaaaagtg	1980
ccacctgacg	tcgacggatc	gggagatctc	ccgatcccct	atggtcgact	ctcagtacaa	2040
tctgctctga	tgccgcatag	ttaagccagt	atctgctccc	tgcttggtg	ttggaggtcg	2100
ctgagtagtg	cgcgagcaaa	atttaagcta	caacaaggca	aggcttgacc	gacaattgca	2160
tgaagaatct	gcttaggggt	aggcgttttg	cgctgcttcg	cgatgtacgg	gccagatata	2220
cgctgctag	gggtctagga	tcgattctag	gaattctcta	gccgcggtct	agggatcccg	2280
gcgcgtatgg	tgcactctca	gtacaatctg	ctctgatgcc	gcatagttaa	gccagtatct	2340
gctccctgct	tgtgtgttg	aggtcgctga	gtagtgcgcg	agcaaaattt	aaagctacaac	2400
aaggcaaggc	ttgaccgaca	attgcatgaa	gaatctgctt	agggttaggc	gttttgcgct	2460
gcttcgcgat	gtacgggcca	gatatacgcg	tatctgaggg	gactaggggtg	tgtttaggcg	2520
aaaagcgggg	cttcggttgt	acgcggttag	gagtcccctc	aggatatagt	agtttcgctt	2580
ttgcataggg	agggggaaat	gtagtcttat	gcaatacact	tgtagtcttg	caacatggta	2640
acgatgagtt	agcaacatgc	cttacaagga	gagaaaaagc	accgtgcatg	ccgattgggtg	2700
gaagtaagg	ggtacgatcg	tgccctatta	ggaaggcaac	agacaggctc	gacatggatt	2760
ggacgaacca	ctgaattccg	cattgcagag	ataattgtat	ttaagtgcct	agctcgatac	2820
aataaacgcc	atttgaccat	tcaccacatt	ggtgtgcacc	tccaagcttg	catgcctgca	2880
ggtaccggtc	cggaattccc	gggtcgacga	gctcactagt	cgtaggggtcg	ccgacatgac	2940
acaaggggtt	gtgaccgggg	tggacacgta	cgcggtgct	tacgaccgtc	agtcgcgcga	3000
gcgcgagaat	tcgagcgag	caagcccagc	gacacagcgt	agcgccaacg	aagacaaggc	3060
ggccgacctt	cagcggaag	tcgagcgga	cgggggccgg	ttcagggttcg	tcgggcattt	3120
cagcgaagcg	ccgggcacgt	cggcgttcgg	gacggcgag	cgcccggagt	tcgaacgcac	3180
cctgaacgaa	tgccgcgccg	ggcggtctca	catgatcatt	gtctatgacg	tgctcgcgctt	3240
ctcgcgcctg	aaggctcatg	acgcgattcc	gattgtctcg	gaattgctcg	ccctgggcgt	3300
gacgattgtt	tccactcagg	aaggcgctct	ccggcaggga	aacgtcatgg	acctgattca	3360
cctgattatg	cggctcgacg	cgctcgacaa	agaatcttcg	ctgaagtcgg	cgaagattct	3420

			O25CIP7SEQ	List.tXt		
cgacacgaag	aaccttcagc	gcgaattggg	cggttacgtc	ggcgggaagg	cgccttacgg	3480
cttcgagctt	gtttcggaga	cgaaggagat	cacgcgcaac	ggccgaatgg	tcaatgtcgt	3540
catcaacaag	cttgcgact	cgaccactcc	ccttacccga	cccttcgagt	tcgagcccg	3600
cgtaatccgg	tgggtgtggc	gtgagatcaa	gacgcacaaa	caccttcctt	tcaagccggg	3660
cagtcaagcc	gccattcacc	cgggcagcat	cacggggcct	tgtaagcgca	tggacgctga	3720
cgccgtgccg	acccggggcg	agacgattgg	gaagaagacc	gcttcaagcg	cctgggaccc	3780
ggcaaccgtt	atgcgaatcc	ttcgggaccc	gcgtattgcg	ggcttcgccg	ctgaggtgat	3840
ctacaagaag	aagccggacg	gcacgccgac	cacgaagatt	gagggttacc	gcattcagcg	3900
cgacccgatc	acgtccggc	cggtcgagct	tgattgcgga	ccgatcatcg	agcccgtga	3960
gtggtatgag	cttcaggcgt	ggttggacgg	cagggggcgc	ggcaaggggc	tttcccgggg	4020
gcaagccatt	ctgtccgcca	tggacaagct	gtactgcgag	tgtggcgccg	tcatgacttc	4080
gaagcgccgg	gaagaatcga	tcaaggactc	ttaccgctgc	cgtcgccgga	aggtggtcga	4140
cccgtccgca	cctgggcagc	acgaaggcac	gtgcaacgtc	agcatggcgg	cactcgacaa	4200
gttcgttgcg	gaacgcatct	tcaacaagat	caggcacgcc	gaaggcgacg	aagagacgtt	4260
ggcgcttctg	tgggaagccg	cccgcgctt	cggcaagctc	actgaggcgc	ctgagaagag	4320
cggcgaacgg	gcgaaccttg	ttgcggagcg	cgccgacgcc	ctgaacgccc	ttgaagagct	4380
gtacgaagac	cgcgcggcag	gcgcgtacga	cggaccgctt	ggcaggaagc	acttccggaa	4440
gcaacaggca	gcgctgacgc	tccggcagca	aggggcggaa	gagcggcttg	ccgaacttga	4500
agccgccgaa	gccccgaagc	ttccccttga	ccaatggttc	cccgaagacg	ccgacgtga	4560
cccgaccggc	cctaagtcgt	ggtggggggc	cgcgtcagta	gacgacaagc	gcgtgttcgt	4620
cgggctcttc	gtagacaaga	tcgttgtcac	gaagtcgact	acgggcaggg	ggcagggaac	4680
gcccacgcag	aagcgcgctt	cgatcacgtg	ggcgaagccg	ccgaccgacg	acgacgaaga	4740
cgacgcccag	gacggcacgg	aagacgtage	ggcgtagcga	gacaccgga	tccttcgagg	4800
ggccctattc	tatagtgtca	cctaaatgct	agagctcgt	gatcagcctc	gactgtgcct	4860
tctagttgcc	agccatctgt	tgtttgcccc	tccccgtgc	cttccttgac	cctggaaggt	4920
gccactccca	ctgtcccttc	ctaataaaat	gaggaaattg	catcgcattg	tctgagtagg	4980
tgtcattcta	ttctgggggg	tggggtgggg	caggacagca	agggggagga	ttgggaagac	5040
aatagcaggc	atgctgggga	tgcggtgggc	tctatggctt	ctgaggcgga	aagaaccagg	5100
tgcccagtca	tagccgaata	gcctctccac	ccaagcggcc	ggagaacctg	cgtgcaatcc	5160
actggggggc	eg					5172

<210> 10
 <211> 6233

025CIP7SEQ Li st.tXt

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Plasmid pCR-XL-TOPO-CMV-pur-attB

```

<400> 10
agcgcccaat acgcaaaccg cctctccccg cgcgttggcc gattcattaa tgcagctggc      60
acgacagggtt tcccgaactgg aaagcgggca gtgagcgcaa cgcaattaat gtgagttagc      120
tcaactcatta ggcaccccag gctttacact ttatgcttcc ggctcgtatg ttgtgtggaa      180
ttgtgagcgg ataacaattt cacacaggaa acagctatga ccatgattac gccaaagctat      240
ttaggtgacg cgttagaata ctcaagctat gcatcaagct tggtaaccgag ctcggatcca      300
ctagtaacgg ccgccagtggt gctggaattc gcccttggcc gcaataaaaat atctttattt      360
tcattacatc tgtgtgttgg ttttttgtgt gaatcgatag tactaacata cgctctccat      420
caaaacaaaa cgaaacaaaa caaactagca aaataggctg tccccagtgc aagtgcagggt      480
gccagaacat ttctctatcg ataggtagcg agctcttacg cgtgctagcc ctcgagcagg      540
atctatacat tgaatcaata ttggcaatta gccatattag tcattgggta tatagcataa      600
atcaatattg gctattggcc attgcatacg ttgtatctat atcataatat gtacatttat      660
attggctcat gtccaatatg accgccatgt tgacattgat tattgactag ttattaatag      720
taatcaatta cgggggtcatt agttcatagc ccataatatgg agttccgcgt tacataactt      780
acggtaaatg gcccgcctgg ctgaccgcc aacgaccccc gccattgac gtcaataatg      840
acgtatgttc ccataagtaac gccaataggg actttccatt gacgtcaatg ggtggagtat      900
ttacggtaaa ctgcccactt ggcagtagat caagtgtatc atatgccaaag tccgccccct      960
attgacgtca atgacggtaa atggcccgcc tggcattatg ccagtagcat gaccttacgg      1020
gactttccta cttggcagta catctacgta ttagtcatcg ctattaccat ggtgatgcgg      1080
ttttggcagt acatcaatgg gcgtggatag cggtttgact cacggggatt tccaagtctc      1140
caccctattg acgtcaatgg gagtttggtt tggcaccaaa atcaacggga ctttccaaaa      1200
tgtcgtaaca actccgcccc attgacgcaa atgggcggta ggcggtgtacg gtgggaggtc      1260
tatataagca gagctcggtt agtgaaccgt cagatcgctt ggagacgcca tccacgctgt      1320
tttgacctcc atagaagaca ccgggaccga tccagcctcc cctcgaagct cgactctagg      1380
ggctcgagat ctgcgatcta agtaagcttg catgcctgca ggtcggccgc cacgaccggt      1440
gccgccacca tcccctgacc cagccccctg acccctcaca aggagacgac cttccatgac      1500
cgagtacaag cccacgggtgc gcctcgccac ccgcgacgac gtccccggg ccgtacgcac      1560
cctcgccgcc gcggttcgcc actacccgcg cagcgccac accgtcgacc cggaccgcca      1620
catcgagcgg gtcaccgagc tgcaagaact cttcctcacg cgcgtcgggc tcgacatcgg      1680

```


						025CIP7SEQ	List.tXt	
caaggtgtgg	gtcgcggacg	acggcgccgc	ggtggcggtc	tggaccacgc	cggagagcgt			1740
cgaagcgggg	gcggtgttcg	cagagategg	cccgcgcatg	gccgagttga	gcggttcccg			1800
gctggcccg	cagcaacaga	tggaaggcct	cctggcgccg	caccggccca	aggagcccg			1860
gtggttcctg	gccaccgtcg	gcgtctcgcc	cgaccaccag	ggcaagggtc	tgggcagcgc			1920
cgtcgtgctc	cccggagtgg	aggcggcga	gcgcgcggg	gtgccgcct	tcctggagac			1980
ctccgcgcc	cgcaacctcc	ccttctacga	gcggctcggc	ttcacctca	ccgcgcagct			2040
cgaagtgcc	gaaggaccgc	gcacctggtg	catgaccgc	aagcccggtg	cctgacgcc			2100
gccccacgac	ccgcagcgcc	cgaccgaaag	gagcgcacga	ccccatggct	ccgaccgaag			2160
ccgaccggg	cgccccgcc	gaccccgac	ccgccccga	ggcccaccga	ctctagagtc			2220
ggggcgccg	gccgcttcga	gcagacatga	taagatacat	tgatgagttt	ggacaaacca			2280
caactagaat	gcagtgaaaa	aaatgcttta	tttgtgaaat	ttgtgatgct	attgctttat			2340
ttgtaacat	tataagctgc	aataaacaag	ttaacaacaa	caattgcatt	cattttatgt			2400
ttcaggttca	gggggaggtg	tgaggaggtt	tttaaagcaa	gtaaaacctc	tacaaatgtg			2460
gtaaaatcga	taaggatcaa	ttcggcttca	ggtaccgtcg	acgatgtagg	tcacggctctc			2520
gaagccgcg	tgccgggtgcc	aggcgctgcc	cttgggctcc	ccgggcgcgt	actccacctc			2580
acccatctgg	tccatcatga	tgaacgggtc	gaggtggcgg	tagttgatcc	cggcgaacgc			2640
gcggcgacc	gggaagccct	cgccctcgaa	accgctgggc	gcggtggtca	cggtagcac			2700
gggacgtgc	acggcgctcg	cggtgcgga	tacgcggggc	agcgtcagcg	ggttctcgac			2760
ggtcacggcg	ggcatgtcga	cagccgaatt	gatccgtcga	ccgatgccct	tgagagcctt			2820
caaccagtc	agctccttcc	ggtgggcgcg	gggcatgact	atcgtcgccg	cacttatgac			2880
tgtcttcttt	atcatgcaac	tcgtaggaca	ggtgccggca	gcgctcttcc	gcttctcgc			2940
tactgactc	gctgcgctcg	gtcgctcggc	tgccgcgagc	ggtatcagct	cactcaaagg			3000
cggtaatacg	gttatccaca	gaatcagggg	ataacgcagg	aaagaacatg	aagggcgaat			3060
tctgcagata	tccatcacac	tgccggccgc	tcgagcatgc	atctagaggg	cccaattcgc			3120
cctatagtga	gtcgtattac	aattcactgg	ccgtcgtttt	acaacgtcgt	gactgggaaa			3180
accctggcgt	tacccaactt	aatcgcttg	cagcacatcc	cccttctgcc	agctggcgta			3240
atagcgaaga	ggcccgacc	gategccctt	cccaacagtt	gcgcagccta	tacgtacggc			3300
agtttaaggt	ttacacctat	aaaagagaga	gccgttatcg	tctgtttgtg	gatgtacaga			3360
gtgatattat	tgacacgccg	gggcgacgga	tggtgatccc	cctggccagt	gcacgtctgc			3420
gtgcagataa	agtctcccg	gaactttacc	cggtggtgca	tatcggggat	gaaagctggc			3480
gcatgatgac	caccgatatg	gccagtgtgc	cggtctccgt	tatcggggaa	gaagtggctg			3540
atctcagcca	ccgcgaaaat	gacatcaaaa	acgccattaa	cctgatgttc	tggggaatat			3600

025CIP7SEQ List.tXt

aaatgtcagg	catgagatta	tcaaaaagga	tcttcaccta	gatccttttc	acgtagaaag	3660
ccagtccgca	gaaacggtgc	tgaccccgga	tgaatgtcag	ctactgggct	atctggacaa	3720
gggaaaacgc	aagcgcaaag	agaaagcagg	tagcttgcag	tgggcttaca	tggcgatagc	3780
tagactgggc	ggttttatgg	acagcaagcg	aaccggaatt	gccagctggg	gcgcctctcg	3840
gtaaggttgg	gaagccctgc	aaagtaaact	ggatggcttt	ctcgccgcca	aggatctgat	3900
ggcgcagggg	atcaagctct	gatcaagaga	caggatgagg	atcgtttcgc	atgattgaac	3960
aagatggatt	gcacgcaggt	tctccggcgc	cttgggtgga	gaggctattc	ggctatgact	4020
gggcacaaca	gacaatcggc	tgctctgatg	ccgccgtgtt	ccggctgtca	gcgcaggggc	4080
gcccggttct	ttttgtcaag	accgacctgt	ccggtgccct	gaatgaactg	caagacgagg	4140
cagcgcggct	atcgtggctg	gccacgacgg	gcgttccttg	cgcagctgtg	ctcgacgttg	4200
tactgaagc	gggaaggac	tggtctctat	tgggcgaagt	gccggggcag	gatctcctgt	4260
catctcacct	tgctcctgcc	gagaaagtat	ccatcatggc	tgatgcaatg	cggcggctgc	4320
atacgcttga	tccggctacc	tgcccattcg	accaccaagc	gaaacatcgc	atcgagcgag	4380
cacgtactcg	gatggaagcc	ggtcttgctg	atcaggatga	tctggacgaa	gagcatcagg	4440
ggctcgcgcc	agccgaactg	ttcgccaggc	tcaaggcgag	catgcccgcg	ggcgaggatc	4500
tcgtcgtgac	ccatggcgat	gcctgcttgc	cgaatatcat	ggtggaaaat	ggccgctttt	4560
ctggattcat	cgactgtggc	cggtcgggtg	tggcggaccg	ctatcaggac	atagcgttgg	4620
ctaccctgta	tattgctgaa	gagcttggcg	gcgaatgggc	tgaccgcttc	ctcgtgcttt	4680
acggtatcgc	cgctcccgat	tcgcagcgca	tcgccttcta	tcgccttctt	gacgagttct	4740
tctgaattat	taacgcttac	aatttcctga	tgcggtatth	tctccttacg	catctgtgcg	4800
gtatttcaca	ccgcatacag	gtggcacttt	tcggggaaat	gtgcgcggaa	cccctatttg	4860
tttatttttc	taaatacatt	caaatatgta	tccgctcatg	agacaataac	cctgataaat	4920
gcttcaataa	tagcacgtga	ggagggccac	catggccaag	ttgaccagtg	ccgttcgggt	4980
gctcaccgcg	cgcgacgtcg	ccggagcggg	cgagttctgg	accgaccggc	tcgggttctc	5040
ccgggacttc	gtggaggacg	acttcgcggg	tgtggtccgg	gacgacgtga	ccctgttcat	5100
cagcgcggtc	caggaccagg	tggtgccgga	caacaccctg	gcctgggtgt	gggtgcgcgg	5160
cctggacgag	ctgtacgccg	agtggtcgga	ggtcgtgtcc	acgaacttcc	gggacgcctc	5220
cgggccggcc	atgaccgaga	tcggcgagca	gccgtggggg	cgggagttcg	ccctgcgcga	5280
cccgccgggc	aactgcgtgc	acttcgtggc	cgaggagcag	gactgacacg	tgctaaaact	5340
tcatthttta	tttaaaagga	tctaggtgaa	gatcctthtt	gataatctca	tgacccaaat	5400
cccttaacgt	gagthtttcg	tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	5460

025CIP7SEQ List.tXt

ttcttgagat	ccttttttttc	tgcgcgtaat	ctgctgcttg	caaacaaaaa	aaccaccgct	5520
accagcgggtg	gtttgtttgc	cggatcaaga	gctaccaact	ctttttccga	aggtaactgg	5580
cttcagcaga	gcgagatac	caaatactgt	ccttctagt	tagccgtagt	taggccacca	5640
cttcaagaac	tctgtagcac	cgcctacata	cctcgctctg	ctaatectgt	taccagtggc	5700
tgctgccagt	ggcgataagt	cgtgtcttac	cgggttgac	tcaagacgat	agttaccgga	5760
taaggcgcag	cggtcgggct	gaacgggggg	ttcgtgcaca	cagcccagct	tggagcgaac	5820
gacctacacc	gaactgagat	acctacagcg	tgagctatga	gaaagcgcca	cgtttcccga	5880
agggagaaag	gcgagacaggt	atccggtaag	cggcagggtc	ggaacaggag	agcgacagag	5940
ggagcttcca	gggggaaacg	cctggatatct	ttatagtcct	gtcgggtttc	gccacctctg	6000
acttgagcgt	cgattttttgt	gatgctcgtc	aggggggcgg	agcctatgga	aaaacgccag	6060
caacgcggcc	tttttacgggt	tcctgggctt	ttgctggcct	tttgctcaca	tgttctttcc	6120
tcggttatcc	cctgattctg	tgataaccg	tattaccgcc	tttgagttag	ctgataccgc	6180
tcgccgcagc	cgaacgaccg	agcgacagca	gtcagtgagc	gaggaagcgg	aag	6233

<210> 11
 <211> 234
 <212> DNA
 <213> Artificial sequence

<220>
 <223> attP containing polynucleotide

<400> 11	
gactagtact	gacggacaca ccgaagcccc ggcggaacc ctcagcggat gccccggggc 60
ttcacgtttt	cccaggtcag aagcggtttt cgggagtagt gcccactg gggtaacctt 120
tgagttctct	cagttggggg cgtagggctg ccgacatgac acaaggggtt gtgaccgggg 180
tggacacgta	cgcgggtgct taacgacgct agtcgcgcga gcgcgactag taca 234

<210> 12
 <211> 26
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer attB-for

<400> 12	
taccgtcgac	gatgtaggtc acggtc 26

<210> 13
 <211> 11
 <212> PRT
 <213> SV40

<400> 13

025CIP7SEQ List.tXt
 cys Gly Gly Pro Lys Lys Lys Arg Lys Val Gly
 1 5 10

<210> 14
 <211> 237
 <212> DNA
 <213> mouse

<400> 14
 gacctggaat atcgcgagta aactgaaaat cacggaaaat gagaaataca cacttttagga 60
 cgtgaaatat ggcgaggaaa actgaaaaag gtggaaaatt tagaaatgtc cactgttagga 120
 cgtggaatat ggcaagaaaa ctgaaaatca tggaaaatga gaaacatcca cttgacgact 180
 tgaaaaatga cgaaatcact aaaaaacgtg aaaaatgaga aatgcacact gaaggac 237

<210> 15
 <211> 275
 <212> DNA
 <213> mouse

<400> 15
 gagtgagtta cactgaaaaa cacatacgtt ggaaaccggc attgtagaac agtgtatatc 60
 aatgagttac aatgagaaaa atggaaaatg ataaaaacca cagtgtagaa catattagat 120
 gtgtgagtta cactgaaaaa cacatttcctt ggaaacggga tttgtagaac tgtgtatatc 180
 aatgagttac aatgagaaac atggaaaatg ataaaaacca cactgtagaa catttttagat 240
 gagtgagtta cactgaaaaa cacatatgtt ggaaa 275

<210> 16
 <211> 9299
 <212> DNA
 <213> mouse

<400> 16
 ggccgctctt ctcgttctgc cagcgggccc tcgtctctcc accccatccg tctgcccgtg 60
 gtgtgtggaa ggcaggggtg cggtctctcc gcccgacgct gccccgcgcg cactttttctc 120
 agtggttcgc gtggtccttg tggatgtgtg aggcgcccggt ttgtgccctc acgtgtttca 180
 ctttggtcgt gtctcgcttg accatgttcc cagagtcggt ggatgtggcc ggtggcggtg 240
 catacccttc cgcgtcgttg tgtgcacgcg ctgtttcttg taagcgtcga ggtgctcctg 300
 gagcgttcca ggtttgtctc ctaggtgcct gcttctgagc tgggtggtggc gctccccatt 360
 ccctggtgtg cctccggtgc tccgtctggc tgtgtgcctt cccgtttgtg tctgagaagc 420
 ccgtgagagg ggggtcagag agagaaggag gggcaagacc ccccttcttc gtcgggtgag 480
 gcgcccaccc cgcgactagt acgcctgtgc gtagggtgtg tgcctgagcg tgcgggtgtg 540
 ggttgaaaag tttctcgaga gactcattgc tttcccgtgg ggagctttga gaggcctggc 600
 tttcgggggg gaccggttgc agggctctcc ctgtccgcgg atgctcagaa tgcccttgga 660

O25CIP7SEQ List.txt

agagaacctt	cctgttgccg	cagaccccc	cgcgcggtcg	cccgcggtgtt	ggtcttctgg	720
tttccctgtg	tgctcgtcgc	atgcatactc	tctcggtggc	cggggctcgt	cggggttttg	780
ggtccgtccc	gccctcagtg	agaaagtttc	cttctctagc	tatcttccgg	aaagggtgcg	840
ggcttcttac	ggtctcgagg	ggtctctccc	gaatggctcc	ctggagggct	cgtccctga	900
ccgcctcccc	cgcgcgcagc	gtttgctctc	tcgtctaccg	cggcccgcgg	cctccccgct	960
ccgagttcgg	ggagggatca	cgcggggcag	agcctgtctg	tcgtcctgcc	gttgetgcgg	1020
agcatgtggc	tcggcttgtg	tggttggtgg	ctggggagag	ggctccgtgc	acacccccgc	1080
gtgcgcgtac	tttccctccc	tcctgagggc	cgcctgcggg	acgggggtgtg	ggtaggcgac	1140
ggtgggctcc	cgggtcccca	cccgctcttc	cgtgcctcac	ccgtgccttc	cgtcgcgtgc	1200
gtccctctcg	ctcgcgtcca	cgaacttggtc	cgtcccgccg	acggcgccct	gcgcgcgcgc	1260
tgggtgcgtgc	tgtgtgcttc	tcgggctgtg	tggttgtgtc	gcctcgcccc	ccccttcccg	1320
cggcagcgtt	cccacggctg	gcgaaatcgc	gggagtcctc	cttccccctc	tcggggctga	1380
gagggctcgt	gtctggcgtt	gattgatctc	gctctcgggg	acgggaccgt	tctgtgggag	1440
aacggctgtt	ggccgcgtcc	ggcgcgacgt	cggacgtggg	gacccactgc	cgtcgggggg	1500
tcttcgtcgg	taggcatacg	tgtgtcgcca	tcggctctctc	tctcgtgtcg	gtgtcgcctc	1560
ctcgggctcc	cggggggccg	tcgtgtttcg	ggtcggctcg	gcgtgcagg	tgtggtggga	1620
ctgctcaggg	gagtgggtgca	gtgtgattcc	cgcgggtttt	gcctcgcgtg	ccctgaccgg	1680
tccgacgccc	gagcgggtctc	tcgggtccctt	gtgaggaccc	ccttccggga	ggggcccgtt	1740
tcggccgccc	ttgccgtcgt	cgcgcggccct	cgttctgctg	tgtcgttccc	ccctccccgc	1800
tcgccgcagc	cggctctttt	tcctctctcc	ccccctctcc	tctgactgac	ccgtggccgt	1860
gctgtcggac	cccccgcatg	ggggcgggccg	ggcacgtacg	cgtccggggcg	gtcaccgggg	1920
tcttgggggg	gggcccaggg	gtaagaaagt	cggctcggcg	ggcgggagga	gctgtggttt	1980
ggagggcgctc	ccggccccgc	ggcgtggcg	gtgtcttgcg	cggctcttga	gagggctgcg	2040
tgcgagggga	aaaggttgcc	ccgcgagggc	aaagggaaag	aggctagcag	tggtcattgt	2100
cccgcgggtg	tgggtggtctg	ttggccgagg	tgcgtctggg	gggctcgtcc	ggccctgtcg	2160
tccgtcggga	aggcgcgtgt	tggggcctgc	cggagtgccg	aggtaggtac	cctggcgggtg	2220
ggattaaccc	cgcgcgcgtg	tcccgggtgtg	gcgggtggggg	ctccggctga	tgtctacctc	2280
cctctccccg	aggctctcagg	ccttctccgc	gcgggctctc	ggccctcccc	tcgttctctc	2340
ctctcgcggg	gttcaagtcg	ctcgtcgacc	tccccctctc	cgtccttcca	tctctcgcgc	2400
aatggcgccg	cccgagttca	cggtaggttc	gtcctccgcc	tccgcttctc	gccgggggct	2460
ggccgctgtc	cggctctctcc	tgcccagacc	ccgttgccgt	ggtcttctct	cgcgggcttc	2520

025CIP7SEQ List.tXt						
gcggactcct	ggcttcgccc	ggaggggtcag	ggggcttccc	ggttccccga	cgttgcgcct	2580
cgctgctgtg	tgcttggggg	gggcccgtg	cggcctccgc	ccgcccgtga	gcccctgccg	2640
caccgcgcgg	tgtgcggttt	cgcgcgcgg	tcagttgggc	cctggcgttg	tgtcgcgtcg	2700
ggagcgtgtc	cgctcgcgg	cggttagacg	cggtgtgcgc	cggtctccga	cggttgccct	2760
atccagggct	cgccccgcc	gacccccgcc	tgcccgtccc	ggtggtggtc	gttggtgtgg	2820
ggagtgaatg	gtgctaccgg	tcattccctc	ccgcgtgggt	tgactgtctc	gccggtgtcg	2880
cgcttctctt	tccgccaacc	cccacgcaa	cccaccaccc	tgctctcccg	gcccggtgcg	2940
gtcgacgttc	cggtctccc	gatgccgagg	ggttcgggat	ttgtgccggg	gacggagggg	3000
agagcgggta	agagaggtgt	cggagagctg	tcccggggcg	acgctcgggt	tggctttgcc	3060
gcgtgcgtgt	gctcgcggac	gggttttgtc	ggacccccgac	gggtcggtc	cgccgcgatg	3120
cactctcccg	ttccgcgcga	gcgcccgcgc	ggctcacccc	cggtttgtcc	tcccgcgagg	3180
ctctccgcgg	ccgcgcctc	ctctctctct	ctcgcgtctc	ctgtcccgcc	tggtcctgtc	3240
ccacccccga	cgctccgctc	gcgcttcctt	acctgggtga	tcctgccagg	tagcatatgc	3300
ttgtctcaaa	gattaagcca	tgcatgtcta	agtacgcacg	gccggtacag	tgaaactgcg	3360
aatggctcat	taaatcagtt	atggttcctt	tggtcgctcg	ctcctctcct	acttgataa	3420
ctgtggtaat	tctagagcta	atacatgccg	acgggcgcgtg	accccccttc	ccgggggggg	3480
atgcgtgcat	ttatcagatc	aaaaccaacc	cggtgagctc	cctcccgggt	ccggccgggg	3540
gtcgggcgcc	ggcggcttgg	tgactctaga	taacctcggg	ccgatcgac	gccccccgtg	3600
gcggcgacga	cccattcgaa	cgtctgccct	atcaactttc	gatggtagtc	gccgtgccta	3660
ccatggtgac	cacgggtgac	ggggaatcag	ggttcgattc	cggagaggga	gcctgagaaa	3720
cggctaccac	atccaaggaa	ggcagcaggc	gcgcaaatta	cccactcccg	accgggggag	3780
gtagtgacga	aaaataacaa	tacaggactc	tttcgaggcc	ctgtaattgg	aatgagtcca	3840
ctttaaatcc	tttaacgagg	atccattgga	gggcaagtct	ggtgccagca	gccgcggtaa	3900
ttccagctcc	aatagcgtat	attaaagttg	ctgcagttaa	aaagctcgta	gttgatctt	3960
gggagcgggc	gggcgggtccg	ccgcgaggcg	agtcaccgcc	cgtccccgcc	cettgcctct	4020
cggcgccccc	tcgatgctct	tagctgagtg	tcccgcgggg	cccgaagcgt	ttactttgaa	4080
aaaattagag	tgttcaaagc	aggcccagac	cgcttgata	ccgcagctag	gaataatgga	4140
ataggaccgc	ggttctatct	tggtggtttt	cggaactgag	gccatgatta	agagggacgg	4200
ccgggggcat	tcgtattgcg	ccgctagagg	tgaaattctt	ggaccggcgc	aagacggacc	4260
agagcgaaag	catttgccaa	gaatgttttc	attaatcaag	aacgaaagtc	ggaggttcga	4320
agacgatcag	ataccgtcgt	agttccgacc	ataaacgatg	ccgactggcg	atgcggcggc	4380
gttattccca	tgacccgcgg	ggcagcttcc	gggaaaccaa	agtctttggg	ttccgggggg	4440

025CIP7SEQ List.tXt

agtatggttg	caaagctgaa	acttaaagga	attgacggaa	gggcaccacc	aggagtgggc	4500
ctgcggctta	atttgactca	acacgggaaa	cctcaccg	cccggacacg	gacaggattg	4560
acagattgat	agctctttct	cgattccgtg	ggtggtggtg	catggccgtt	cttagttggt	4620
ggagcgattt	gtctggttaa	ttccgataac	gaacgagact	ctggcatgct	aactagttac	4680
gcgacccccg	agcggtcggc	gtcccccaac	ttcttagagg	gacaagtggc	gttcagccac	4740
ccgagattga	gcaataacag	gtctgtgatg	cccttagatg	tccggggctg	cacgcgcgt	4800
acactgactg	gctcagcgtg	tgctaccct	gcgcggcag	gcgcgggtaa	ccggttgaac	4860
cccattcgtg	atggggatcg	gggattgcaa	ttattcccca	tgaacgagga	attcccagta	4920
agtgcgggtc	ataagcttgc	gttgattaa	tccctgccct	ttgtacacac	cgcccgtcgc	4980
tactaccgat	tggatggtt	agtgaggccc	tggatcggc	ccgcgcggg	tcgcccacg	5040
gccctggcgg	agcgtgaga	agacggtcga	acttgactat	ctagaggaa	taaaagtcgt	5100
aacaaggtt	ccgtaggtga	acctgcggaa	ggatcattaa	acgggagact	gtggaggagc	5160
ggcggcgtgg	cccgtctctc	ccgtcttgtg	tgtgtcctcg	ccgggaggcg	cgtgcgtccc	5220
gggtcccgtc	gcccgcgtgt	ggagcgaggt	gtctggagtg	aggtgagaga	aggggtgggt	5280
gggtcggtc	tgggtccgtc	tgggaccgcc	tccgatttcc	cctccccctc	ccctctccct	5340
cgtccggtc	tgacctcgcc	accctaccgc	ggcggcggct	gctcgcgggc	gtcttgcttc	5400
tttcccgtcc	ggctcttccg	tgtctacgag	ggcggttacg	tcgttacggg	tttttgacct	5460
gtcccggggg	cgttcggctg	tggggcgcg	cgttttctgc	tcccggcacc	catccccgcc	5520
gcggtctctg	cttttctacg	ttggctgggg	cgtttgtcgc	gtgtgggggg	atgtgagtgt	5580
cgcgtgtggg	ctcgcccgtc	ccgatgccac	gcttttcttg	cctcgcgtgt	cctccccgct	5640
cctgtcccgg	gtacctagct	gtcgcgttcc	ggcgcgagg	tttaaggacc	ccgggggggt	5700
cgcctgccg	ccccaggggt	cgggggcgcg	tggggcccg	agggaagtcg	gtcgttcggg	5760
cggctctccc	tcagactcca	tgacctctct	ccccccgctg	ccgccgttcc	cgaggcggcg	5820
gtcgtgtggg	ggggtggatg	tctggagccc	cctcgggcgc	cgtggggggc	cgaccgcgc	5880
cgcggcgttg	cccgatttcc	gcgggtcggt	cctgtcgggtg	ccggtcgtgg	gttcccgtgt	5940
cgttcccgtg	tttttccgct	cccgaccctt	tttttttcc	ccccccaca	cgtgtctcgt	6000
ttcgttcctg	ctggccggcc	tgaggctacc	cctcggtcca	tctgttctcc	tctctctccg	6060
gggagaggag	ggcgggtggtc	gttgggggac	tgtgccgtcg	tcagcaccgc	tgagttcgt	6120
cacacccgaa	ataccgatac	gactcttagc	ggtggatcac	tcggctcgtg	cgtcgatgaa	6180
gaacgcagct	agctgcgaga	attaatgtga	attgcaggac	acattgatca	tcgacacttc	6240
gaacgcactt	gcggcccccg	gttcctccc	gggctacgcc	tgtctgagcg	tcggttgacg	6300

025CIP7SEQ List.tXt

atcaatcgcg	tcaccgcgtg	cggtgggtgc	tgcgcggctg	ggagtttgct	cgcagggcca	6360
accccccaac	cggggtcggg	ccctcegtct	cccgaagtgc	agacgtgtgg	gcggttgtcg	6420
gtgtggcgcg	cgcgcgcgcg	tcgcggagcc	tgggtctccc	cgcgcacccg	cgctcgcggc	6480
ttcttcccg	tccgccgttc	ccgccctcgc	ccgtgcaccc	cggtcctggc	ctcgcgtcgg	6540
cgctcccg	accgctgcct	caccagtctt	tctcggtccc	gtgcccctg	ggaaccacc	6600
gcgccccgt	ggcgcccg	ggtgggcgcg	tccgcacctg	ctctggtcga	ggttggcggt	6660
tgagggtgtg	cgtgcgcgca	ggtggtggtc	ggtcccctgc	ggccgcgggg	ttgtcgggggt	6720
ggcggtcgac	gagggccggg	cggtcgcctg	cggtggttgt	ctgtgtgtgt	ttgggtcttg	6780
cgctggggga	ggcggggtcg	accgctcgcg	gggttggcgc	ggtcgcgcgg	cgcgcgcac	6840
cctccggctt	gtgtggagg	agagcgagg	cgagaacgga	gagaggtggt	atccccgggtg	6900
gcgttgcgag	ggagggtttg	gcgtcccg	tccgtccgtc	cctccctccc	tcgggtggcg	6960
ccttcgcgc	gcacgcggcc	gctagggcg	gtcggggccc	gtggccccc	tggtctcttct	7020
tcgtctccg	ttctcttca	ccggggcggt	accgctccg	gcgcgcggcc	gcgggacgcc	7080
gcggcgctcg	tgcgccgatg	cgagtcaccc	ccgggtgttg	cgagttcggg	gagggagagg	7140
gcctcgcgtg	ccggttgctg	cccggtcttc	ctggggggga	cccggcgtct	gtgggctgtg	7200
cgtcccggg	ggtgcgtgtg	agtaagatcc	tccacccccg	ccgccctccc	ctccgcgcgg	7260
cctctcgggg	acccctgag	acggttcgcc	ggctcgtcct	cccgtgcgcg	cgggtgcctg	7320
ctctttcccg	ccgcctcct	cgtctctctt	ttcccgcggc	tgggcgcgtg	tcccccttt	7380
ctgaccgcga	cctcagatca	gacgtggcga	cccgtggaat	ttaagcatat	tagtcagcgg	7440
aggaagaa	actaaccagg	attccctcag	taacggcgag	tgaacaggga	agagcccagc	7500
gccgaatccc	cgcgcgcgtg	cgcggcggtg	gaaatgtggc	gtacggaaga	ccactcccc	7560
ggcgccgctc	gtggggggcc	caagtccctc	tgatcgaggc	ccagcccgtg	gacggtgtga	7620
ggccggtagc	ggccccggcg	cgcggggctc	gggtcttccc	ggagtcgggt	tgcttgggaa	7680
tgcagcccaa	agcgggtggt	aaactccatc	taaggctaaa	taccggcacg	agaccgatag	7740
tcaacaagta	ccgtaaggga	aagttgaaaa	gaactttgaa	gagagagttc	aagagggcgt	7800
gaaaccgtta	agaggtaaac	gggtgggggtc	cgcgcagtc	gcccggagga	ttcaaccgcg	7860
cggcgcgctg	ccggccgtgc	ccggtgggtc	cggcgatct	ttcccgctcc	ccgttccctc	7920
cgaccctcc	accgcgcgtg	cgttccctc	ttctccccc	cgtccggcgc	ctccggcggc	7980
gggcgcgggg	ggtggtgtgg	tgggtggcg	cgggcggggc	cgggggtggg	gtcggcgggg	8040
gaccgcccc	ggcggcgac	cggccgcgc	cgggcgcact	tccaccgtgg	cgggtgcgcg	8100
cgaccggtc	cgggacggcc	gggaaggccc	ggtggggaag	gtggctcggg	gggggcggcg	8160
cgtctcagg	cgcgccgaac	cacctcacc	cgagtgttac	agccctccgg	ccgcgctttc	8220

025CIP7SEQ List.tXt

gccgaatccc	ggggccgagg	aagccagata	cccgtcgccg	cgtctctccct	ctccccccgt	8280
ccgcctcccg	ggcgggctg	ggggtggggg	ccgggccgcc	cctcccacgg	cgcgaccgct	8340
ctcccacccc	cctccgtcgc	ctctctcggg	gcccgtggg	gggcggggcg	gactgtcccc	8400
agtgcgcccc	gggcgtcgtc	gcgcgcgtcg	gtcccggggg	gaccgtcggg	cacgcgtctc	8460
ccgacgaagc	cgagcgcacg	gggtcggcgg	cgatgtcggc	taccaccccg	accggtcttg	8520
aaacacggac	caaggagtct	aacgcgtgcg	cgagtcaggg	gctcgtccga	aagccgcggt	8580
ggcgcaatga	aggtgaaggg	ccccgccggg	ggggccgagg	tgggatcccg	aggcctctcc	8640
agtccgccga	gggcgcacca	ccggcccgtc	tcgcccgcgg	cgccggggag	gtggagcacg	8700
agcgtacgcg	ttaggacccg	aaagatgggt	aactatgctt	gggcagggcg	aagccagagg	8760
aaactctggt	ggaggtccgt	agcggtcctg	acgtgcaa	cggtcgtccg	acctgggtat	8820
aggggcgaaa	gactaatcga	accatctagt	agctgggtcc	ctccgaagtt	tccttcagga	8880
tagctggcgc	tctcgtcccc	gacgtacgca	gttttatccg	gtaaagcgaa	tgattagagg	8940
tcttggggcc	gaaacgatct	caacctattc	tcaaacttta	aatgggtaag	aagcccggct	9000
cgtcggcgtg	gagccggggc	tggaatgcga	gtgcctagt	ggccactttt	ggtaagcaga	9060
actggcgtg	cgggatgaac	cgaacgcggg	gttaaggcgc	ccgatgccga	cgtcatcag	9120
acccagaaa	aggtgttggg	tgatatagac	agcaggacgg	tggccatgga	agtcgggaatc	9180
cgctaaggag	tgtgtaacaa	ctcacctgcc	gaatcaacta	gccctgaaaa	tggatggcgc	9240
tggagcgtcg	ggcccatacc	cggcgcgtcg	cgcagtcgga	acggaacggg	acgggagcg	9299

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2006/006752**A. CLASSIFICATION OF SUBJECT MATTER***C12N 15/11(2006.01)I, C12N 15/06(2006.01)I, C12N 15/00(2006.01)I*

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC8 C12N 15/1 I, A61K 67/027

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

NCBI Pubmed, Esp@snet, Delphion

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X Y	US 2004/255345 A1 (RAPP, J C) 16 DECEMBER 2004 See the abstract, pages 2, 12, and 18	1, 5-17, 19-24 2-4, 18
Y	US 2005/003414 A1 (ALEX, H) 6 JANUARY 2005 See the abstract and pages 1-2	2-4, 18
Y	DE JONG, G et al 'Mammalian artificial chromosome pilot production facility large-scale isolation of functional satellite DNA-based artificial chromosomes' Cytometry, Vol 35, no 2, pp 129-133 1 FEBRUARY 1999 See the whole document	2-4, 18



Further documents are listed in the continuation of Box C



See patent family annex

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

07 JULY 2006 (07 07 2006)

Date of mailing of the international search report

10 JULY 2006 (10.07.2006)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No 82-42-472-7140

Authorized officer

KIN, Ji Yun

Telephone No 82-42-481-8288



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/006752

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2004/255345 A1	2004-12-16	EP 01608219 A2	2005-12-28
		US 2004226057 A1	2004-11-11
		WO 2004080162	2004-9-23
US 2005/003414 A1	2005-1-6	EP 01608219 A2	2005-12-28
		US 20040210954 A1	2004 10 26
		WO 2005040215 A2	2005-5-6